

1 Generalized determinant

Alexander Barvinok [1] defines a symmetrized determinant of a matrix nxn over a ring (which need not be commutative) as:

$$\text{sdet}_n(X) = \frac{1}{n!} \cdot \sum_{\sigma, \tau \in S_n} \text{sgn}(\sigma) \cdot \text{sgn}(\tau) \cdot \prod_{i=1}^n x_{\tau(i), \sigma(i)}$$

I was interested in its properties, so I examined the definition as follows:

$$\begin{aligned} \text{sdet}_n(X) &= \frac{1}{n!} \cdot \sum_{\sigma, \tau \in S_n} \text{sgn}(\sigma) \cdot \text{sgn}(\tau) \cdot \prod_{i=1}^n x_{\tau(i), \sigma(i)} = \\ &= \frac{1}{n!} \cdot \sum_{\sigma, \tau \in S_n} \text{sgn}(\sigma) \cdot \text{sgn}(\tau) \cdot \prod_{i=1}^n x_{\tau(i), (\sigma \circ (\tau^{-1} \circ \tau))(i)} = \\ &= \frac{1}{n!} \cdot \sum_{\tau \in S_n} \text{sgn}(\tau) \cdot \sum_{\sigma \in S_n} \text{sgn}(\sigma) \cdot \prod_{i=1}^n x_{\tau(i), ((\sigma \circ \tau^{-1}) \circ \tau)(i)} = \\ &= \frac{1}{n!} \cdot \sum_{\tau \in S_n} \text{sgn}(\tau) \cdot \sum_{\pi \in (S_n \tau^{-1})} \text{sgn}(\pi \circ \tau) \cdot \prod_{i=1}^n x_{\tau(i), (\pi \circ \tau)(i)} = \\ &= \frac{1}{n!} \cdot \sum_{\tau \in S_n} \text{sgn}(\tau) \cdot \sum_{\pi \in S_n} \text{sgn}(\pi) \cdot \text{sgn}(\tau) \cdot \prod_{i=1}^n x_{\tau(i), (\pi \circ \tau)(i)} = \\ &= \frac{1}{n!} \cdot \sum_{\tau \in S_n} \sum_{\pi \in S_n} \text{sgn}(\pi) \cdot \prod_{i=1}^n x_{\tau(i), \pi(\tau(i))} = \\ &= \sum_{\pi \in S_n} \text{sgn}(\pi) \cdot \frac{1}{n!} \cdot \sum_{\tau \in S_n} \prod_{i=1}^n x_{\tau(i), \pi(\tau(i))} = \\ &= \sum_{\pi \in S_n} \text{sgn}(\pi) \cdot \prod_{i=1}^n (x_{1, \pi(1)}, x_{2, \pi(2)}, \dots, x_{n, \pi(n)}) \end{aligned}$$

The average product is defined as the average value of products over all permutations of the factors:

$$\prod_{i=1}^n (a_1, a_2, \dots, a_n) = \frac{1}{n!} \cdot \sum_{\nu \in S_n} a_{\nu(1)} \cdot a_{\nu(2)} \cdot \dots \cdot a_{\nu(n)}$$

2 Comparison

As a result, the only difference between a normal determinant and a symmetrized determinant is the following:

$$\det_n(X) = \sum_{\pi \in S_n} \text{sgn}(\pi) \cdot x_{1,\pi(1)} \cdot x_{2,\pi(2)} \cdot \dots \cdot x_{n,\pi(n)}$$
$$\text{sdet}_n(X) = \sum_{\pi \in S_n} \text{sgn}(\pi) \cdot \prod^{\text{ave}}(x_{1,\pi(1)}, x_{2,\pi(2)}, \dots, x_{n,\pi(n)})$$

If the ring were commutative, then $\det_n(X)$ would be equal to $\text{sdet}_n(X)$.

Paper [2] shows that calculating the normal determinant (also called Cayley determinant) over a non-commutative ring is NP-hard, but the symmetrized determinant can be always computed efficiently.

3 References

- [1] A. Barvinok. New Permanent Estimators via Non-Commutative Determinants. Online: <https://arxiv.org/abs/math/0007153>
- [2] V. Arvind, S. Srinivasan. On the hardness of the noncommutative determinant. Online: <https://arxiv.org/abs/0910.2370>