

Computer Science Track Core Course

Linear programming — part 2

Martin Dvorak
IST Austria
Kolmogorov group

2021-06-10

Linear program

Definitions

We are given an input $A \in \mathbb{Q}^{m \times n}$, $b \in \mathbb{Q}^m$, $c \in \mathbb{Q}^n$ and we search for $\mathbf{x} \in \mathbb{Q}^n$ that optimizes...

$$\begin{array}{ll} \text{minimize} & c' \mathbf{x} \\ \text{s.t.} & A\mathbf{x} \geq b \end{array}$$

$$\begin{array}{ll} \text{maximize} & c' \mathbf{x} \\ \text{s.t.} & A\mathbf{x} \leq b \end{array}$$

$$\begin{array}{ll} \text{minimize} & c' \mathbf{x} \\ \text{s.t.} & A\mathbf{x} = b \\ \text{where} & \mathbf{x} \geq \mathbf{0} \end{array}$$

$$\begin{array}{ll} \text{maximize} & c' \mathbf{x} \\ \text{s.t.} & A\mathbf{x} = b \\ \text{where} & \mathbf{x} \geq \mathbf{0} \end{array}$$

Integer linear program

Definitions

We are given an input $A \in \mathbb{Z}^{m \times n}$, $b \in \mathbb{Z}^m$, $c \in \mathbb{Z}^n$ and we search for $\mathbf{x} \in \mathbb{Z}^n$ that optimizes...

$$\begin{array}{ll} \text{minimize} & c' \mathbf{x} \\ \text{s.t.} & A\mathbf{x} \geq b \end{array}$$

$$\begin{array}{ll} \text{maximize} & c' \mathbf{x} \\ \text{s.t.} & A\mathbf{x} \leq b \end{array}$$


$$\begin{array}{ll} \text{minimize} & c' \mathbf{x} \\ \text{s.t.} & A\mathbf{x} = b \\ \text{where} & \mathbf{x} \geq \mathbf{0} \end{array}$$

$$\begin{array}{ll} \text{maximize} & c' \mathbf{x} \\ \text{s.t.} & A\mathbf{x} = b \\ \text{where} & \mathbf{x} \geq \mathbf{0} \end{array}$$

Example of LP

Separating points

We are searching for a line $y = ax + b$ that separates “disks” (top) from “squares” (bottom).



Variables: $a \leq 0, b \leq 0$

Point (x_1, y_1) above the line: $x_1 \cdot a + b \leq y_1$

“disk”

Point (x_2, y_2) below the line: $x_2 \cdot a + b \geq y_2$

“square”

Objective: minimize 0

Example of ILP


Separating points partially

We are searching for a line $y = ax + b$ that separates given points — “disks” (top) from “squares” (bottom) — with as few exceptions as possible.

Let us ignore all vertical and nearly-vertical solutions.

M is a very large integer.

Variables:



Example of ILP


Separating points partially

We are searching for a line $y = ax + b$ that separates given points — “disks” (top) from “squares” (bottom) — with as few exceptions as possible.

Let us ignore all vertical and nearly-vertical solutions.

M is a very large integer.

Variables: $a \leq 0$, $b \leq 0$; $\forall i : p_i \in \{0, 1\}$



Example of ILP

Separating points partially


We are searching for a line $y = ax + b$ that separates given points — “disks” (top) from “squares” (bottom) — with as few exceptions as possible.

Let us ignore all vertical and nearly-vertical solutions.

M is a very large integer.

Variables: $a \leq 0$, $b \leq 0$; $\forall i : p_i \in \{0, 1\}$

Point (x_1, y_1) above the line:



Example of ILP

Separating points partially


We are searching for a line $y = ax + b$ that separates given points — “disks” (top) from “squares” (bottom) — with as few exceptions as possible.

Let us ignore all vertical and nearly-vertical solutions.

M is a very large integer.

Variables: $a \leq 0, b \leq 0; \forall i: p_i \in \{0, 1\}$

Point (x_1, y_1) above the line: $x_1 \cdot a + b - M \cdot p_i \leq y_1$ “disk”



Example of ILP

Separating points partially

We are searching for a line $y = ax + b$ that separates given points — “disks” (top) from “squares” (bottom) — with as few exceptions as possible.


Let us ignore all vertical and nearly-vertical solutions.

M is a very large integer.

Variables: $a \leq 0, b \leq 0; \forall i: p_i \in \{0, 1\}$

Point (x_1, y_1) above the line: $x_1 \cdot a + b - M \cdot p_i \leq y_1$ “disk”

Point (x_2, y_2) below the line:



Example of ILP

Separating points partially

We are searching for a line $y = ax + b$ that separates given points — “disks” (top) from “squares” (bottom) — with as few exceptions as possible.

Let us ignore all vertical and nearly-vertical solutions.

M is a very large integer.


Variables: $a \leq 0, b \leq 0; \forall i: p_i \in \{0, 1\}$

Point (x_1, y_1) above the line: $x_1 \cdot a + b - M \cdot p_i \leq y_1$

“disk”

Point (x_2, y_2) below the line: $x_2 \cdot a + b + M \cdot p_i \geq y_2$

“square”



Example of ILP

Separating points partially

We are searching for a line $y = ax + b$ that separates given points — “disks” (top) from “squares” (bottom) — with as few exceptions as possible.

Let us ignore all vertical and nearly-vertical solutions.

M is a very large integer.

Variables: $a \leq 0, b \leq 0; \forall i: p_i \in \{0, 1\}$


Point (x_1, y_1) above the line: $x_1 \cdot a + b - M \cdot p_i \leq y_1$

“disk”

Point (x_2, y_2) below the line: $x_2 \cdot a + b + M \cdot p_i \geq y_2$

“square”

Objective:



Example of ILP

Separating points partially

We are searching for a line $y = ax + b$ that separates given points — “disks” (top) from “squares” (bottom) — with as few exceptions as possible.

Let us ignore all vertical and nearly-vertical solutions.

M is a very large integer.

Variables: $a \leq 0, b \leq 0; \forall i : p_i \in \{0, 1\}$


Point (x_1, y_1) above the line: $x_1 \cdot a + b - M \cdot p_i \leq y_1$

“disk”

Point (x_2, y_2) below the line: $x_2 \cdot a + b + M \cdot p_i \geq y_2$

“square”

Objective: minimize $\sum_i p_i$



Example of ILP

Separating points partially

Variables: $a \leq 0, b \leq 0; \forall i : p_i \in \{0, 1\}$


Point (x_1, y_1) above the line: $x_1 \cdot a + b - M \cdot p_i \leq y_1$

“disk”

Point (x_2, y_2) below the line: $x_2 \cdot a + b + M \cdot p_i \geq y_2$

“square”

Objective: minimize $\sum_i p_i$



Example of ILP

Separating points partially

Variables: $a \leq 0, b \leq 0; \forall i : p_i \in \{0, 1\}$


Point (x_1, y_1) above the line: $x_1 \cdot a + b - M \cdot p_i \leq y_1$

“disk”

Point (x_2, y_2) below the line: $x_2 \cdot a + b + M \cdot p_i \geq y_2$

“square”

Objective: minimize $\sum_i p_i$



Unfortunately, there is no general algorithm for solving ILP in polynomial time

Example of ILP


Separating points partially

Variables: $a \leq 0, b \leq 0; \forall i : p_i \in \{0, 1\}$

Point (x_1, y_1) above the line: $x_1 \cdot a + b - M \cdot p_i \leq y_1$ “disk”

Point (x_2, y_2) below the line: $x_2 \cdot a + b + M \cdot p_i \geq y_2$ “square”

Objective: minimize $\sum_i p_i$



Unfortunately, there is no general algorithm for solving ILP in polynomial time (we don't know any such algorithm — but most importantly, it cannot exist unless $P = NP$).

Example of ILP


Separating points partially

Variables: $a \leq 0, b \leq 0; \forall i : p_i \in \{0, 1\}$

Point (x_1, y_1) **above** the line: $x_1 \cdot a + b - M \cdot p_i \leq y_1$ “disk”

Point (x_2, y_2) **below** the line: $x_2 \cdot a + b + M \cdot p_i \geq y_2$ “square”

Objective: minimize $\sum_i p_i$



Unfortunately, there is no general algorithm for solving ILP in polynomial time (we don't know any such algorithm — but most importantly, it cannot exist unless $P = NP$).

Does it mean, however, that our problem cannot be solved in polynomial time?

Combinations of vectors (linear, affine, convex)

Definition

Let V be a vector space over F . Consider vectors $\mathbf{x}_1, \dots, \mathbf{x}_n \in V$.

Combinations of vectors (linear, affine, convex)

Definition

Let V be a vector space over F . Consider vectors $\mathbf{x}_1, \dots, \mathbf{x}_n \in V$.

- We say that $\mathbf{y} \in V$ is a **linear combination** of $\mathbf{x}_1, \dots, \mathbf{x}_n$ if:

$$\exists \alpha_1, \dots, \alpha_n \in F : \sum_{i=1}^n \alpha_i \mathbf{x}_i = \mathbf{y}$$

Combinations of vectors (linear, affine, convex)

Definition

Let V be a vector space over F . Consider vectors $\mathbf{x}_1, \dots, \mathbf{x}_n \in V$.

- We say that $\mathbf{y} \in V$ is a **linear combination** of $\mathbf{x}_1, \dots, \mathbf{x}_n$ if:

$$\exists \alpha_1, \dots, \alpha_n \in F : \sum_{i=1}^n \alpha_i \mathbf{x}_i = \mathbf{y}$$

- We say that $\mathbf{y} \in V$ is an **affine combination** of $\mathbf{x}_1, \dots, \mathbf{x}_n$ if:

$$\exists \alpha_1, \dots, \alpha_n \in F : \sum_{i=1}^n \alpha_i = 1 \wedge \sum_{i=1}^n \alpha_i \mathbf{x}_i = \mathbf{y}$$

Combinations of vectors (linear, affine, convex)

Definition

Let V be a vector space over F . Consider vectors $\mathbf{x}_1, \dots, \mathbf{x}_n \in V$.

- We say that $\mathbf{y} \in V$ is a **linear combination** of $\mathbf{x}_1, \dots, \mathbf{x}_n$ if:

$$\exists \alpha_1, \dots, \alpha_n \in F : \sum_{i=1}^n \alpha_i \mathbf{x}_i = \mathbf{y}$$

- We say that $\mathbf{y} \in V$ is an **affine combination** of $\mathbf{x}_1, \dots, \mathbf{x}_n$ if:

$$\exists \alpha_1, \dots, \alpha_n \in F : \sum_{i=1}^n \alpha_i = 1 \wedge \sum_{i=1}^n \alpha_i \mathbf{x}_i = \mathbf{y}$$

- Suppose now that F is a totally ordered field.


Denote the set of $\varphi \in F$ such that $\varphi \geq 0$ by the symbol F_0^+ .

We say that $\mathbf{y} \in V$ is a **convex combination** of $\mathbf{x}_1, \dots, \mathbf{x}_n$ if:

$$\exists \alpha_1, \dots, \alpha_n \in F_0^+ : \sum_{i=1}^n \alpha_i = 1 \wedge \sum_{i=1}^n \alpha_i \mathbf{x}_i = \mathbf{y}$$

Combinations of vectors (linear, affine, convex)

Example



Combinations of vectors (linear, affine, convex)

Quiz

Let $d \in \mathbb{Z}$ such that $d \geq 2$.

What is the **maximum** possible amount (largest set) of:

- **Linearly** independent vectors in \mathbb{Z}_2^d over \mathbb{Z}_2 ?
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{C} ?
- **Convexly** independent vectors in \mathbb{Q}^d over \mathbb{Q} ?

Combinations of vectors (linear, affine, convex)

Quiz

Let $d \in \mathbb{Z}$ such that $d \geq 2$.

What is the **maximum** possible amount (largest set) of:

- **Linearly** independent vectors in \mathbb{Z}_2^d over \mathbb{Z}_2 ?
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{C} ?
- **Convexly** independent vectors in \mathbb{Q}^d over \mathbb{Q} ?

Combinations of vectors (linear, affine, convex)

Quiz

Let $d \in \mathbb{Z}$ such that $d \geq 2$.

What is the **maximum** possible amount (largest set) of:

- **Linearly** independent vectors in \mathbb{Z}_2^d over \mathbb{Z}_2 ?
 d
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{C} ?
 $d + 1$
- **Convexly** independent vectors in \mathbb{Q}^d over \mathbb{Q} ?

Combinations of vectors (linear, affine, convex)

Quiz

Let $d \in \mathbb{Z}$ such that $d \geq 2$.

What is the **maximum** possible amount (largest set) of:

- **Linearly** independent vectors in \mathbb{Z}_2^d over \mathbb{Z}_2 ?
 d
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{C} ?
 $d + 1$
- **Convexly** independent vectors in \mathbb{Q}^d over \mathbb{Q} ?
 ∞

Combinations of vectors (linear, affine, convex)

Quiz

Let $d \in \mathbb{Z}$ such that $d \geq 2$.

What is the **maximum** possible amount (largest set) of:

- **Linearly** independent vectors in \mathbb{Z}_2^d over \mathbb{Z}_2 ?
 d
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{C} ?
 $d + 1$
- **Convexly** independent vectors in \mathbb{Q}^d over \mathbb{Q} ?
 ∞
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{R} ?
- **Linearly** independent vectors in \mathbb{R}^d over \mathbb{Q} ?

Combinations of vectors (linear, affine, convex)

Quiz

Let $d \in \mathbb{Z}$ such that $d \geq 2$.

What is the **maximum** possible amount (largest set) of:

- **Linearly** independent vectors in \mathbb{Z}_2^d over \mathbb{Z}_2 ?
 d
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{C} ?
 $d + 1$
- **Convexly** independent vectors in \mathbb{Q}^d over \mathbb{Q} ?
 ∞
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{R} ?
 $2d + 1$
- **Linearly** independent vectors in \mathbb{R}^d over \mathbb{Q} ?

Combinations of vectors (linear, affine, convex)

Quiz

Let $d \in \mathbb{Z}$ such that $d \geq 2$.

What is the **maximum** possible amount (largest set) of:

- **Linearly** independent vectors in \mathbb{Z}_2^d over \mathbb{Z}_2 ?
 d
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{C} ?
 $d + 1$
- **Convexly** independent vectors in \mathbb{Q}^d over \mathbb{Q} ?
 ∞
- **Affinely** independent vectors in \mathbb{C}^d over \mathbb{R} ?
 $2d + 1$
- **Linearly** independent vectors in \mathbb{R}^d over \mathbb{Q} ?
 ∞

Combinations of vectors (linear, affine, convex)

Basic properties

$$\{\text{convex combin.}\} \subseteq \{\text{affine combin.}\} \subseteq \{\text{linear combin.}\}$$

Combinations of vectors (linear, affine, convex)

Basic properties

$$\{\text{convex combin.}\} \subseteq \{\text{affine combin.}\} \subseteq \{\text{linear combin.}\}$$

Let V be a vector space and $X \subseteq V$.

- X is convexly dependent. $\implies X$ is affinely dependent.
 $\implies X$ is linearly dependent.
- X is linearly independent. $\implies X$ is affinely independent.
 $\implies X$ is convexly independent.

Combinations of vectors (linear, affine, convex)

Basic properties

$$\{\text{convex combin.}\} \subseteq \{\text{affine combin.}\} \subseteq \{\text{linear combin.}\}$$

Let V be a vector space and $X \subseteq V$.

- X is convexly dependent. $\implies X$ is affinely dependent.
 $\implies X$ is linearly dependent.
- X is linearly independent. $\implies X$ is affinely independent.
 $\implies X$ is convexly independent.

Let us have a matrix $A \in F^{m \times n}$ and a vector $b \in F^m$.

We search for a solution $x \in F^n$.

- $Ax = 0$: Any linear combination of solutions is a solution.
- $Ax = b$: Any affine combination of solutions is a solution.
- $Ax \leq b$: Any convex combination of solutions is a solution.

In this example, F must be a totally ordered field.

Convex hull

Definition and exercises

Def. Let V be a vector space over a totally ordered field. Consider a set $X \subseteq V$. We define a convex hull of X as:

$$\text{conv}(X) = \{\mathbf{y} \in V \mid \mathbf{y} \text{ is a convex combin. of some } \mathbf{x}_1, \mathbf{x}_2, \dots \in X\}$$

Convex hull

Definition and exercises

Def. Let V be a vector space over a totally ordered field. Consider a set $X \subseteq V$. We define a convex hull of X as:

$$\text{conv}(X) = \{\mathbf{y} \in V \mid \mathbf{y} \text{ is a convex combin. of some } \mathbf{x}_1, \mathbf{x}_2, \dots \in X\}$$

Ex. Prove this set identity:

$$\text{conv}(\text{conv}(X)) = \text{conv}(X)$$

Convex hull

Definition and exercises

Def. Let V be a vector space over a totally ordered field. Consider a set $X \subseteq V$. We define a convex hull of X as:

$$\text{conv}(X) = \{\mathbf{y} \in V \mid \mathbf{y} \text{ is a convex combin. of some } \mathbf{x}_1, \mathbf{x}_2, \dots \in X\}$$

Ex. Prove this set identity:

$$\text{conv}(\text{conv}(X)) = \text{conv}(X)$$

Ex. Let $v \in V$. If $M \subseteq V$, we define $M + v$ as $\{m + v \mid m \in M\}$.

Prove this set identity:

$$\text{conv}(X) + v = \text{conv}(X + v)$$

Convex hull

Definition and exercises

Def. Let V be a vector space over a totally ordered field. Consider a set $X \subseteq V$. We define a convex hull of X as:

$$\text{conv}(X) = \{\mathbf{y} \in V \mid \mathbf{y} \text{ is a convex combin. of some } \mathbf{x}_1, \mathbf{x}_2, \dots \in X\}$$

Ex. Prove this set identity:

$$\text{conv}(\text{conv}(X)) = \text{conv}(X)$$

Ex. Let $v \in V$. If $M \subseteq V$, we define $M + v$ as $\{m + v \mid m \in M\}$. Prove this set identity:

$$\text{conv}(X) + v = \text{conv}(X + v)$$

Ex. Express a d -dimensional simplex (triangle, tetrahedron,...) as:

- A convex hull of $d + 1$ points.
- An intersection of $d + 1$ closed half-spaces.

Polyhedral vertices

Equivalent formulations

Def. Let $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$ where $m \geq n$. Consider a set $P \subseteq \mathbb{R}^n$ defined by $P = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} \leq b\}$, that is, a polyhedron.

Let $\mathbf{p} \in P$. The following conditions are equivalent formulations of \mathbf{p} being a vertex of the polyhedron P .

- No vector $\mathbf{y} \neq \mathbf{0}$ satisfies both $\mathbf{p} + \mathbf{y} \in P$ and $\mathbf{p} - \mathbf{y} \in P$.
- We have $\mathbf{p} \notin \text{conv}(P \setminus \{\mathbf{p}\})$.
- There exists a hyperplane H of dimension $n - 1$ such that $P \cap H = \{\mathbf{p}\}$.

Recall that $H = \{\mathbf{x} \in \mathbb{R}^n \mid h'\mathbf{x} = r\}$ for some $h \in \mathbb{R}^n$, $r \in \mathbb{R}$.

- There is a cost vector $c \in \mathbb{R}^n$ such that \mathbf{p} is the unique maximum of the corresponding cost function, that is, $\forall \mathbf{x} \in (P \setminus \{\mathbf{p}\})$ we have $c'\mathbf{p} > c'\mathbf{x}$.
- There are n linearly independent constraints tight ($=$) at \mathbf{p} .

Ex. Prove their equivalence.

Thanks for your attention!

Questions?