

Which Stems Will Beaver Harvest? A Final Project for FOR 796.

Abstract

Enter the text of your abstract here.

Introduction

Methods

(Mahoney and Stella 2020)

```
str(tree_data)
```

```
## 'data.frame':  10368 obs. of  15 variables:
## $ Site      : chr  "Alder Pond" "Alder Pond" "Alder Pond" "Alder Pond" ...
## $ Transect  : int   1 1 1 1 1 1 1 1 1 1 ...
## $ Plot      : chr   "A" "A" "A" "A" ...
## $ OU        : chr   "U" "U" "U" "U" ...
## $ Species   : chr   "EH" "AB" "AB" "RM" ...
## $ DBHClass  : chr   "0-1" "0-1" "0-1" "0-1" ...
## $ HeightClass: chr   "0-1" "0-1" "0-1" "0-1" ...
## $ Gnaw      : chr   "0" "0" "0" "0" ...
## $ Stump     : chr   "0" "0" "0" "0" ...
## $ Slope     : int  10 10 10 10 10 10 10 10 10 10 ...
## $ LightDots : int    0 0 0 0 0 0 0 0 0 0 ...
## $ LakeRiver : chr   "Lake" "Lake" "Lake" "Lake" ...
## $ Elevation : int  327 327 327 327 327 327 327 327 327 327 ...
## $ Lat       : num  43.9 43.9 43.9 43.9 43.9 ...
## $ Long      : num -73.7 -73.7 -73.7 -73.7 -73.7 ...
```

Results

Discussion

Mahoney, Michael J., and John C. Stella. 2020. "Stem Size Selectivity Is Stronger Than Species Preferences for Beaver, a Central Place Forager." *Forest Ecology and Management* 475: 118331. <https://doi.org/10.1016/j.foreco.2020.118331>.