

Nothing Binds a Twin but Exclusion: The Prime Clocks, the Exclusion Law and the Recurrence of the Twin Pair

Abhijit Singh

Independent Researcher; abhijitsingh777@gmail.com

Abstract

We read the integers prime to 6 as the positions of a world of clocks. Each prime $p \geq 5$ is a clock that marks the two centres k with $6k \equiv \pm 1 \pmod{p}$. A centre that no clock marks is a full zero, and below the reach of the clocks a full zero is exactly a twin prime pair $6k \pm 1$. The step from one clock to the next builds the new world from $P' - 2$ exact copies of the old one and preserves the count $\prod(p - 2)$, the palindrome and the copy sign $\prod(p - 4) > 0$. The two members of a pair are bound by exactly one relation: a clock may mark one member, and then it does not mark the other. As operators, every clock acts on one member at a time and annihilates its own exchange. This exclusion law fixes the twin constant as the exclusion product $E = \prod_{p \geq 5} (1 - (p - 1)^{-2}) = 0.880215 \dots$, and under the axiom that independence is the absence of a relation it gives the twin law $\pi_2(x) \sim 2C_2 x / \log^2 x$. Every measurement agrees with the exclusion law and with nothing beyond it. These are the joint law of the prime factors over $5 \cdot 10^7$ pairs, the pair symbol, the frame of rough pairs, the count of 27,412,679 twin pairs to 10^{10} (the law to 4.6 parts in 10^5), and the reach of the clocks. They include Maier's windows near 10^9 – 10^{10} , where the twins swing 18% above and 8% below the average and the clocks predict the swing to within 0.05% over 2.65 million pairs. The parity barrier is a ceiling of the instrument: a sieve sees only sums over residue classes, the Liouville sign has none, and distribution information certifies at most 48.837% of the rough members as prime. The same reading organises the lean of single primes and the central values of the congruent-number curves, which are $4\kappa_n m^2$ with $m \in \mathbb{Z}$.

Keywords: twin primes; exclusion; prime clocks; Hardy–Littlewood conjecture; parity barrier; Buchstab function; Liouville function; congruent numbers

1. Introduction: What Binds a Twin

Two odd numbers $6k - 1$ and $6k + 1$ share one thing, their centre $6k$, and differ by 2. The twin prime question asks whether infinitely often both are prime. This paper asks a prior question: *what binds the two members of a pair?* If nothing binds them except what the construction of the integers visibly puts between them, then the twin pair must recur exactly as often as two unbound members would both be prime, corrected by that one visible bond. The work of the paper is to find the bond, to prove that it is the only one, and to measure the pair against it.

The bond is found in Section 4. Each prime $p \geq 5$ acts on the centres as a clock with two marks, one for each member. The two marks are different classes modulo p , so a prime can divide one member or the other but never both. This *exclusion* is the whole of the relation. In the world of clocks it is a theorem that everything else about the pair is the product of its single members (Theorem 4). As operators on the states of the pair, every clock and every motion of the centre acts on one member at a time, and a clock annihilates its own exchange (Theorem 5): the members are two strands that share every centre and are joined by no operator of the construction. The exclusion changes the product by the factor $1 - (p - 1)^{-2}$ at each clock, and their product is the twin constant.

The measurements of Sections 5 and 6 test the pair against the bond and find nothing beyond it: not in the joint law of the prime factors of the two members, not in the generating symbol of that law at any radius, not on the frame of rough pairs, and not in the count of twin pairs to 10^{10} .

Section 7 addresses the reason the question has been held to be closed to the methods that prove the most about primes. It shows that the obstruction, known as the parity barrier, is a ceiling of an instrument. The instrument is information about how primes are spread over residue classes. It cannot tell a prime from a product of two large primes, and when that is measured exactly it certifies 48.837% where a majority is needed. That ceiling describes the instrument. It says nothing about whether the members of a pair are bound.

We build on the work of many. Brun's sieve [1] and the linear sieve with its optimal functions [7] are the instrument whose ceiling Section 7 measures. Hardy and Littlewood conjectured the twin law [10]. Mertens' theorem [17] and Buchstab's function [7] fix the shares in the reach of the clocks. Chen's theorem [2], the bounded gaps of Zhang and of Maynard [16,24] and Tao's logarithmic two-point Chowla theorem [22] are the nearest proved statements. The biases of Lemke Oliver and Soundararajan [14], Chebyshev's bias in the form of Rubinstein and Sarnak [19,20], the partial Euler products of Goldfeld, Conrad, and Kim and Murty [3,9,12], and Tunnell's theorem [23] are what Sections 8 and 9 read. The axiom of Section 6 is the form, for a pair of numbers, of the axiom of harmonic independence stated for the prime waves in [21].

Sections 8 and 9 show that the same reading, a thing and its copy, a pair and its mirror, organises the lean of single primes, the rank of elliptic curves at the centre, and the central value of the congruent-number curves. Section 10 records, for each problem, which of its claims the generating step returns whole.

Main results.

In one line each:

- (i) The step from the clocks up to P to the clocks up to P' builds the new world from $P' - 2$ exact copies of the old one (Theorem 2). The count $\prod(p - 2)$, the palindrome and the copy sign $\prod(p - 4) > 0$ are preserved (Theorems 1 and 3).
- (ii) The two members of a pair are related only by exclusion at each clock; everything else is the product of the members (Theorem 4). Every operator of the clocks is block-diagonal in the members, and the mirror exchanges the members while fixing every clock (Theorem 5).
- (iii) Under the axiom that independence is the absence of a relation (Axiom 1), the twin law holds and the twin pair recurs at every level (Theorem 6).
- (iv) In Maier's windows, where the twins swing 18% above and 8% below the average, the clocks predict the swing to within 0.05% (Table 3).
- (v) Twins are forced by majority wherever primes outnumber the other rough numbers (Theorem 7). Distribution information certifies at most 48.837% (Proposition 3). In the reach of the clocks, $u = 2$, the bound is the equality $\text{twins} = U$, and each member keeps $e^\gamma/2$ of its share over a turn (Proposition 4).
- (vi) A single is its own copy $\lfloor \sqrt{x} \rfloor$ times; a pair never is (Theorem 8). The squares sit on the + side, and so does the lean of prime pairs (Theorem 9 and Proposition 5).
- (vii) The central value of the congruent-number curve is $4\kappa_n m^2$ with $m \in \mathbb{Z}$ (Theorem 10).

2. The World of Clocks

Definition 1 (centres, clocks, full zeros). *A centre is an integer k . It carries the pair $(6k - 1, 6k + 1)$, the lower and upper member. For a prime $p \geq 5$ let $r_p \equiv 6^{-1} \pmod{p}$. The clock p marks the centres*

$$M_p = \{k : k \equiv r_p \text{ or } k \equiv -r_p \pmod{p}\} = \{k : p \mid (6k - 1)(6k + 1)\}.$$

At level P the full zeros are $Z_P = \{k : k \notin M_p \text{ for all primes } 5 \leq p \leq P\}$. The pattern is periodic with period, the turn, $Q_P = \prod_{5 \leq p \leq P} p$.

The two marks of a clock are r_p and $-r_p$; they differ by $2r_p \equiv 3^{-1} \pmod{p}$. The mark $k \equiv r_p$ is the lower member divisible by p , and $k \equiv -r_p$ the upper.




Figure 1. The clocks 5, 7, 11, 13 over the centres $0 \leq k \leq 72$. Each clock marks two centres per turn, one for each member of the pair. Black squares are the full zeros that are twin primes. Grey squares are full zeros beyond the reach of these clocks, where a member may be a product of two primes above 13. Left of the dotted line every full zero is a twin prime pair.

Proposition 1 (the mirror window is the twin question). *Let $P < P'$ be consecutive primes and $k \geq 1$ with $6k - 1 > P$ and $6k + 1 < P'^2$. Then $k \in Z_P$ if and only if $6k - 1$ and $6k + 1$ are both prime.*

Proof. A number below P'^2 that is prime to 6 and to every prime up to P has no prime factor below P' , so it is 1 or a prime. $\square$

The *reach* of the clocks up to P is therefore the range of centres $6k + 1 < P'^2$. The window of centres around 0 is the *mirror window*: it is where the actual primes live. Every other window of a turn is a rotation of the clocks, in which a full zero is a pair of numbers free of small factors but not necessarily prime (Figure 3).

3. What the Step Preserves

Theorem 1 (count and palindrome). *In each turn of level P there are exactly $\prod_{5 \leq p \leq P} (p - 2)$ full zeros, and $Z_P = -Z_P$.*

Proof. Going from P to the next prime P' , each full zero k of a turn of level P lifts to the P' centres $k + jQ_P$, $0 \leq j < P'$. These have distinct residues modulo P' , because Q_P is prime to P' , so exactly two of them are marked by the new clock. Hence $|Z_{P'}| = (P' - 2)|Z_P|$ per turn. The marks $\pm r_p$ are symmetric under $k \mapsto -k$, so each level is a palindrome. $\square$

Theorem 2 (the step builds the world from copies of itself). *Let $P < P'$ be consecutive primes and c a residue class modulo P' . If $c \equiv \pm r_{P'}$, the class contains no full zero of level P' . Otherwise*

$$\{j \in \mathbb{Z} : c + P'j \in Z_{P'}\} = \{j \in \mathbb{Z} : c + P'j \in Z_P\},$$

which is the full-zero set of level P read along the affine map $j \mapsto c + P'j$: a copy of level P with the same count per turn, symmetric about its own centre.

Proof. The clock P' marks exactly the classes $\pm r_{P'}$. On every other class it marks nothing, so a centre $c + P'j$ is a full zero of level P' if and only if it is one of level P . Since P' is invertible modulo Q_P , the map $j \mapsto c + P'j$ runs once through all residues modulo Q_P in a period, which gives the count. $\square$

The new world is made of the old one: $P' - 2$ copies and two empty classes (Figure 2). This is the precise sense in which the pattern at every level is the same simple rule recurring. The step also preserves the full distribution of marks.




Figure 2. Level 13 read class by class modulo 13. Eleven classes are exact copies of level 11, each with 135 full zeros. The two classes $\pm r_{13}$ are the marks of the new clock and are empty. The same holds exactly for $13 \rightarrow 17$ (1485 per class) and $17 \rightarrow 19$ (22275 per class).

Theorem 3 (marking law and copy sign). *In a turn of level P , the number of centres marked by exactly j clocks is the coefficient of x^j in $\prod_{5 \leq p \leq P} ((p-2) + 2x)$. In particular the copy sign $(-1)^{\#\text{clocks marking } k}$ sums over a turn to $\prod_{5 \leq p \leq P} (p-4) > 0$.*

Proof. In the lifting of Theorem 1, each centre lifts to $p-2$ centres that the new clock leaves unmarked and 2 that it marks. Set $x = -1$ for the sign. $\square$

Both identities were checked exactly over full turns for $P = 7, 11, 13, 17, 19$ (sign sums 3, 21, 189, 2457, 36855).

Windows do not return whole.

Inside each copy of Theorem 2, a window of L consecutive centres of level P' appears as a window of length about L/P' . A statement of the form “every window of length L_P holds a full zero” therefore passes from level P to level P' only if $L_{P'} \leq P' L_P$. The reach grows like P^2 while the copies grow like P , so window statements about the reach never return whole under the step: each application hands the question to a shorter window one modulus deeper. The step returns whole the counts over a turn, not the windows.

The strongest window statement, that the longest run $H(P)$ of centres without a full zero is shorter than the reach, has been computed exactly for all rotations of the clocks up to 41 (Table 1). Asked of all rotations, it is at least as demanding as a two-class form of Jacobsthal’s problem. For one class, the best proved bound is still of the order of the square of the largest clock [11], and the longest runs are known to exceed the average spacing by more than a logarithm [6]. The twin question asks for much less: it concerns only the mirror window. That window is not a worst case. At $P = 23$ it holds 35 full zeros, against a mean of 28.87 and a maximum of 39 over the 37,182,145 windows of the turn (Figure 3).

Remark 1 (window statistics over a turn). *Over a full turn, the joint law of two full zeros at distance h is exact: $\rho_2(h) = \prod_p (p - c_p(h)) / p$, where $c_p = 2$ if $p \mid h$, $c_p = 3$ if $3h \equiv \pm 1 \pmod{p}$, and $c_p = 4$ otherwise. This gives the variance of the number of full zeros in a window exactly. It is 3.781 at $P = 19$ and 5.157 at $P = 23$, against the enumerated 3.76 and 5.15. The ratio of variance to mean is 0.18 to 0.32 for $P \leq 1601$, so full zeros are three to five times more evenly spread than independent points. The counts are Gaussian over the turn, and their lower tail is thinner than the Gaussian at the exact variance. Compare the singular-series variance of primes in short intervals [8, 18]. Brun’s method, subtracting the marks clock by clock, certifies a full*

zero in every window only up to a length $P^{e(P)}$, with $e(P) = 4.17$ at $P = 29$ and 8.94 at $P = 10^6$ [1]. The dimension-two sieve [5] is the refined form of the same instrument.

Table 1. Exact longest runs over all rotations of the clocks up to P , against the reach. The ratio stays between 0.31 and 0.42. Following only the sparsest windows from level to level reproduces $H(P)$ exactly up to $P = 37$. At 41 the longest run comes from a window that was not among the sparsest at the lower levels.

P	11	13	17	19	23	29	31	37	41
longest run $H(P)$	6	10	17	24	33	42	57	87	90
reach $(P^2 - P)/6$	18.3	26	45.3	57	84.3	135.3	155	222	273.3




Figure 3. Full zeros in all windows of 135 centres over a turn of the clocks up to 23. The mirror window, the one that contains the actual numbers, holds 35, near the top of the distribution; the fewest in any rotation is 18.

4. The Exclusion Law

Theorem 4 (exclusion law). *Let k be a uniformly chosen centre of a turn of level P . For each clock $p \leq P$ let $X_p = 1$ if $p \mid 6k - 1$ and $Y_p = 1$ if $p \mid 6k + 1$, and 0 otherwise. Then:*

- (i) *the pairs (X_p, Y_p) for different clocks are independent;*
- (ii) $\Pr(X_p = 1) = \Pr(Y_p = 1) = 1/p$;
- (iii) $\Pr(X_p = 1, Y_p = 1) = 0$.

The two members are bound only by (iii). In particular

$$\Pr(k \in Z_P) = \prod_{5 \leq p \leq P} \left(1 - \frac{2}{p}\right) = \underbrace{\prod \left(1 - \frac{1}{p}\right)}_{\text{lower member}} \underbrace{\prod \left(1 - \frac{1}{p}\right)}_{\text{upper member}} \underbrace{\prod \left(1 - \frac{1}{(p-1)^2}\right)}_{\text{exclusion}}.$$

Proof. (i) is the Chinese remainder theorem over the turn. (ii) holds because each of the classes $\pm r_p$ is one class of p . (iii) holds because $r_p \not\equiv -r_p$ for $p \geq 5$: no prime divides two numbers that differ by 2. The factorisation is the identity $1 - 2/p = (1 - 1/p)^2 (1 - (p-1)^{-2})$. $\square$

Corollary 1 (the twin constant is the exclusion product). *The share of full zeros relative to the product of the members' survival shares tends, as $P \rightarrow \infty$, to*

$$E = \prod_{p \geq 5} \left(1 - \frac{1}{(p-1)^2}\right) = 0.880215\dots, \quad \frac{3}{4}E = C_2 = \prod_{p \geq 3} \left(1 - \frac{1}{(p-1)^2}\right) = 0.66016\dots$$

The factor $\frac{3}{4}$ is the clock 3, already absorbed by writing every pair as $6k \pm 1$.

Theorem 4 is a complete list of the relations that the clocks contain. The only mechanism in the construction that ties one member to the other is a clock. A clock ties them in exactly one way, by taking one member and thereby leaving the other.

4.1. The Exclusion Operator

The exclusion law has an exact form as a statement about operators. A state of the pair over a turn of level P is a vector $(x_-, x_+) \in \mathbb{C}^{\mathbb{Z}/Q_P} \oplus \mathbb{C}^{\mathbb{Z}/Q_P}$, the lower and the upper member. Put $\Gamma = \text{diag}(I, -I)$ and $P_{\pm} = \frac{1}{2}(I \pm \Gamma)$, the projections onto the two members, let Σ exchange the members, $\Sigma(x_-, x_+) = (x_+, x_-)$, and let S and R be the shift $k \mapsto k + 1$ and the reflection $k \mapsto -k$ of the centres.

Definition 2 (the exclusion operator). *The exclusion operator of the clock p is $E_p = \text{diag}(\delta_{r_p}, \delta_{-r_p})$, multiplication on the lower member by the indicator of the class r_p and on the upper member by the indicator of $-r_p$. The algebra of the clocks $\mathcal{E}_P$ is generated by the E_p , $5 \leq p \leq P$, and the common shift $\text{diag}(S, S)$. The mirror of the pair is $\iota = \Sigma \text{diag}(R, R)$.*

Theorem 5 (the clock acts on one member). (i) *Every $E \in \mathcal{E}_P$ commutes with Γ , so $P_+EP_- = P_-EP_+ = 0$: no clock, no product or sum of clocks and no motion of the centre carries a state of one member into a state of the other.*
(ii) *A clock annihilates its own exchange: $E_p(\Sigma E_p \Sigma) = 0$.*
(iii) *The mirror exchanges the members and fixes every clock: $\iota \Gamma \iota^{-1} = -\Gamma$ and $\iota E_p \iota^{-1} = E_p$.*

Proof. (i) The generators are block-diagonal, block-diagonal operators form an algebra, and each of them commutes with Γ ; since $P_{\pm}$ are polynomials in Γ , $P_+EP_- = EP_+P_- = 0$. (ii) $\Sigma E_p \Sigma = \text{diag}(\delta_{-r_p}, \delta_{r_p})$, and $\delta_{r_p} \delta_{-r_p} = 0$ because $2r_p \equiv 3^{-1} \not\equiv 0 \pmod{p}$. (iii) $\Sigma \Gamma \Sigma = -\Gamma$, $\text{diag}(R, R)$ commutes with Γ , and $R \delta_a R = \delta_{-a}$, so $\iota E_p \iota^{-1} = \Sigma \text{diag}(\delta_{-r_p}, \delta_{r_p}) \Sigma = E_p$. $\square$

Remark 2 (the same arrangement in a geometry). *The arrangement of Theorem 5 is that of the two strands of the medial antiprism, a stack of triangles each turned through 180° against the next, with every vertex joined to every side midpoint of the neighbouring layers. The two strands share every layer and no line, and the central inversion of the stack exchanges them while carrying every bundle of lines onto a bundle of lines, as ι exchanges the members and fixes every clock.*

5. The Pair Measured Against the Exclusion Law

The joint law of prime factors.

For the $5 \cdot 10^7$ centres with $6k + 1 \leq 3 \cdot 10^8$ let N_{ab} count the centres with $\Omega(6k - 1) = a$ and $\Omega(6k + 1) = b$. Put $D_{ab} = N_{ab}K / (N_a N_b)$, where K is the number of centres and N_a, N_b are the marginal counts. If the members were unbound, every D_{ab} would be 1. Figure 4 shows the table. The prime-prime entry is $D_{11} = 0.8833$, against the exclusion product $E = 0.8802$. The other entries show the smooth negative association that exclusion produces: a clock taken by one member cannot be taken by the other, so large factor counts avoid each other.




Figure 4. Left: the joint law of $\Omega(6k-1)$ and $\Omega(6k+1)$ relative to independence, over $5 \cdot 10^7$ pairs. The prime–prime entry equals the exclusion product to 0.3%. Right: the pair symbol $G(z, w)$ tested for factorisation on the circle $z = w = r$. As $r \rightarrow 0$ the symbol isolates the prime–prime coefficient and the defect tends to the exclusion constant; at $r = 1$ it is 1.

The pair symbol.

Define

$$G(z, w; x) = \sum_{6k+1 \leq x} z^{\Omega(6k-1)} w^{\Omega(6k+1)}.$$

For fixed z and w with $|z|, |w| \leq 1$, the map $n \mapsto z^{\Omega(n)}$ is completely multiplicative and bounded. Exchanging z and w is the mirror of the pair. The number of twin centres is the single coefficient $[z^1 w^1]G$, so “the members are unbound” is the statement that $G(z, w)$ factorises as $G(z, 1)G(1, w)/K$ up to the exclusion. Measured on the circle $z = w = r$, the defect of factorisation is 0.973 at $r = 0.5$, 0.937 at 0.25, 0.907 at 0.1 and 0.888 at 0.02. As $r \rightarrow 0$ it tends to the exclusion constant, as it must if exclusion is the only bond (Figure 4).

The rough frame.

Fix z and $y = z^u$. Call a pair *rough* when both members are free of every prime up to z . On this frame each member is either prime or a product of primes above z , and Theorem 4 has already been applied to every clock up to z . Let R be the share of twins among rough pairs divided by the product of the prime shares of the two members. Table 2 and Figure 5 give R for z up to 1000 and y up to 10^9 . It is 1 to within 1% throughout: beyond the clocks already accounted for, the members decide their primality as two unbound numbers.

Table 2. The share of twins among rough pairs over the product of the members’ prime shares.

	$z = 100$	$z = 200$	$z = 400$	$z = 1000$
R at $u = 2.5$	1.0045	1.0051	1.0082	1.0053
R at $u = 3.0$	1.0016	1.0090	1.0043	1.0029
rough pairs at $u = 3.0$	19,296	115,416	741,104	8,775,234




Figure 5. The frame ratio R of Table 2. The members of a rough pair are prime or not as two unbound numbers would be.

Maier's windows.

Maier showed that the number of primes in intervals of length $(\log x)^\lambda$ swings above and below its average infinitely often [15]. His windows $(Qk, Qk + y]$, with $Q = \prod_{p \leq z} p$ and $y = z^u$, start at whole turns of the clocks up to z . So every clock marks them exactly as it marks $(0, y]$: they are copies of the mirror window, and the swing is a clock effect. The exclusion law then predicts the count in each window from its own marks. Each member free of the primes up to z is prime with probability $1/(M_z \log n)$, $M_z = \prod_{p \leq z} (1 - \frac{1}{p})$, and each such pair with probability $E_{>z}/(M_z^2 \log n \log(n+2))$, $E_{>z} = \prod_{p > z} (1 - \frac{1}{(p-1)^2})$. Table 3 counts all of Maier's windows with $10^9 < Qk < 10^{10}$ for $z = 13$ (299,699 windows) and $z = 17$ (17,629), against the same number of windows placed at random. The twins swing by +18% and −8%. The clocks predict the swing, to 0.9995 ± 0.0006 over 2.65 million pairs, and they predict the single members to within 0.2% in every row. In Maier's own windows, once the clocks are counted, the members are prime independently.

Table 3. Twin pairs in Maier's windows near 10^9 – 10^{10} : the swing against the average, and the count against the prediction of the clocks.

	twins	clock prediction	twins / average	twins / prediction
$z = 17, u = 2$	16,013	16,152.0	1.183	0.991 ± 0.008
$z = 17, u = 2.5$	51,381	51,686.3	0.921	0.994 ± 0.004
$z = 17, u = 3$	228,873	229,357.8	0.993	0.998 ± 0.002
$z = 13, u = 2$	145,134	145,369.7	1.082	0.998 ± 0.003
$z = 13, u = 2.5$	484,039	484,565.7	1.000	0.999 ± 0.001
$z = 13, u = 3$	1,728,629	1,728,284.3	0.986	1.000 ± 0.001
all	2,654,069	2,655,415.8		0.9995 ± 0.0006

6. The Axiom and the Twin Law

The construction contains one bond between the members of a pair, and Theorem 4 proves that the bond is exclusion. Under the axiom below, independence is what remains when the relations have been listed.

Axiom 1 (independence is the absence of a relation). *Two numbers are related only through a mechanism present in their construction. For the members $6k \pm 1$ of a pair, the only mechanism is a clock, and a clock relates them only by exclusion (Theorems 4 and 5). A dependence between the members beyond exclusion, in any stretch of the integers, would be a relation without a clock.*

Read on initial segments, Axiom 1 gives the law of Hardy and Littlewood (Theorem 6). The measurements of Sections 5 and 7 test it everywhere a further bond could appear.

Theorem 6 (the twin law). *Under Axiom 1, the number of twin prime pairs up to x satisfies*

$$\pi_2(x) \sim \frac{1}{6} \int_2^x \left(\frac{3}{\log t} \right)^2 E dt = 2C_2 \int_2^x \frac{dt}{\log^2 t} \sim \frac{2C_2 x}{\log^2 x},$$

and there are infinitely many twin primes.

Proof. The centres up to x number $x/6$. Among the numbers $\equiv \pm 1 \pmod{6}$ near t , the primes have density $3/\log t$, because every prime above 3 lies in these two classes, which hold a third of the integers. By Theorem 4 and Axiom 1, the two members are prime as two unbound numbers, corrected at every clock by the exclusion factor, whose product is E . Summing over the centres gives the integral. Finally $\frac{3}{2}E = 2C_2$ by Corollary 1. $\square$

The density $3/\log t$ and the constant $2C_2$ agree with the prediction of Hardy and Littlewood [10]. In the present reading, that constant is nothing but the exclusion of the clocks.

The count.

Table 4 and Figure 6 compare the law with the count of twin pairs to 10^{10} . From 10^7 on, the count and the law agree to within 4 parts in 10^3 . At 10^{10} they agree to 4.6 parts in 10^5 .

Table 4. The twin law against the count of twin prime pairs.

x	twin pairs $\leq x$	law $2C_2 \int_2^x dt / \log^2 t$	ratio
10^4	205	214.2	0.9570
10^5	1,224	1,248.7	0.9802
10^6	8,169	8,248.0	0.9904
10^7	58,980	58,753.8	1.0039
10^8	440,312	440,367.8	0.99987
10^9	3,424,506	3,425,308.2	0.99977
10^{10}	27,412,679	27,411,416.5	1.00005




Figure 6. The count of twin pairs up to x divided by the twin law, at quarter decades from 10^3 to 10^{10} .

Where the burden lies.

In the convention of analytic number theory, carrying the exclusion law from the full turn of the clocks to the initial stretch $[1, x]$ is the content of the Hardy–Littlewood conjecture for twins, and is regarded as open. This paper asks instead what that transfer could fail by. A failure would be a correlation between $6k - 1$ and $6k + 1$ carried by no clock. It would have to show up somewhere, and we have looked for it in every place it could appear:

- per clock (Theorem 4);
- jointly over all clocks, in the full law of $(\Omega(6k - 1), \Omega(6k + 1))$;

- in the pair symbol at every radius;
- on the frame of rough pairs, after all small clocks have acted;
- in the count to 10^{10} .

The only departure from the product measured anywhere is the exclusion constant, to four decimals. The sign version of the axiom, the independence of the parities $\lambda(n)$ and $\lambda(n+2)$ of the number of prime factors of neighbours, has been proved in logarithmic mean [22]. Gaps between primes are known to be bounded infinitely often [16,24].

7. The Instrument and Its Ceiling

This section proves that twins are forced by a majority, with no appeal to independence. It then measures exactly how much of that majority can be certified by information about the distribution of primes in residue classes, the information on which sieve methods rest [7].

Theorem 7 (twins by majority). *Fix z and y . Let U be the number of rough pairs with $6k+1 \leq y$, and let A and B be the numbers of those whose lower, respectively upper, member is prime. Then the number of twin pairs among them is at least $A + B - U$. In particular, if more than half of the rough members on each side are prime, the range $(z, y]$ contains a twin pair.*

Proof. For indicators, $[\text{lower prime}][\text{upper prime}] \geq [\text{lower prime}] + [\text{upper prime}] - 1$ pointwise. Sum over the rough pairs. $\square$

Proposition 2 (the majority horizon). *Among the numbers up to y free of the primes up to $z = y^{1/u}$, the primes have share $1/(u\omega(u))$ asymptotically, where ω is Buchstab's function [7]. This share exceeds $\frac{1}{2}$ exactly for $u < u^*$, with $u^* = 3.565$.*

On the frame of rough pairs the measured prime share of each member exceeds one half up to $u \approx 3.8$ for z between 30 and 170, and approaches $1/(u\omega(u))$ as z grows (Table 5 and Figure 7). Theorem 7 then forces twins in every such range, with a guaranteed share of at least $2x - 1$ of the frame, where x is the prime share of each member.

Table 5. Prime share of the members of rough pairs, and the share of twins it forces by Theorem 7. At $z = 1000$, $u = 3$ the forced share is 0.247; the actual share is 0.390.

	$u = 3.0$	$u = 3.5$	$u = 3.75$	$u = 4.0$
$z = 30$: prime share / forced twins	0.700 / 0.383	0.587 / 0.165	0.545 / 0.085	0.506 / 0.009
$z = 60$	0.666 / 0.325	0.563 / 0.127	0.524 / 0.048	0.488 / –
$z = 170$	0.645 / 0.289	0.548 / 0.097	0.510 / 0.019	0.476 / –
$z = 1000$	0.624 / 0.247			

Proposition 3 (the ceiling of the instrument). *Suppose only the distribution of primes in residue classes is used, at the full level, and the frame count U is known exactly. Then the largest share of rough members that can be certified prime is*

$$\max_u \frac{1}{u\omega(u)} \cdot \frac{f(u)}{e^{\gamma}\omega(u)} = 0.48837, \quad \text{attained at } u = 3.16,$$

where f is the lower function of the linear sieve, which is optimal for this information [7].

Allowing the two members different levels of roughness, the best certification of $A + B$ against U is:

- 0.97881 at level $\frac{1}{2}$;
- 0.98034 at level $\frac{4}{7}$;
- 0.98058 at level $\frac{3}{4}$;
- 0.99975 at the full level.

The value 1 is approached only as one member's roughness reaches the square root of the range. There that member is automatically prime, and Theorem 7 reads $\text{twins} \geq \text{twins}$.




Figure 7. The object and the instrument. Blue: the prime share of rough numbers, above one half for $u < 3.565$, where Theorem 7 forces twins. Green: measured on the pair frame. Orange: the largest share that distribution information can certify, which never reaches one half (maximum 0.48837).

Reading.

A sieve sees only divisibility by small numbers. It cannot tell a prime from a product of two large primes, because both are free of small factors. Proposition 3 measures that blindness exactly. In the symmetric frame the instrument falls 1.16% short of a majority. In the best asymmetric frame it closes the gap only where the inequality has become a tautology. Chen's method, which adds the switch of the two members, reaches a prime and a product of at most two primes [2] and stops at the same place. This is the parity barrier. It is a statement about what the instrument can see: the instrument cannot certify the majority that Proposition 2 computes. It says nothing about whether the members of a pair are bound, which is the question of Sections 4 and 5. In terms of sums: a sieve receives about a sequence only its sums over the multiples of each d [7]. The Liouville sign $\lambda(n) = (-1)^{\Omega(n)}$ has sum $\lambda(d)L(x/d)$ there, which is small: up to 10^7 it is at most 1.0% of x/d for $d \leq 1000$, and at most 1.2% of the count in every class modulo $q \leq 60$. So $1 + \lambda$ and $1 - \lambda$ give a sieve the same data, while one vanishes on the primes and the other doubles them. The parity of the two members is itself unbound. Tao proved that $\lambda(n)\lambda(n+2)$ has logarithmic mean zero [22], and over the 1.67 million twin centres $1000 \leq k \leq 1,666,666$ the mean of $\lambda(6k-1)\lambda(6k+1)$ is $+0.00001 \pm 0.0008$.

The reach as a frame.

The reach of the clocks (Proposition 1) is a frame of rough pairs with $z = P$ and $y = P'^2$. So $u = 2 \log P' / \log P$, which tends to 2. There Theorem 7 becomes an equality. Buchstab's function at $u = 2$ fixes the share of each member that the frame keeps against the turn.

Proposition 4 (the frame at $u = 2$). *Let $P < P'$ be consecutive primes with $P \geq 5$. Take the centres of the reach, those with $6k - 1 > P$ and $6k + 1 < P'^2$. Let n be their number, U the number of rough pairs among them, and A and B the numbers of those whose lower, respectively upper, member is prime.*

- (i) $A = B = U$, and the number of twin pairs among these centres is exactly $A + B - U = U$.
- (ii) The prime share $1/(u \omega(u))$ of Proposition 2 equals 1 for $1 < u \leq 2$.
- (iii) As $P \rightarrow \infty$,

$$\frac{A}{n \prod_{5 \leq p \leq P} (1 - 1/p)} \rightarrow e^\gamma \omega(2) = \frac{e^\gamma}{2} = 0.89054,$$

and the same holds for B .

Proof. (i) By Proposition 1 a rough member of the reach is prime, so $A = U = B$. By Theorem 7 the number of twin pairs is at least $A + B - U$, and it is at most U . (ii) Buchstab's function satisfies $u\omega(u) = 1$ for $1 \leq u \leq 2$ [7]. (iii) The lower members are the primes $q \equiv 5 \pmod{6}$ with $P < q < P'^2 - 2$. By the prime number theorem for the progressions modulo 6 [4], $A \sim P'^2/(4\log P')$. Also $n \sim P'^2/6$, and by Mertens' theorem [17] $\prod_{5 \leq p \leq P}(1 - 1/p) = 3 \prod_{p \leq P}(1 - 1/p) \sim 3e^{-\gamma}/\log P$. Since $P' < 2P$, $\log P'/\log P \rightarrow 1$, so the ratio tends to $e^\gamma/2$. The upper members are the primes $q \equiv 1 \pmod{6}$ and have the same count. $\square$

Over a full turn the share of full zeros is exactly $\prod_{5 \leq p \leq P}(1 - 2/p)$, and the share of each member is $\prod(1 - 1/p)$ (Theorem 4). The reach is not a turn. By Proposition 4(iii), each member keeps $e^\gamma\omega(2)$ of its turn share there. In Buchstab's terms the turn share is the limit $u \rightarrow \infty$, where $e^\gamma\omega(u) \rightarrow 1$ [7]. Under Axiom 1 the pair keeps the product of its members' factors. The twin law (Theorem 6) at $x = P'^2$ gives

$$U \sim \frac{2C_2P'^2}{\log^2(P'^2)} \sim \left(\frac{e^\gamma}{2}\right)^2 n \prod_{5 \leq p \leq P} \left(1 - \frac{2}{p}\right), \quad \frac{e^{2\gamma}}{4} = 0.79305.$$

Table 6 measures the reach up to $P = 31601$, where $P'^2 > 10^9$:

- The pair keeps the product of its members' factors to within 1.5% for every $P \geq 211$, and to within 0.3% at $P = 31601$.
- The members keep 0.9387 of their turn share. To within 0.04%, this is $e^\gamma/2$ times $\pi(y) \log y/y = 1.054$ at $y = P'^2$.
- The pair keeps 0.8839, and its factor falls with P toward 0.7931.

Table 6. The reach of the clocks up to P as a frame at $u = 2$: the centres with $6k - 1 > P$ and $6k + 1 < P'^2$. The turn share is $n \prod_{5 \leq p \leq P}(1 - 2/p)$. Lower and upper: the prime members divided by $n \prod_{5 \leq p \leq P}(1 - 1/p)$. The limits are $e^{2\gamma}/4$ and 1 under the twin law, and $e^\gamma/2$ by Proposition 4(iii).

P	centres n	twins U	turn share	$U/\text{turn share}$	lower	upper	pair/(lower·upper)
23	135	29	28.87	1.0044	1.0415	1.0113	0.9537
101	1,750	208	197.08	1.0554	1.0233	1.0009	1.0304
211	8,252	685	699.47	0.9793	0.9919	0.9841	1.0034
503	43,095	2,631	2,731.23	0.9633	0.9828	0.9778	1.0024
1009	170,859	8,332	8,856.47	0.9408	0.9685	0.9673	1.0043
2003	673,685	27,075	28,938.79	0.9356	0.9609	0.9594	1.0148
5003	4,180,845	130,803	143,413.65	0.9121	0.9516	0.9514	1.0075
10007	16,695,011	440,825	490,176.00	0.8993	0.9462	0.9461	1.0045
20011	66,803,404	1,510,202	1,698,922.62	0.8889	0.9413	0.9412	1.0033
31601	166,495,140	3,420,966	3,870,157.36	0.8839	0.9387	0.9387	1.0031
limit				0.7931	0.8905	0.8905	1

What the frame at $u = 2$ decides.

In the reach, Theorem 7 reads twins = U . This is the degenerate case of Proposition 3, where the inequality has become twins $\geq$ twins, and it does not give the size of U . The count U is the number of twin pairs between P and P'^2 . The share over a full turn, $n \prod_{5 \leq p \leq P}(1 - 2/p)$, is not that count. In the reach each member keeps $e^\gamma\omega(2)$ of its turn share, and at $P = 31601$ the twin pairs number 0.884 of the turn share. That $U > 0$ for every P means a twin pair between every prime and the square of the next. That is a statement about the twin primes themselves, and it implies that there are infinitely many. The reach is therefore the place where Axiom 1 acts. The members' factors are theorems. That the pair keeps their product is the exclusion law carried from the turn into the reach.

8. Singles Copy Themselves; Pairs Never Do

Let λ be Liouville's function, the parity of the number of prime factors, and $L(x) = \sum_{n \leq x} \lambda(n)$.

Theorem 8 (a single is its own copy; a pair is not). (i) For every $x \geq 1$, $\sum_{n \leq x} \lambda(n) \lfloor x/n \rfloor = \lfloor \sqrt{x} \rfloor$.
(ii) For every even D the equation $n(n+D) = m^2$ has finitely many solutions, all with $n < D^2/8 + 1$. Hence $\sum_{n \leq x} \sum_{d|n(n+D)} \lambda(d)$ is bounded in x . For twins, $n(n+2) = (n+1)^2 - 1$ is never a square, and the sum is 0 for every x .
(iii) For every $x \geq 1$, $L(x) = \sum_{m \leq x} \mu(m) \lfloor \sqrt{x/m} \rfloor$.

Proof. (i) $\sum_{d|n} \lambda(d)$ is 1 when n is a square and 0 otherwise; sum over $n \leq x$. (ii) $(n+D/2-m)(n+D/2+m) = D^2/4$ has finitely many factorisations. (iii) is the Möbius inversion of (i). $\square$

The sign-weighted count of single numbers is exactly the number of self-copies, the squares. That is the origin of the lean of the primes by one half. A pair's product is never its own copy, so a pair has no lean of its own: any lean of prime pairs must come from the copies of its members. Identity (iii) was checked exactly at eight values of x up to 10^5 . Its first term, $\sqrt{x} \sum_{m \leq x} \mu(m) m^{-1/2}$, is the source of the logarithmic mean of $L(x)/\sqrt{x}$, measured as -0.682 against $1/\zeta(\frac{1}{2}) = -0.685$.

Theorem 9 (the squares sit on the + side). Every prime $q \geq 5$ has $q^2 \equiv 1 \pmod{24}$, so q^2 is the upper member $6k+1$ of its triad $6k-1, 6k, 6k+1$, on the + side. For a pair $(n, n+D)$ of numbers prime to 6:

- if $D \equiv 2 \pmod{6}$, only the upper member can be a prime square ($3 \mid q^2 + 2$);
- if $D \equiv 4 \pmod{6}$, only the lower member can be ($3 \mid q^2 - 4$);
- if $D \equiv 0 \pmod{6}$, both can be when $n \equiv 1 \pmod{6}$, and neither when $n \equiv 5 \pmod{6}$.

Proposition 5 (the lean of a pair lives on its + member). Suppose the pairs of prime powers $(n, n+D)$ are balanced over residue classes in the logarithmically weighted count. Then the class lean of prime pairs of gap $D \equiv 2 \pmod{6}$ is a function of the residue symbol of the upper member alone. The same holds for the lower member when $D \equiv 4$, and for the pairs on the + side when $D \equiv 0$. The model in which a pair's lean is the product of its members' leans is incompatible with this for every modulus $q \geq 5$.

Proof. This is the setting of the biases among prime pairs studied in [14]. Under the balance, prime pairs carry the negative of the prime-power pairs that contain a prime square. By Theorem 9, for $D \equiv 2$ these are $(q^2 - D, q^2)$. Their centres c have $c + D/2$ a nonzero square modulo q , and they are spread uniformly over those classes. The two symbols are linearly independent on the allowed classes for every $q \geq 5$, which gives the incompatibility. $\square$




Figure 8. Class lean of prime pairs over 428 moduli $q \leq 3000$, per member. For twins the lower member carries no lean (-42 ± 112 against $+3$) and the upper member carries it all (-767 ± 113 against -723). Cousins show the same with the members exchanged. For gap 6 the lower member keeps an excess of about -190 on both sides, about 2.5 standard errors in combination.

The lean across L -functions.

For an L -function with normalised local sign $\lambda(p)$, under the generalised Riemann hypothesis the lean of the primes $\sum_{p \leq x} \lambda(p) \log p / \sqrt{p}$ grows like $(-\nu/2 - r) \log x$. Here ν is the slope of the self-copy sum $\sum \text{tr}(\text{Frob}_p^2) \log p / p$ and r is the order of the zero at the centre. This is the form in which the rank enters the partial Euler products [3,9,12], and the square term is the source of Chebyshev's bias [19,20]. We measured both slopes separately, with primes to 10^9 , on 17 objects (Figure 9):

- six complex characters, which have $\nu = 0.000$ and no lean;
- six real characters, the triad sign among them, with $\nu = 0.999$ and lean -0.49 to -0.50 ;
- two curves of rank 0, with $\nu = -1.00$ and lean $+0.49$;
- a curve of rank 1, with lean -0.48 ($r = 0.98$);
- the symmetric squares, with $\nu = 1.998$ with complex multiplication and 1.000 without.

On the congruent-number curves the rank read from the lean, over 608 curves, sorts into clusters with medians 1.00 and 1.98 . These agree with Tunnell's criterion [23] and with the parity of the root number. The curve $n = 210$ was read as rank 2 from its primes before any point was known; a search then found two independent points, and three for $n = 1254$.




Figure 9. Lean of the primes against the self-copy count, for 17 L -functions. All lie on the lines $-\nu/2 - r$ with $r = 0$ or $r = 1$. Open circles: a curve without complex multiplication (level 11) and its symmetric square.

9. At the Centre: The Congruent-Number Curves

For squarefree n let $E_n : y^2 = x^3 - n^2x$. Its coefficients follow from those of E_1 by the twist $a_p(E_n) = \left(\frac{n}{p}\right)a_p(E_1)$. For E_1 itself, $a_p = 0$ for $p \equiv 3 \pmod{4}$; otherwise write $p = a^2 + b^2$ with a odd, and then $a_p = \pm 2a$, the sign $+$ exactly when $a + b \equiv 1 \pmod{4}$ [13]. Tunnell's formula [23] expresses $L(E_n, 1)\sqrt{n}$ as $\kappa_n(A - 2B)^2$, where A and B count representations of n by two ternary forms, with $\kappa_n = 0.163879$ for odd n and 0.327757 for even n . We confirmed the formula on all 608 squarefree $n \leq 1000$.

Theorem 10 (the value at the centre is a square). *For squarefree n , $L(E_n, 1)\sqrt{n} = 4\kappa_n m^2$ with $m \in \mathbb{Z}$. Hence $L(E_n, 1)$ is either 0 or at least $4\kappa_n/\sqrt{n}$.*

Proof. For odd n , every representation $x^2 + 2y^2 + 8z^2 = n$ (and likewise for the second form) has x odd, so the representations fall into sign classes of even size and A, B are even; for even n the same holds with $n/2$. Hence $A - 2B = 2m$. $\square$

The value at the mirror point is a copy of a count, a square, and it cannot drift continuously to zero. Under the generalised Riemann hypothesis, Hadamard's product for the completed L -function, which is even about the centre, gives exactly $\sum_{\gamma > 0} \gamma^{-2} = S_2/S_0$. Here S_0 and S_2 are rapidly converging sums over the coefficients, and S_0 carries the quantised central value. Hence $\gamma_1 \geq (S_2/S_0)^{-1/2}$.

Table 7 shows one fact read five ways: a nearly balanced count, a small central value, a zero near the centre, a misread rank, and a slow Euler product.

Table 7. The congruent-number curves $n = 347, 563, 587, 659, 683, 691, 787, 827, 922$ against controls. A zero pair near the centre adds about $2 \operatorname{sinc}(\gamma_1 \log x)$ to the rank read from the primes. The Euler products of the controls column are those of $n = 1, 3, 11, 17, 19, 930$. The value $\gamma_1 \approx 0.05\text{--}0.1$ was predicted from the readings before the zeros were computed.

	the 9 curves read as rank 1.2–1.5	12 controls
Tunnell balance $ A - 2B $	4 (smallest nonzero for $n > 300$)	24–36
lowest zero above the centre γ_1	0.053–0.091	0.323–0.537
GRH floor $(S_2/S_0)^{-1/2}$, smallest of 247	0.0375 ($n = 563$)	
Euler product at 10^9 / limit	0.38, 0.50 ($n = 347, 563$)	within 18%

The contrast with ζ .

Take the finite pattern $\sum_{n \leq N} n^{-s}$ with its mirror, cut at $N = \lfloor \sqrt{t/2\pi} \rfloor$. Its off-line pairs, found by complete scans of $10^3 \leq t \leq 10^5$ and of windows near 10^6 and 10^7 , number 525. The remainder of ζ closes every one of them back onto the line. All 525 sit at heights where the prime clocks are anti-aligned: the sum $\sum_{p \leq 400} \cos(t \log p) / \sqrt{p}$ lies at a median percentile of 0.9% among random heights. The closest of them, at $t = 71732.91$, leaves $|Z| = 0.00055$ between two zeros. For the congruent-number curves, Theorem 10 forbids a near-merge at the centre from becoming a merge except exactly. No such floor between neighbouring zeros is known for ζ .

10. Which Claims the Step Returns Whole

A self-reproducing world proves its own persistence in one step: combined with a machine that runs it, the world produces the world again. The same test sorts the claims of this paper.

Returned whole by one step.

- the count $\prod (p - 2)$;
- the palindrome;
- the marking law and the copy sign;
- the family of affine copies (Theorem 2);
- the exclusion law in the world of clocks (Theorem 4);
- the mirror symmetry of the zeros of ζ as a set;
- the twisting family of the congruent-number curves;
- the square at the centre (Theorem 10).

Not returned whole, and why.

- *Window claims about the reach* (Section 3): the reach grows like P^2 , the copies like P .
- *The statement that each zero of ζ is its own mirror image*: the mirror alone returns only the set. Functions with the mirror but without the Euler product, such as Epstein zeta functions and the Davenport–Heilbronn function, have zeros off the line.
- *The pair $(m - 1, m + 1)$ under multiplication*: shift and multiplication do not commute. Seen from the centre, and modulo any prime, the pair is the unit pair ± 1 , and every clock's marks are the square roots of 1. Squaring respects both the mirror and multiplication, and in its terms a twin centre is a square m^2 whose predecessor $m^2 - 1$ splits into exactly two primes.

In every case the claim returns whole under the mirror or under multiplication. What is sought is the operation that is both. For the pair, Theorem 4 shows that nothing beyond these two acts on it.

11. Methods

All computations are exact integer computations unless stated otherwise.

- *Clock identities* (Theorems 1–3) were checked by complete enumeration of turns for $P \leq 19$, and the window statistics over the full turn for $P = 23$. The longest runs $H(P)$ come from exhaustive search over all rotations.
- *Twin counts* to 10^{10} come from a segmented sieve of Eratosthenes. The law was integrated numerically.
- *The joint law of Ω* was computed for all $n \leq 3 \cdot 10^8$ by counting prime powers.
- *The rough frames* used a sieve to 10^9 , primality from the same sieve, and roughness by marking multiples of the primes up to z .
- *The reach as a frame* (Table 6) used a sieve of the numbers $6k \pm 1$ up to 10^9 by the primes up to 31623. Each clock marks its two classes, and $\pi(P'^2) = 50,799,680$ at $P' = 31607$.
- *The Buchstab function and the sieve functions f, F* were integrated from their delay equations with step 10^{-4} . Checks: $\omega(3) = (1 + \log 2)/3$, $\omega(10) = e^{-\gamma}$ to five digits, and $f(4) = 0.97835$.
- *Class leans* were computed over 428 prime moduli $q \leq 3000$ with the weight $\log^2 m / \sqrt{m}$ at the centre m . Standard errors come from the spread of the per-modulus slopes.
- *The self-copy sums* used primes to 10^9 . The coefficients of the curve of level 11 come from its eta product to 10^7 .
- *Central values* used the rapidly converging series, with the conductor ($32n^2$ for odd n , $16n^2$ for even n) confirmed by invariance in the cutoff parameter to $3 \cdot 10^{-14}$. The zeros are sign changes of the completed L -function on the line, by Gauss–Laguerre quadrature with 48 nodes and step 0.002. They are computed, not certified.

12. Conclusions

The paper set out to find what binds the two members of a twin pair. It found one thing. A clock can take one member, and in doing so it leaves the other. That bond is proved exactly, it fixes the twin constant, and every measurement of the pair shows it and nothing else. Read with the axiom that independence is the absence of a relation, it gives the twin law and the recurrence of the twin pair at every level of the clocks. The one obstruction that has kept the question closed to the methods that know the most about primes is, on inspection, a limit of those methods' sight: they cannot tell one factor from two, and so they cannot certify the majority that the object itself carries. In the reach of the clocks the majority bound is an equality. There each member keeps $e^\gamma/2$ of its turn share, and the twins keep the product of the two factors (Proposition 4, Table 6): this is the exclusion law carried into the reach.

Funding: This research received no external funding.

Data Availability Statement: The computations are described in Section 11; the scripts are available from the author on request.

Conflicts of Interest: The author declares no conflicts of interest.

References

1. V. Brun, La série $1/5 + 1/7 + 1/11 + 1/13 + 1/17 + 1/19 + 1/29 + 1/31 + 1/41 + 1/43 + 1/59 + 1/61 + \dots$, où les dénominateurs sont “nombres premiers jumeaux” est convergente ou finie, *Bull. Sci. Math.* 43 (1919) 100–104, 124–128.
2. J. R. Chen, On the representation of a larger even integer as the sum of a prime and the product of at most two primes, *Sci. Sinica* 16 (1973) 157–176.
3. K. Conrad, Partial Euler products on the critical line, *Canad. J. Math.* 57 (2005) 267–297.
4. H. Davenport, *Multiplicative Number Theory*, 3rd ed., revised by H. L. Montgomery, Graduate Texts in Mathematics 74, Springer, New York, 2000.
5. H. G. Diamond, H. Halberstam, *A Higher-Dimensional Sieve Method*, with procedures for computing sieve functions by W. F. Galway, Cambridge Tracts in Mathematics 177, Cambridge University Press, 2008.
6. K. Ford, B. Green, S. Konyagin, J. Maynard, T. Tao, Long gaps between primes, *J. Amer. Math. Soc.* 31 (2018) 65–105.

7. J. Friedlander, H. Iwaniec, *Opera de Cribro*, American Mathematical Society Colloquium Publications 57, 2010.
8. P. X. Gallagher, On the distribution of primes in short intervals, *Mathematika* 23 (1976) 4–9.
9. D. Goldfeld, Sur les produits partiels eulériens attachés aux courbes elliptiques, *C. R. Acad. Sci. Paris Sér. I Math.* 294 (1982) 471–474.
10. G. H. Hardy, J. E. Littlewood, Some problems of ‘Partitio numerorum’; III: On the expression of a number as a sum of primes, *Acta Math.* 44 (1923) 1–70.
11. H. Iwaniec, On the problem of Jacobsthal, *Demonstratio Math.* 11 (1978) 225–231.
12. S. Kim, M. R. Murty, From the Birch and Swinnerton-Dyer conjecture to Nagao’s conjecture, *Math. Comp.* 92 (2023) 385–408.
13. N. Koblitz, *Introduction to Elliptic Curves and Modular Forms*, 2nd ed., Graduate Texts in Mathematics 97, Springer, New York, 1993.
14. R. J. Lemke Oliver, K. Soundararajan, Unexpected biases in the distribution of consecutive primes, *Proc. Natl. Acad. Sci. USA* 113 (2016) E4446–E4454.
15. H. Maier, Primes in short intervals, *Michigan Math. J.* 32 (1985) 221–225.
16. J. Maynard, Small gaps between primes, *Ann. of Math.* 181 (2015) 383–413.
17. F. Mertens, Ein Beitrag zur analytischen Zahlentheorie, *J. Reine Angew. Math.* 78 (1874) 46–62.
18. H. L. Montgomery, K. Soundararajan, Primes in short intervals, *Comm. Math. Phys.* 252 (2004) 589–617.
19. M. Rubinstein, P. Sarnak, Chebyshev’s bias, *Experiment. Math.* 3 (1994) 173–197.
20. P. Sarnak, Letter to B. Mazur on Chebyshev’s bias for $\tau(p)$, 2007, <https://publications.ias.edu/sites/default/files/MazurLtrMay08.PDF>.
21. A. Singh, The square-root horizon: squares, reversal and prime clocks in the distribution of the primes and the zeros of the Riemann zeta function, *Preprints* 2026, 2026100430, doi:10.20944/preprints202610.0430.v1.
22. T. Tao, The logarithmically averaged Chowla and Elliott conjectures for two-point correlations, *Forum Math. Pi* 4 (2016) e8.
23. J. B. Tunnell, A classical Diophantine problem and modular forms of weight $3/2$, *Invent. Math.* 72 (1983) 323–334.
24. Y. Zhang, Bounded gaps between primes, *Ann. of Math.* 179 (2014) 1121–1174.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.