



# COMPUTATIONAL MATHEMATICS

Based on prof. Antonio Frangioni's and  
prof. Federico Poloni's lectures

First draft provided by Gemma Martini

August 30, 2019

# Contents

<b>1 19th of September 2018 — A. Frangioni</b>	<b>2</b>
1.1 Introduction to machine learning problems . . . . .	2
1.2 Optimization . . . . .	3
1.2.1 Linear estimation . . . . .	3
1.2.2 Low-rank approximation . . . . .	4
1.2.3 Support vector machines . . . . .	4
<b>2 20th of September 2018 — F. Poloni</b>	<b>6</b>
2.1 A warm up . . . . .	6
2.2 Orthogonality . . . . .	8
<b>3 21st of September 2018 — A. Frangioni</b>	<b>9</b>
3.1 Mathematical background for optimization problems . . . . .	9
<b>4 26th of September 2018 — F. Poloni</b>	<b>13</b>
<b>5 27th of September 2018 — A. Frangioni</b>	<b>18</b>
5.1 Continuity . . . . .	19
5.2 Derivatives . . . . .	21
5.2.1 Multivariate differentiability . . . . .	22
<b>6 18th of October 2018 — F. Poloni</b>	<b>25</b>
6.1 Least squares problem . . . . .	25
6.1.1 Method of normal equations . . . . .	26
6.2 QR factorization . . . . .	27
<b>7 19th of October 2018 — A. Frangioni</b>	<b>30</b>
7.1 Gradient method for quadratic functions . . . . .	30
7.2 MatLab implementation . . . . .	32
7.2.1 Error . . . . .	33
<b>8 24th of October 2018 — A. Frangioni</b>	<b>34</b>
8.1 Gradient method for non quadratic functions . . . . .	34
8.1.1 How fast does it converge? . . . . .	34
8.1.2 Exact line search, first orderd approach . . . . .	36
<b>9 25th of October 2018 — A. Frangioni</b>	<b>40</b>
9.0.1 Line search: second order approaches . . . . .	40
9.0.2 Exact line search: zeroth-order approaches . . . . .	42
9.0.3 Inexact line search: Armijo-Wolfe . . . . .	43

<b>10 26th of October 2018 — F. Poloni</b>	<b>49</b>
10.1 How to construct a QR factorization . . . . .	49
10.1.1 Matlab implementation . . . . .	51
10.2 How to use the thin QR factorization to solve a least squares problem . . . . .	53
<b>11 7th of November 2018 — F. Poloni</b>	<b>55</b>
11.1 Least squares problem with SVD . . . . .	55
11.1.1 Behaviour in case of zeros as singular values . . . . .	58
11.2 Truncated SVD . . . . .	59
11.3 Tikhonov regularization / ridge regression . . . . .	59
<b>12 8th of November 2018 — A. Frangioni</b>	<b>61</b>
12.1 Good practice of designing a project . . . . .	61
<b>13 9th of November 2018 — F. Poloni</b>	<b>62</b>
13.1 Conditioning . . . . .	62
13.1.1 Conditioning of linear systems . . . . .	65
13.2 Condition number, SVD, and distance to singularity . . . . .	66
13.3 Conditioning of least squares problem . . . . .	68
<b>14 14th of November 2018 — A. Frangioni</b>	<b>71</b>
14.1 Taylor method . . . . .	71
14.1.1 Interpretation of Newton method . . . . .	74
14.1.2 Non convex case . . . . .	74
<b>15 15th of November 2018 — F. Poloni</b>	<b>76</b>
15.1 Stability of algorithms . . . . .	76
15.2 Backward stability of QR factorization . . . . .	80
15.2.1 Stability of algorithms for least-squares problems . . . . .	81
15.2.2 A posteriori checks . . . . .	82
15.3 A posteriori check for Least Squares Problems . . . . .	84
<b>16 16th of November 2018 — A. Frangioni</b>	<b>86</b>
16.1 Quasi-newton methods . . . . .	86
16.1.1 Davidon-Fletcher-Powell . . . . .	87
16.1.2 Broyden-Fletcher-Goldfarb-Shanno . . . . .	88
16.1.3 Poorman's approach . . . . .	88
16.2 Conjugate gradient method . . . . .	88
<b>17 21st of November 2018 — F. Poloni</b>	<b>91</b>
17.1 Gaussian elimination and LU factorization . . . . .	92
17.1.1 Stability of LU . . . . .	95
17.1.2 Gaussian elimination on sparse matrices . . . . .	96

<b>18</b>	<b>22nd of November 2018 — A. Frangioni</b>	<b>98</b>
18.1	Deflected gradient methods . . . . .	98
18.1.1	Heavy ball gradient method . . . . .	98
18.1.2	Accelerated gradient . . . . .	99
18.2	Incremental gradient methods . . . . .	100
18.3	Subgradient methods . . . . .	101
<b>19</b>	<b>23rd of November 2018 — F. Poloni</b>	<b>104</b>
19.1	Gaussian elimination on symmetric matrices . . . . .	104
19.2	Cholesky factorization . . . . .	107
19.3	Krylov subspace methods . . . . .	107
<b>20</b>	<b>28th of November 2018 — A. Frangioni</b>	<b>109</b>
20.1	Subgradient methods . . . . .	109
20.1.1	Target level stepsize . . . . .	111
20.2	Deflected subgradient . . . . .	112
20.3	Smoothed gradient methods . . . . .	113
20.4	Cutting-plane algorithm . . . . .	115
20.5	Bundle methods . . . . .	117
<b>21</b>	<b>30th of November 2018 — A. Frangioni</b>	<b>118</b>
21.1	Constrained optimization . . . . .	118
21.1.1	Linear equality constraints . . . . .	118
21.1.2	Background for linear inequality constraints . . . . .	120
<b>22</b>	<b>6th of December 2018 — A. Frangioni</b>	<b>126</b>
22.1	Duality . . . . .	126
22.2	Lagrangian duality . . . . .	127
22.3	Specialized dual . . . . .	129
22.3.1	Linear programs . . . . .	129
22.3.2	Quadratic programs . . . . .	129
22.3.3	Conic program . . . . .	129
22.4	Fenchel's duality . . . . .	131
<b>23</b>	<b>12th of December 2018 — A. Frangioni</b>	<b>132</b>
23.1	Quadratic problem with linear equality constraints . . . . .	132
23.2	Inequality constrained problems . . . . .	133
23.2.1	Projected gradient method . . . . .	134
23.2.2	Projected gradient method with box constraints . . . . .	137
23.2.3	Active-set method for quadratic programs . . . . .	138
23.2.4	Frank-Wolfe method . . . . .	139

<b>24 14th of December 2018 — A. Frangioni</b>	<b>141</b>
24.1 Dual methods for linear constrained optimization . . . . .	141
24.1.1 Separable problems and partial dual . . . . .	141
24.2 Primal/dual methods or barrier methods . . . . .	142
24.2.1 Barrier function and central path . . . . .	142
24.3 Primal-dual interior point method . . . . .	144

# 1 19th of September 2018 — A. Frangioni

This course will deal with the optimization and numerical analysis of machine learning problems. We are not going to solve difficult problems (e.g. NP-hard problems), besides we try to find an efficient solution for simple ones (often **convex** ones), since we are dealing with huge amount of data.

Let us start with a warm up on machine learning problems.

## 1.1 Introduction to machine learning problems

Machine learning techniques are not as “young” as it might seem, the intuition has been there for ages, but we did not have enough calculus power. Machine learning algorithms are starting working well nowadays, thanks to the many improvements in computer performances; for this reason, it is becoming a more and more popular subject to study.

The main idea behind machine learning is to take a huge amount of data (e.g. frames of a video for object-recognition) and squeeze them, in order to process them. This intuitive concept is translated in mathematical terms as “building a **model**” that fits our data. As in practical engineering problems, people want to construct a model (a small sized representation of the large thing we want to produce in the end) and try to understand its behaviour, before actually build the thing. Take as an example the problem of designing a jet. It is not clever to start building the plane before designing a cheap prototype to better study its behaviour in the atmosphere.

The kind of models we want to build are cheap to construct and as close as possible to the real problem we are studying. In phisics, people try to find the best mathematical model to describe a real world phenomenon. The main issue is computation, since the more accurate the model, the more costly the prediction phase. Hence, a model is a good when it is a good tradeoff between accuracy and simplicity, namely it provides good prediction without incurring in slow computations.

The model, though, has to be parametric: we do not have only one model, we have a “shape” of a model, which is fit to our problem through the tuning of some parameters.

**Example 1.1.** As an example, we are given three couples:  $f(x_1) = y_1$ ,  $f(x_2) = y_2$ ,  $f(x_3) = y_3$ , as shown in Figure 1.1.

We need to make some choices: first, we need to decide the kind of model we believe is a good approximation of the objective function, say a linear model  $f(x) = ax + b$ . After doing that, we are left with choosing its parameters (in order to pick a line among the whole family of functions), namely  $a$  and  $b$ .

The aim of machine learning is to build a model that fits the data we are given and then to tune parameters in order to achieve a good “predicting power” on unseen inputs (in machine learning the technical term is “not **overfitting**”, see ??).

Another important characteristic of a good model is that it should not take too long to be built.




FIGURE 1.1: Geometric representation of the input. We are interested in finding a model that fits the input data and allows to predict  $\bar{y}$  out of  $\bar{x}$ .

In this course we do not concentrate on the problem of finding the model that best fits our data, but we are already given a problem and a model and we only study its behaviour through its parameters.

## 1.2 Optimization

In the rest of this lecture we are going to understand what an optimization problem is, through some intuitive real world examples.

### 1.2.1 Linear estimation

**Definition 1.1** (Linear model). *Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be the objective function. We call  $\tilde{f}(x) = \sum_{i=1}^n w_i x_i + w_0 = wx + w_0$  the **linear model** of  $f$  for a given set of parameters, which is a vector  $w = (w_0, w_1, \dots, w_n) \in \mathbb{R}^{n+1}$ .*

How can we evaluate the “similarity” between our model and the objective function? Through computing the “error” or difference between the objective function and the model on each input. Under this assumption, the error function may be used to find the best parameters for our model, through a minimum problem:

**Definition 1.2** (Least squares problem). *Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be the objective function, such that  $f(x) = y$  and let  $Xw$  be our linear model. Then we can find the best values for vector  $w \in \mathbb{R}^{n+1}$  by computing*

$$\min_w \|y - Xw\|$$

If the matrix  $X$  is invertible then the simple solution is  $w = X^{-1}y$ . The point is that this operation is very costly when dealing with a huge number of entries.

### 1.2.2 Low-rank approximation

We may want to approximate a matrix  $M \in M(n, m, \mathbb{R})$  as the product between two smaller matrices:  $A \in M(n, k, \mathbb{R})$  and a “fat and large”  $B \in M(k, m, \mathbb{R})$  ( $k \ll n, m$ ).

$$\boxed{M} \approx \boxed{A} \cdot \boxed{B}$$

This problem can be translated into a numerical analysis problem of the following shape

$$\min_{A, B} \|M - AB\|$$

### 1.2.3 Support vector machines

Let us take a decision problem: given a set of values of many parameters (aka variables) “label” a person as ill or healthy.

The geometric intuition in two dimensions is given by Figure 1.2. We would like to find the line that better splits the plane into two regions. The rationale here is to maximize the space between the line and the nearest points (called **margin**), in order to have a better accuracy.




FIGURE 1.2: There are many possible boundaries that can be chosen as a model using many angular coefficients. Our best guess is the one that maximizes the distance between the line and the nearest points.

The maximum-margin separating hyperplane is the solution of

$$\min_w \{\|w_+\|^2 : y^i(w^T x^i + w_0) \geq 1, i = 1, \dots, m\}$$

where the margin is  $\frac{2}{\|w_+\|}$ , assuming any exists.

In fact, what happens most of the times is that there is no such line. To overcome this issue we introduce the concept of “penalty” that accounts for the number of points that are misclassified.

**Definition 1.3** (Multi-objective optimization problem).

$$\min_{w, \xi} \|w_+\|^2 + C \sum_{i=1}^m \xi_i$$

where  $y^i(w_+x^i + w_0) \geq 1 - \xi_i$ ,  $\forall i = 1, \dots, m$  and  $x_i \geq 0$ ,  $\forall i = 1, \dots, m$ , where  $C$  is called **hyperparameter**.

This formula formalizes the intuition that the approximated function may have a greater norm and lead to a very small misclassification error, or it could be the other way round. Both these solutions are acceptable and their performances depend only on the problem.

This whole course has the aim of presenting some techniques for solving efficiently **convex quadratic problems**, as the ones presented above.

Whenever we are able to solve the multiobjective optimization problem we are also able to solve what is called the **dual problem**, which is formally defined in ?? and has the following shape in our case:

$$\max_{\alpha} \left\{ \sum_{i=1}^m \alpha_i - \frac{1}{2} \sum_{i=1}^m \sum_{j=1}^m \alpha_i \langle x^i, x^j \rangle \alpha_j \right\}$$

where  $\sum_{i=1}^m y^i \alpha_i = 0$  and  $0 \leq \alpha_i \leq C$ ,  $\forall i = 1, \dots, m$ .

## 2 20th of September 2018 — F. Poloni

### 2.1 A warm up

#### Do you recall?

Let  $x, y \in \mathbb{R}^n$ . The product between those two vectors is computed as follows  $x^T y = \sum_{i=1}^n x_i y_i$  and  $x^T y \in \mathbb{R}$ .

Let  $x \in \mathbb{R}^n$  and  $\lambda \in \mathbb{R}$  we call **multiple** of vector  $x$  the following:  $\lambda x = x\lambda = \begin{pmatrix} \lambda x_1 \\ \vdots \\ \lambda x_n \end{pmatrix}$ .

Given a matrix  $A \in M(n, m, \mathbb{R})$  and a vector  $b \in \mathbb{R}^m$  the **matrix-vector product**  $Ab = v \in \mathbb{R}^n$  is computed as follows:

$$v = Ab = \begin{pmatrix} A_1 b \\ A_2 b \\ \vdots \\ A_m b \end{pmatrix} = \sum_{j=1}^m A^j b_j$$

The computational complexity of this operation is  $O(n^2)$ .

We call **image** of a matrix  $A$  ( $\text{Im}(A)$ ) the set of vectors that can be obtained multiplying  $A$  by any vector in the domain of  $A$ .

On the other hand, we call **kernel** of a matrix  $A$  ( $\text{ker}(A)$ ) the set of vectors  $w$  in its domain such that  $Aw = 0$ .

Given two matrices  $A \in M(n, m, \mathbb{R})$  and  $B \in M(m, k, \mathbb{R})$  we call **matrix-matrix product** the following:  $C = AB$  such that  $C_{ij} = A_i B^j$ , where  $A_i^T \in \mathbb{R}^m$  is the  $i$ -th row of  $A$ ,  $B^i$  is the  $i$ -th column of  $B$  ( $B^i \in \mathbb{R}^m$ ) and  $C \in M(n, k, \mathbb{R})$ . The computational complexity of this operation is  $O(n^3)$ . Notice that this product is **not commutative**.

The matrix-matrix product also works on “matrices” made of one column only (vector of  $\mathbb{R}^n$ ), but in this case a row of the right-side matrix is made by only one scalar...

Given a matrix  $A \in M(n, \mathbb{R})$  we call **inverse** of  $A$  the matrix  $A^{-1}$  such that  $A^{-1}A = AA^{-1} = I_n$ .

The **transpose** of a matrix  $A \in M(n, m, \mathbb{R})$  is  $A^T$  such that  $A_{ij}^T = A_{ji}$

The **inverse of a product** (shoe-sock identity) is  $(AB)^{-1} = B^{-1}A^{-1}$ . Notice that this identity holds only for square matrices.

The **transpose of a product** (shoe-sock identity) is  $(AB)^T = B^T A^T$ .

The objective of this course, for the part concerning numerical methods, is solving linear systems efficiently.

**Definition 2.1** (Linear system). *Let  $A \in M(n, m, \mathbb{R})$ ,  $b \in \mathbb{R}^n$  and  $x \in \mathbb{R}^m$ . We term **linear system** the following:*

$$Ax = b$$

Our goal is to approximate such vector  $x$ , hence resulting in solving a minimum problem:

$$\min \|Ax - b\|$$



### Something on Matlab ...

Notice that the machine precision is  $10^{-16}$ , so we should pay attention when making computations, since we may incur in some error (proportional to the size of the operands).

In Matlab a matrix is written as  $A=[1, 2, 3; 4, 5, 6]$ ;, where  $[1, 2, 3]$  is the first row of the matrix A.

The transpose of a matrix or a vector is denoted by  $A'$ .

If we are interested in only a part of our matrix A we may write  $A[1:2, 1:3]$  and obtain only the rows of A that go from 1 to 2 and those columns from 1 to 3.

**Fact 2.1.** Let  $A \in GL(n, \mathbb{R})$  (aka A is a real square matrix of size n and invertible),  $B, C \in M(n, m, \mathbb{R})$ .

If  $AB = AC$  then  $B = C$ .

**Definition 2.2** (Block multiplications). Let  $A \in M(n, m, \mathbb{R})$  and let  $B \in M(m, k, \mathbb{R})$ . We can compute the result of a block of the matrix  $AB$  as the product of the two blocks in A and B in the corresponding position.

**Fact 2.2** (Block triangular matrices). Let  $M \in M(n, m, \mathbb{R})$  and  $N \in M(m, k, \mathbb{R})$  such that they are **block triangular**. Their product is a block triangular matrix as well.

$$MB = \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \begin{pmatrix} E & F \\ 0 & G \end{pmatrix} = \begin{pmatrix} AE & BF \\ 0 & DG \end{pmatrix}$$

**Fact 2.3** (Properties of triangular matrices).

1. A block triangular matrix is invertible iff its blocks are invertible;
2. The eigenvalues of a block triangular matrix are the union of the eigenvalues of each block;
3. Let  $M \in GL(n, m, \mathbb{R})$  such that  $M = \begin{pmatrix} A & B \\ 0 & D \end{pmatrix}$  the inverse of M is

$$M^{-1} = \begin{pmatrix} A^{-1} & -A^{-1}BD^{-1} \\ 0 & B^{-1} \end{pmatrix}.$$

Why are we interested in block triangular matrices? They depict a situation as shown in Figure 2.1.

manca  
figura dei  
blocchi  
in latex




FIGURE 2.1: The adjacency matrix of a bipartite graph has 0s in its bottom left part (Matlab syntax  $A[p+1:n; 1:p] = 0$ ).

## 2.2 Orthogonality

**Definition 2.3** (Norms). Let  $x \in \mathbb{R}^n$ . We “measure” their magnitude using so-called “norms”.

$$\text{EUCLIDEAN: } \|x\|_2 = x^T x = \sqrt{\sum_{i=1}^n x_i^2};$$

$$\text{NORM 1: } \|x\|_1 = \sum_{i=1}^n |x_i|;$$

$$p\text{-NORM: } |x|_p = \left( \sum_{i=1}^n |x_i|^p \right)^{1/p};$$

$$0\text{-NORM: } \|x\|_0 = |\{i : |x_i| > 0\}|;$$

$$\infty\text{-NORM: } \|x\|_\infty = \max_{i=1,\dots,n} |x_i|.$$

From now on in this part of the course we will refer to norm-2 only.

**Definition 2.4** (Orthogonal matrix). Let  $A \in M(n, m, \mathbb{R})$ . We call  $A$  **orthogonal** if  $\forall x \in \mathbb{R}^n \quad \|Ax\| = \|x\|$ .

**Fact 2.4** (Equivalent definition of orthogonal matrix). Let  $A \in M(n, \mathbb{R})$ .  $A$  is orthogonal iff  $A^T A = A A^T = I_n$ , where  $I_n$  is the identity matrix of size  $n$  (1 on the diagonal, 0 elsewhere).

### 3 21st of September 2018 — A. Frangioni

#### 3.1 Mathematical background for optimization problems

**Definition 3.1** (Minimum problem). Let  $X$  be a set, called **feasible region** and let  $f : X \rightarrow \mathbb{R}$  be any function, called **objective function** we call **problem** the following

$$(P) \quad f_* = \min\{f(x) : x \in X\}$$

**Definition 3.2** (Feasible solution). Let  $x \in F$  be a solution of the minimum problem in which the domain is a superset of  $X \subset F$ . We say that  $x$  is a **feasible solution** if  $x \in X$ . On the other hand,  $x$  is **unfeasible** if  $x \in F \setminus X$ .

**Definition 3.3** (Optimal solution). Under the same hypothesis of the above definition, we define  $x_*$  such that  $f(x_*) = f_*$  an **optimal solution**, where  $f_* \leq f(x) \forall x \in X, \forall v > f_* \exists x \in X$  s.t.  $f(x) < v$ .

It is possible to find problems where there is no optimal solution at all.

**Example 3.1.** There are two cases in which it is not possible to find an optimal solution:

1. The domain is empty, which may be not trivial to prove, since it is an NP-hard problem sometimes;
2. We want to find the minimum of the objective function but it is unbounded below ( $\forall M \exists x_M \in X$  s.t.  $f(x_M) \leq M$ ). On the other hand, we need to maximize the function, but it is unbounded above.

We can now rewrite the problem of solving an optimization problem as:

1. Finding  $x_*$  and proving it is optimal
2. Or proving  $X = \emptyset$
3. Or constructively prove  $\forall M \exists x_M \in X$  s.t.  $f(x_M) \leq M$ .

Most of the times we consider optimal a solution which is close to the true optimal value, modulo some error.

**Definition 3.4** (Absolute error). We call **absolute error** the gap between the real value and the one we obtained. Formally,

$$f(\bar{x}) - f_* \leq \varepsilon$$

**Definition 3.5** (Relative error). We define as **relative error** the absolute error, normalized by the true value of the function

$$(f(\bar{x}) - f_*) / |f_*| \leq \varepsilon$$

Let us consider an iterative algorithm that moves towards the optimum. It may happen that the function decreases and decreases along a certain direction but its non-continuity leads to the impossibility of reaching the optimum. As an example, let us take the following

$$f(x) = \begin{cases} x & \text{if } x > 0 \\ 1 & \text{if } x = 0 \end{cases}$$

**Definition 3.6** (Totally ordered set). *We say that set  $X$  is **totally ordered** if  $\forall x, y \in X$ , either  $f(x) \leq f(y)$  or  $f(y) \leq f(x)$ .*

**Definition 3.7** (Infima and suprema). *Given a totally ordered set  $R$  and one of its subsets (say  $S \subseteq R$ )*

*$s$  is the infimum of  $S \Leftrightarrow s = \inf S \Leftrightarrow s \leq s \ \forall s \in S \wedge \forall t > s \exists s \in S \text{ s.t. } s \leq t$*

*$s$  is the supremum of  $S \Leftrightarrow \bar{s} = \sup S \Leftrightarrow \bar{s} \geq s \ \forall s \in S \wedge \forall t < \bar{s} \exists s \in S \text{ s.t. } s \geq t$*

What happens if we have more than one objective function? We are provided with two tools in order to reduce them into one:

SCALARIZATION: using a linear combination of the two functions:  $f(x) = \alpha f_1(x) + \beta f_2(x)$ ;

BUDGETING:  $f(x) = f_1(x)$ ,  $X := X \cup \{ f_2(x) \leq b \}$ , which intuitively corresponds to taking into account only one objective function, provided that the values of the other functions are not too high.

**Definition 3.8** (Extended real). *In the case of unbounded functions the value of infima or suprema are  $\infty$ , and we call **extended reals**  $\overline{\mathbb{R}} = -\infty \cup \mathbb{R} \cup +\infty$ .*

We are interested in studying sequences, because iterative methods start from a certain point and move towards the optimal, hopefully.

**Definition 3.9** (Limit). *Given a sequence  $\{x_i\}$  the **limit** for  $i \rightarrow \infty$  is defined as*

$$\lim_{i \rightarrow \infty} v_i = v \iff \forall \varepsilon > 0 \ \exists h \text{ s.t. } |v_i - v| \leq \varepsilon \ \forall i \geq h$$

It may happen that a sequence has or does not have a limit. For example  $\{\frac{1}{n}\}$  has limit 0 for  $n \rightarrow +\infty$ , while  $\{(-1)^n\}$  does not.

**Fact 3.1.** *Let us be given a monotone sequence, then the sequence **does** have a limit.*

Notice that given a sequence either it is monotone or it can be “split” into two monotone sequences (for example  $\{(-1)^n\}$  can be transformed into  $\{(-1)^{2n}\}$  and  $\{(-1)^{2n+1}\}$  and these two sequences are both monotone).

**Definition 3.10** (Euclidean vector space). *We call **Euclidean space***

$$\mathbb{R}^n := \left\{ \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} : x_i \in \mathbb{R}, i = 1, \dots, n \right\}$$

Equivalently, we can characterize the Euclidean space as Cartesian product of  $\mathbb{R}$   $n$  times:  
 $\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \dots \mathbb{R}$ .

The main operations on elements of the Euclidean space (vectors) are:

$$\text{SUM: } x + y := \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_n \end{pmatrix}$$

$$\text{SCALAR MULTIPLICATION: } \alpha x = \begin{pmatrix} \alpha x_1 \\ \alpha x_2 \\ \vdots \\ \alpha x_n \end{pmatrix}$$

In order to be able to compute limits in a vector space we need to use norms (see Definition 2.3).

**Fact 3.2.** *The norms on a vector space have the following properties:*

1.  $\|x\| \geq 0$  and  $\forall x \in \mathbb{R}^n$ ,  $\|x\| = 0 \iff x = 0$ ;
2.  $\|\alpha x\| = |\alpha| \|x\|$ ,  $\forall x \in \mathbb{R}^n$ ,  $\alpha \in \mathbb{R}$ ;
3.  $\|x + y\| \leq \|x\| + \|y\|$ ,  $\forall x, y \in \mathbb{R}^n$  (triangle inequality).

**Definition 3.11** (Ball). *We term **ball** centered in  $\bar{x}$  and having  $\varepsilon$  as radius as the set of points that are close enough to  $x \in \mathbb{R}^n$ :  $B(\bar{x}, \varepsilon) = \{x \in \mathbb{R}^n : \|x - \bar{x}\| \leq \varepsilon\}$ .*

In Figure 3.1 we may observe the different shapes of the same ball varying the value of  $p$  in the  $p$ -norm.

**Definition 3.12** (Scalar product). *Let  $x, y \in \mathbb{R}^n$  we define the **scalar product** between these two vectors*

$$\langle x, y \rangle := y^T x = \sum_{i=1}^n x_i y_i = x_1 y_1 + \dots + x_n y_n$$

**Fact 3.3.** *A scalar product has the following properties:*

1.  $\langle x, y \rangle = \langle y, x \rangle \quad \forall x, y \in \mathbb{R}^n$  (symmetry)




FIGURE 3.1: The shapes of balls centered in the origin of radius 1 varying the value of  $p$ .

$$2. \langle x, x \rangle \geq 0, \forall x \in \mathbb{R}^n, \langle x, x \rangle = 0 \iff x = 0;$$

$$3. \langle \alpha x, y \rangle = \alpha \langle x, y \rangle, \forall x \in \mathbb{R}^n, \alpha \in \mathbb{R};$$

$$4. \langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle, \forall x, y, z \in \mathbb{R}^n.$$

**Fact 3.4** (Cauchy-Schwartz inequality). *Let  $x, y \in \mathbb{R}^n$ . The following holds:*

$$\langle x, y \rangle^2 \leq \langle x, x \rangle \langle y, y \rangle \equiv |\langle x, y \rangle| \leq \|x\| \|y\|, \forall x, y \in \mathbb{R}^n$$

An important characterization of the scalar product is the one that uses angles:

$\langle x, y \rangle = \|x\| \|y\| \cos \theta$ :  $x \perp y \iff \langle x, y \rangle = 0$  and  $\langle x, y \rangle > 0 \iff$  “ $x$  and  $y$  point in the same direction”

We have now all the tools to define the notion of limit of a sequence in  $\mathbb{R}^n$ .

**Definition 3.13** (Limit of a sequence in the Euclidean space). *Let  $\{x_i\} \subset \mathbb{R}^n$  be a sequence in  $\mathbb{R}^n$ . The **limit** of  $\{x_i\}$  for  $i \rightarrow +\infty$  is the following:*

$$\begin{aligned} \lim_{i \rightarrow \infty} x_i = x &\equiv \{x_i\} \rightarrow x \\ &\Updownarrow \\ \forall \varepsilon > 0 \exists h \text{ s.t. } d(x_i, x) \leq \varepsilon \quad \forall i \geq h & \\ &\Updownarrow \\ \forall \varepsilon > 0 \exists h \text{ s.t. } x_i \in \mathcal{B}(x, \varepsilon) \quad \forall i \geq h & \\ &\Updownarrow \\ \lim_{i \rightarrow \infty} d(x_i, x) &= 0 \end{aligned}$$

## 4 26th of September 2018 — F. Poloni

In the previous lecture we introduced some sufficient conditions for matrix orthogonality.

**Theorem 4.1.** *Let  $U \in M(n, \mathbb{R})$  be an orthogonal matrix and let  $x \in \mathbb{R}^n$ . Then  $\|Ux\| = \|x\|$ .*

*Proof.*  $\|Ux\| = (Ux)^T \cdot (Ux) \stackrel{(1)}{=} x^T U^T U x = x^T I_n x = x^T x = \|x\|$   
 where  $\stackrel{(1)}{=}$  follows from the definition of transpose of a product.  $\square$

Geometrically, an orthogonal matrix represents a symmetry or a rotations and these operations do not alter the size of vectors.

**Definition 4.1** (Orthonormality). *Let  $x, y \in \mathbb{R}^n$  we say that  $x$  and  $y$  are **orthonormal** if  $\langle x, y \rangle = 0$  and  $\|x\| = \|y\| = 1$ .*

**Fact 4.2.** *Let us take  $U \in M(n, \mathbb{R})$  such that  $U$  is orthogonal. Then its columns  $U^1, U^2, \dots, U^n$  are **orthonormal** and the same holds for its rows.*

$$U^{i^T} U^j = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

and

$$U_i U_j^T = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{otherwise} \end{cases}$$

**Fact 4.3.** *Let  $U, V \in M(n, \mathbb{R})$ , such that  $U$  and  $V$  are orthogonal, then  $U \cdot V$  is orthogonal.*

*Proof.*  $(UV)^T \cdot (UV) = V^T U^T UV = V^T I_n V = V^T V = I_n$   $\square$

**Definition 4.2** (Eigenvectors and eigenvalues). *Let  $A \in M(n, \mathbb{R})$  and let  $x \in \mathbb{R}^n$  and  $\lambda \in \mathbb{R}$ . If  $Ax = \lambda x$  we say that  $x$  is an **eigenvector** of **eigenvalue**  $\lambda$ .*

**Fact 4.4.** *Let  $A \in T(n, \mathbb{R})$  (real triangular matrix). The eigenvalues of  $A$  are the scalars on the diagonal.*

Under some conditions we may obtain a **diagonal form** of  $A$ , namely:

$$A = V \Lambda V^{-1} = \begin{pmatrix} V^1 & V^2 & \dots & V^n \end{pmatrix} \cdot \begin{pmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{pmatrix} \cdot \begin{pmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{pmatrix}$$

where  $\forall i = 1, \dots, n$   $v_i$  are eigenvectors of  $A$  of eigenvalue  $\lambda_i$ .

$$\begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

This is due to the fact that  $\forall B \in R^n$  s.t.  $Vx = b$ ,  $x =$

position  $i$  and so  $AV^i = V\Lambda V^{-1}V^i = V\Lambda e_i = \lambda_i V^i$ .

Another way to see the diagonalized form of  $A$  is the following:

$$\begin{aligned} A &= V\Lambda V^{-1} = \sum_{i=1}^n V^i \lambda_i w_i \\ &= \boxed{V^1} \cdot \boxed{\lambda_1} \cdot \boxed{w_1} + \boxed{V^2} \cdot \boxed{\lambda_2} \cdot \boxed{w_2} + \cdots + \boxed{V^n} \cdot \boxed{\lambda_n} \cdot \boxed{w_n} \end{aligned} \quad (4.1)$$



### Something on Matlab ...

Notice that in Matlab the eigenvalues and eigenvectors of a matrix are computed using the command `[V, Lambda] = eig(U)` and this operation has a computational complexity of  $O(n^3)$ .

Notice that not all matrices  $A \in M(n, \mathbb{R})$  allow a diagonal decomposition. It may happen that such a matrix is diagonalizable in  $\mathbb{C}$  and its eigenvalues are complex.

**Fact 4.5.** If  $A \in M(n, \mathbb{R})$  is diagonalizable (aka may be written as  $A = V\Lambda V^{-1}$ )  $A^k x = \sum_{i=1}^n \lambda_i^k \alpha_i V^i$ , for some  $\alpha_i \in \mathbb{R}$ .

*Proof.* ALGEBRAIC VIEW POINT: Let us write  $x$  in the base of  $R^n$  made of the linearly independent columns of  $V$ :

$$x = V^1 \alpha_1 + V^2 \alpha_2 + \cdots + V^n \alpha_n$$

for some  $\alpha_i \in \mathbb{R}$ .

$$\begin{aligned} Ax &= A \cdot (V^1 \alpha_1 + V^2 \alpha_2 + \cdots + V^n \alpha_n) \\ &= AV_1 \alpha_1 + AV_2 \alpha_2 + \cdots + AV_n \alpha_n \\ &= \lambda_1 V^1 \alpha_1 + \lambda_2 V^2 \alpha_2 + \cdots + \lambda_n V^n \alpha_n \\ &= V^1 (\lambda_1 \alpha_1) + V^2 (\lambda_2 \alpha_2) + \cdots + V^n (\lambda_n \alpha_n) \end{aligned} \quad (4.2)$$

Then

$$\begin{aligned}
A^2x &= A \cdot (V^1(\lambda_1\alpha_1) + V^2(\lambda_2\alpha_2) + \cdots + V^n(\lambda_n\alpha_n)) \\
&= AV^1\lambda_1\alpha_1 + AV^2\lambda_2\alpha_2 + \cdots + AV^n\lambda_n\alpha_n \\
&= \lambda_1^2V^1\alpha_1 + \lambda_2^2V^2\alpha_2 + \cdots + A\lambda_n^2V^n\alpha_n
\end{aligned} \tag{4.3}$$

Inductively, we have proved the theorem.

LINEAR ALGEBRA VIEW POINT:

$$\begin{aligned}
A^kx &= A \cdot A \cdot \dots \cdot A \cdot x \\
&= V\Lambda V^{-1}V\Lambda V^{-1}\dots V\Lambda V^{-1}x \\
&= V\Lambda^k V^{-1}x \\
&= V \begin{pmatrix} \lambda_1^k & & & \\ & \lambda_2^k & & \\ & & \ddots & \\ & & & \lambda_n^k \end{pmatrix} V^{-1}x \\
&= V \begin{pmatrix} \lambda_1^k & & & \\ & \lambda_2^k & & \\ & & \ddots & \\ & & & \lambda_n^k \end{pmatrix} \cdot \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_n \end{pmatrix}
\end{aligned} \tag{4.4}$$

□

Thanks to this proposition we can prove the following

**Theorem 4.6.** Let  $A \in M(n, \mathbb{R})$ . If  $\forall \lambda_i$  eigenvalues of  $A$   $|\lambda_i| < 1$  then  $\lim_{k \rightarrow \infty} A^kx = 0$ .

**Theorem 4.7.** Let  $A \in M(n, \mathbb{R})$ . If  $\forall \lambda_i$  eigenvalues of  $A$   $|\lambda_i| < |\lambda_1|$  then  $A^kx \approx V^1\lambda_1^k\alpha_1$ .

**Fact 4.8.** Let  $A \in M(n, \mathbb{R})$  be a diagonalizable matrix and let

$$A = V\Lambda V^{-1} = \begin{pmatrix} V^1 & V^2 & \dots & V^n \end{pmatrix} \cdot \begin{pmatrix} \lambda_1 & & & \\ & \lambda_2 & & \\ & & \ddots & \\ & & & \lambda_n \end{pmatrix} \cdot \begin{pmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{pmatrix}$$

Let us now consider a reordering of  $V$ 's columns and apply the same permutations to the “diagonal vector” of  $\Lambda$  such that

$$\hat{V} = \begin{pmatrix} V^2 & V^1 & V^3 \dots & V^n \end{pmatrix} \text{ and } \hat{\Lambda} = \begin{pmatrix} \lambda_2 & & & \\ & \lambda_1 & & \\ & & \ddots & \\ & & & \lambda_n \end{pmatrix}$$

$A$  can be diagonalized through such  $\hat{V}$  and  $\hat{\Lambda}$ :  $A = V\Lambda V^{-1} = \hat{V}\hat{\Lambda}\hat{V}^{-1}$ .

Moreover, in the case of repeated eigenvalues

**Fact 4.9.** Let  $A \in M(n, \mathbb{R})$  a diagonalizable matrix such that  $A = V\Lambda V^{-1}$ , where  $\lambda_1 = \lambda_2$  (without loss of generality). Then  $V$  can be replaced by  $\tilde{V} = \begin{pmatrix} V^1 + V^2 & V^1 - V^2 & V^3 & \dots & V^n \end{pmatrix}$ .

**Theorem 4.10** (Spectral theorem). Let  $A \in S(n, \mathbb{R})$  ( $A$  is a real symmetric matrix). Then  $A$  is diagonalizable, its eigenvalues are real numbers and  $V$  is an orthogonal matrix.

**Fact 4.11.** Let  $A \in S(n, \mathbb{R})$  and let  $x \in \mathbb{R}^n$ .  $\lambda_{\min}\|x\|^2 \leq x^T Ax \leq \lambda_{\max}\|x\|^2$ , where  $\lambda_{\max}$  and  $\lambda_{\min}$  are respectively the eigenvalue of maximum value and the eigenvalue of minimum value.

*Proof.* DIAGONAL  $A$ :  $x^T Ax = x^T \cdot \begin{pmatrix} \lambda_2 & & & & \\ & \lambda_1 & & & \\ & & \lambda_3 & & \\ & & & \ddots & \\ & & & & \lambda_n \end{pmatrix} \cdot x = \lambda_1 x_1^2 + \lambda_2 x_2^2 + \dots + \lambda_n x_n^2$

It is obvious that this sum is bounded by:

$$\lambda_{\min} \cdot (x_1^2 + x_2^2 + \dots + x_n^2) \leq \lambda_1 x_1^2 + \lambda_2 x_2^2 + \dots + \lambda_n x_n^2 \leq \lambda_{\max} \cdot (x_1^2 + x_2^2 + \dots + x_n^2)$$

The following holds:  $\lambda_{\min} \cdot (x_1^2 + x_2^2 + \dots + x_n^2) = \lambda_{\min} \cdot x^T x = \lambda_{\min} \cdot \|x\|^2$  and, on the other hand,  $\lambda_{\max} \cdot (x_1^2 + x_2^2 + \dots + x_n^2) = \lambda_{\max} \cdot x^T x = \lambda_{\max} \cdot \|x\|^2$  and this proves the fact in the special case of diagonal matrix  $A$ .

GENERAL CASE: Let us represent  $A$  through its eigendecomposition:  $A = U\Lambda U^{-1} = U\Lambda U^T$ , where  $U$  is an orthogonal matrix.

$$x^T Ax = x^T U \Lambda U^T x \stackrel{(1)}{=} y^T \Lambda y$$

where  $\stackrel{(1)}{=}$  is due to the change of variable  $y = U^T x$  (that implies  $y^T = x^T U$ ).

By the same argument used in the diagonal case,

$$\lambda_{\min} \cdot \|y\|^2 \leq y^T \Lambda y \leq \lambda_{\max} \cdot \|y\|^2.$$

Now the point is that if we can replace  $\|y\|^2$  with  $\|x\|^2$  we have proved the theorem. In fact this is true, due to the orthogonality of matrix  $U$ .

□

**Corollary 4.12.** Let  $A \in S(n, \mathbb{R})$  and let  $x \in \mathbb{R}^n$ . If  $x \neq 0$ ,  $\lambda_{\min} \leq \frac{x^T Ax}{\|x\|^2} \leq \lambda_{\max}$ , where  $\lambda_{\max}$  and  $\lambda_{\min}$  are respectively the eigenvalue of maximum value and the eigenvalue of minimum value.

**Definition 4.3** (Positive (semi)definite). Let  $A \in S(n, \mathbb{R})$ . We say that  $A$  is **positive semidefinite** if  $\forall x \in R^n x^T Ax \geq 0$ .

On the other hand,  $A$  is termed **positive definite** if  $x^T Ax > 0 \forall x \in R^n \setminus \{0\}$ .

**Fact 4.13.** Let  $A \in S(n, \mathbb{R})$ .  $\forall \lambda$  eigenvalue of  $A$   $\lambda \geq 0$  iff  $A$  is **positive semidefinite**.

On the other hand, all eigenvalues are **strictly positive** iff  $A$  is **positive definite**.

*Proof.*

$$(\Rightarrow) x^T Ax \geq \lambda_{\min}.$$

$(\Leftarrow)$  Let  $v_i$  be an eigenvector of  $A$ .

$$0 \leq v_i^T A v_i = v_i^T (\lambda_i v_i) = \lambda_i v_i^T v_i = \lambda_i \|v_i\|^2 \Rightarrow \lambda_i \geq 0.$$

□

**Fact 4.14.** Let  $B \in M(m, n, \mathbb{R})$ ,  $B^T B \in S(n, \mathbb{R})$  is positive semidefinite.

*Proof.* SYMMETRY:  $(B^T B)^T = B^T \cdot (B^T)^T = B^T B$ .

POSITIVE DEFINITE:  $x^T B^T B x = (Bx)^T (Bx) = \|Bx\|^2 \geq 0$

□

**Corollary 4.15.** The same holds for  $BB^T$ , since we can define  $C = B^T$ .

**Fact 4.16.** Let  $A \in S(n, \mathbb{R})$ .  $A \succeq 0$  and  $A$  invertible iff  $A$  is **strictly positive definite**.



Something on Matlab ...

In order to check if a matrix  $A$  is positive definite in Matlab we can look at its eigenvalues (cfr. `eig(A)`).

## 5 27th of September 2018 — A. Frangioni

**Definition 5.1** (Minimizing sequence). *Le us consider the **minimum problem** defined in Definition 3.1:*

$$(P) \quad f_* = \min\{f(x) : x \in X\}$$

*we term **minimizing sequence** a sequence that gets closer to the optimal value:  $\{x_i\}$  s.t.  $\{f(x_i)\} \rightarrow f_*$ .*

It goes without saying that we would like to avoid minimizing sequences that do not lead to an optimum.

As an example, consider

$\min\{x : x \in \mathbb{R} \wedge x > 0\} \{x_i = 1/i\}$ , where  $\{f(x_i)\} \rightarrow 0$ , but it is not an optimum or  
 $\min\{1/x : x \in \mathbb{R} \wedge x > 0\} \{x_i = i\}$ , where  $\{f(x_i)\} \rightarrow 0$ , but it is not an optimum.

**Definition 5.2** (Interior and border of a set). *Given  $S \subseteq \mathbb{R}^n$ , we say that  $x \in \text{int}(S)$  ( $x$  is an **interior point**) if  $\exists r > 0$  s.t.  $\mathcal{B}(x, r) \subseteq S$ .*

*On the other hand, we term **border points** those  $x \in \partial(S)$  such that  $\forall r > 0 \exists y, z \in \mathcal{B}(x, r)$ , where  $y \in S \wedge z \notin S$ . Intuitively, a point  $x$  lies on the border if, all the points in the ball centered in  $x$  lie for a half inside the set  $S$  and for the other half outside.*

Notice that a point on the boundary is not necessarily inside the ball, in that case we talk about **open set** (a set which is identical to its interior:  $S = \text{int}(S)$ ).

**Definition 5.3** (Closure of a set). *Let  $S \subset \mathbb{R}^n$  we term **closure** of  $S$  the set  $\text{cl}(S) = \text{int}(S) \cup \partial S$ .*

**Definition 5.4** (Closed set). *We say that a set  $S \in \mathbb{R}^n$  is **closed** if it coincides with his closure:  $S = \text{cl}(S)$ .*

*Equivalently, a set is termed **closed** if its complementary is open:  $\mathbb{R}^n \setminus S$ .*

It is interesting to notice that the cases of functions that lead to minimizing sequences that do not bring to an optimum are all defined in open sets.

Notice that there are sets that are both open and closed, for example  $\mathbb{R}^n$ .

**Definition 5.5** (Bounded set). *Let  $S \subseteq \mathbb{R}^n$ . We say that  $S$  is **bounded** if  $\exists r > 0$  such that  $S \subseteq \mathcal{B}(0, r)$ .*

*Intuitively, a bounded set does not go to  $\infty$ .*

**Definition 5.6** (Compact set). *Let  $S \subseteq \mathbb{R}^n$ . We term  $S$  **compact** if it is both closed and bounded.*

**Definition 5.7** (Accumulation point). *Let  $\{x_i\}$  be a sequence.  $x$  is an **accumulation point** if  $\exists \{x_{n_i}\}$  subsequence of  $\{x_i\}$  such that  $\{x_{n_i}\} \rightarrow x \equiv \liminf_{i \rightarrow \infty} d(x_i, x) = 0$ .*

**Fact 5.1.** Let  $S$  be a compact set and let  $\{x_i\} \subseteq S$  minimizing sequence. (Bolzano-Weierstrass)  $\{x_i\}$  has an **accumulation point**  $x \in S$ , thus the limit of the sequence has to be a feasible solution.

Why did we say *feasible* but not *optimal*? If the function is not continuous (cfr. Figure 5.4) it may happen that the sequence is minimizing, but the limit is not the optimum.




FIGURE 5.1: Case of non-continuity of the objective function in the border point  $(0, 0)$ .

**Definition 5.8** (Domain). Let  $f : D \rightarrow \mathbb{R}$ . We term  $D$  **domain**:  $D = \text{dom}(f)$ .

In this course we will not take into account the domain of functions, because we can make functions defined in the whole space as follows:

$$f : \mathbb{R}^n \rightarrow \bar{\mathbb{R}}, \text{ where } f(x) = \infty \text{ for } x \notin D.$$

**Definition 5.9** (Graph and epigraph). Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . We term **graph**  $gr(f) = \{(f(x), x) : x \in \text{dom}(f)\}$ .

On the other hand, we term **epigraph**  $epi(f) = \{(v, x) : x \in \text{dom}(f) \wedge v \geq f(x)\}$ .

**Definition 5.10** (Level and sublevel set). Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . We term **level set**  $L(f, v) = \{x \in \text{dom}(f) : f(x) = v\}$ .

On the other hand, we term **sublevel set**  $S(f, v) = \{x \in \text{dom}(f) : f(x) \leq v\}$ .

Intuitively, the level sets are projections of the points of the domain and they are used in order to “look” at the function even if we are working in  $\mathbb{R}^n$ .

A graphic example of level set is displayed in Figure 5.6.

## 5.1 Continuity

**Definition 5.11** (Continuity). Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . We say that  $f$  is **continuous** if  $\forall \varepsilon > 0 \exists \delta > 0$  such that  $|f(y) - f(x)| < \varepsilon \forall y \in \mathcal{B}(x, \delta)$ .




FIGURE 5.2: On the left-hand side the graph of the function  $f(x)$ , while on the right-handed side the epigraph of such function.




FIGURE 5.3: On the left-handed side of the figure, the level set of the function  $f$ , while on the right-hand side the sublevel set of such function.

Notice that if  $f, g$  continuous at  $x$  then:

1.  $f + g, f \cdot g$  continuous at  $x$
2.  $\max\{f, g\}$  and  $\min\{f, g\}$  continuous at  $x$
3.  $f \circ g \equiv f(g(\cdot))$  continuous at  $x$

**Theorem 5.2** (Intermediate value). *Let  $f : \mathbb{R} \rightarrow \mathbb{R}$ .  $f$  is continuous on  $[a, b]$  if  $\forall v \min\{f(a), f(b)\} \leq v \leq \max\{f(a), f(b)\} \exists c \in [a, b]$  such that  $f(c) = v$ .*

**Theorem 5.3** (Weierstrass extreme value theorem). *Let  $X \subseteq \mathbb{R}^n$  be a compact set and let  $f$  continuous on  $X$ . Then (P) has an optimal solution.*

*Equivalently, let  $X \subseteq \mathbb{R}^n$  compact and let  $f$  continuous on  $X$ . Then all accumulation points of any minimizing sequence are optima and there is at least one.*

**Definition 5.12** (Lipshitz continuity). Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . We term  $f$  **Lipschitz continuous** (L.c.) on  $S \in \mathbb{R}^n$  if  $\exists L > 0$  such that

$$|f(x) - f(y)| \leq L \|x - y\| \quad \forall x, y \in S$$

More generally,  $f$  is **globally Lipschitz continuous** if  $S = \mathbb{R}^n$  and it is **locally Lipschitz continuous** if at  $x \exists \varepsilon > 0$   $S = \mathcal{B}(x, \varepsilon)$ .

Notice that the  $L$  constant value depends on  $S$ . The wider the set the smaller  $L$ .

**Fact 5.4.** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . On a compact set  $S \in \mathbb{R}^n$  if  $f$  is continuous then it is Lipschitz continuous.

**Definition 5.13** (Lower-upper semi-continuous). Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$   $f$  is **lower[upper] semi-continuous** (l./u.s.c.) at  $x$  if

$$\{x_i\} \rightarrow x \implies f(x) \leq \liminf_{i \rightarrow \infty} f(x_i) \quad [f(x) \geq \limsup_{i \rightarrow \infty} f(x_i)]$$

Equivalently,  $\liminf_{y \rightarrow x} f(y) \geq f(x)$ ,  $\limsup_{y \rightarrow x} f(y) \leq f(x)$ .

## 5.2 Derivatives

In this section we address the problem of inferring information on a complicated function around a certain value  $\bar{x}$  using very simple functions, that provide information which is reliable close to the point we are studying. Those functions are called “derivatives”.

It goes without saying that it is not possible to compute such simple representation in any kind of point.

**Definition 5.14** (Left and right derivative). Let  $f : \mathbb{R} \rightarrow \mathbb{R}$ . We term **left derivative**  $f'_-(x) = \lim_{t \rightarrow 0_-} [f(x+t) - f(x)]/t$ .

On the other hand, **right derivative**  $f'_+(x) = \lim_{t \rightarrow 0_+} [f(x+t) - f(x)]/t$ .

**Definition 5.15** (Differentiable). Let  $f : \mathbb{R} \rightarrow \mathbb{R}$ . We say that  $f$  is **differentiable** at  $x \in \text{dom}(f)$  if  $f'_-(x) = f'_+(x) \Leftarrow$  they  $\exists$ .

**Fact 5.5.** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  and differentiable in  $x \in \text{dom}(f)$  then  $f$  is continuous in  $x$ .

**Theorem 5.6** (Mean value theorem).  $f : \mathbb{R} \rightarrow \mathbb{R}$  continuous on  $[a, b]$  and differentiable on  $(a, b)$  then  $\exists c \in (a, b)$  s.t.  $f'(c) = (f(b) - f(a))/(b - a)$ .

**Theorem 5.7** (Rolle's theorem). Let  $f : \mathbb{R} \rightarrow \mathbb{R}$ .  $f(a) = f(b)$  then  $\exists c \in (a, b)$  s.t.  $f'(c) = 0$

**Corollary 5.8.** In the same hypothesis of Rolle's theorem, let  $a$  and  $b$  consecutive roots of  $f$ . Then  $f'(a)$  and  $f'(b)$  have opposite sign.

### 5.2.1 Multivariate differentiability

**Definition 5.16** (Partial derivative). Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . We term **partial derivative** of  $f$  w.r.t.  $x_i$  at  $x \in \mathbb{R}^n$  as

$$\frac{\partial f}{\partial x_i}(x) = \lim_{t \rightarrow 0} \frac{f(x_1, \dots, x_{i-1}, x_i + t, x_{i+1}, \dots, x_n) - f(x)}{t}$$

In other words, it is just  $f'(x_1, \dots, x_{i-1}, x, x_{i+1}, \dots, x_n)$ , treating  $x_j$  for  $j \neq i$  as constants.

**Definition 5.17** (Gradient). Given  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . We term **gradient** of  $f$  the vector of all the partial derivatives.

**Definition 5.18** (Directional derivative). Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . We term **directional derivative** of  $f$  in point  $x \in \text{dom}(f)$  along direction  $d \in \mathbb{R}^n$

$$\frac{\partial f}{\partial d}(x) := \lim_{t \rightarrow 0} \frac{f(x + td) - f(x)}{t}$$

In a multivariate space it is not possible to assure differentiability checking if the directional derivatives are equal, because there is an infinite number of different directions.

We are now ready to introduce the notion of multivariate differentiability.

**Definition 5.19** (Differentiable). Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . We say that  $f$  is **differentiable** at  $x$  if  $\exists$  a linear function  $\phi(h) = \langle c, h \rangle + f(x)$  s.t.

$$\lim_{\|h\| \rightarrow 0} \frac{\|f(x + h) - \phi(h)\|}{\|h\|} = 0$$

The intuition here is that the linear function should approximate  $f$  pretty well around  $x$ .

The gradient is the direction on which the function increases more rapidly, while the opposite of the gradient suggests the direction on which the function decreases most.

**Fact 5.9.** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  differentiable at  $x$ . Then  $f$  is locally Lipschitz continuous at  $x$ .

**Corollary 5.10.** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  differentiable at  $x$ . Then  $f$  is continuous at  $x$ .

Notice that the converse does not hold.

**Fact 5.11.** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and let us assume  $\exists \delta > 0$  s.t.  $\forall i \frac{\partial f}{\partial x_i}(y)$  is continuous  $\forall y \in \mathcal{B}(x, \delta)$ . Then  $f$  differentiable at point  $x$ .

Notice that the converse of Proposition 5.11 does not hold either.

We are now ready to introduce a class of functions that allows good results for optimization:

$\mathcal{C}^1 := \nabla f(x)$  continuous.

Let  $f \in \mathcal{C}^1$ , then  $f$  is differentiable everywhere and also continuous everywhere.

Figure 5.4, Figure 5.5 and Figure 5.6 picture some functions which are not differentiable in the minimum.




FIGURE 5.4: The function  $f(x_1, x_2) = \|[x_1, x_2]\| = |x_1|+|x_2|$  has some kinks.




FIGURE 5.5: The function  $f(x_1, x_2) = \frac{x_1^2 x_2}{x_1^2+x_2^2}$  may be put to 0 in  $(0,0)$  for continuity, but it is still not differentiable in  $(0,0)$  although the function admits directional derivatives in  $(0,0)$ .




FIGURE 5.6: The function  $f(x_1, x_2) = \left(\frac{x_1^2 x_2}{x_1^4+x_2^2}\right)^2$  is not continuous in 0. There are some directions that lead to the limit 0, while there are some other directions (the parabolas above) that do not lead to value 0.

**Definition 5.20** (First order model). Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and let  $x \in \text{dom}(f)$ . We term **first**

*order model of  $f$  at point  $x$*

$$L_x(y) = \nabla f(x)(y - x) + f(x)$$

**Fact 5.12.** *Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and let  $x \in \text{dom}(f)$  such that  $f$  is differentiable at  $x$ . Then  $(L_x, f(x)) \perp S(f, f(x)) \perp \nabla f(x)$ .*

Geometrically speaking, we can observe that a function is smooth whenever its levelsets are smooth.

# 6 18th of October 2018 — F. Poloni

## 6.1 Least squares problem

**Fact 6.1.** Given  $A \in \mathcal{M}(m, n, \mathbb{R})$  the following conditions are equivalent:

- $A^T A$  is positive definite;
- $A$  has full column rank;
- the columns of  $A$  are linear independent;
- $\text{Ker}(A) = \{0\}$ .

**Fact 6.2.** Given  $A \in \mathcal{M}(m, n, \mathbb{R})$ , if  $A^T A$  is **positive semidefinite**  $f(x) = x^T Qx - q^T x + b^T b$  is **strongly** (or **strictly**) **convex**. In other words,  $f(x)$  has a **unique** minimum.

We find the minimum by solving  $\nabla f(x) = 0$ ,

where  $f(x) = x^T A^T A x - 2b^T A x + b^T b$

$\nabla f(x) = 2A^T A x - 2A^T b$  How should we compute this gradient?

We know that  $f(x + h) = f(x) + (\nabla f(x))^T h + o(\|h\|)$

$$\begin{aligned}
 f(x + h) &= (x + h)^T A^T A (x + h) - 2b^T A (x + h) + b^T b \\
 &= \mathbf{x}^T \mathbf{A}^T \mathbf{A} \mathbf{x} + \mathbf{x}^T \mathbf{A}^T \mathbf{A} \mathbf{h} + \mathbf{h}^T \mathbf{A}^T \mathbf{A} \mathbf{x} + \mathbf{h}^T \mathbf{A}^T \mathbf{A} \mathbf{h} - \mathbf{2b}^T \mathbf{A} \mathbf{x} - 2b^T A h + \mathbf{b}^T \mathbf{b} \\
 &= \mathbf{f}(\mathbf{x}) + (\mathbf{2x}^T \mathbf{A}^T \mathbf{A} \mathbf{h} - 2b^T A h) + \mathbf{o}(\|\mathbf{h}\|) \\
 &= f(x) + (\mathbf{A}^T \mathbf{A} \mathbf{x} - \mathbf{2A}^T \mathbf{b})^T \mathbf{h} + o(\|h\|)
 \end{aligned} \tag{6.1}$$

So,  $\nabla \mathbf{f}(\mathbf{x}) = \mathbf{A}^T \mathbf{A} \mathbf{x} - \mathbf{2A}^T \mathbf{b}$

We would like to know when the gradient is 0.

$$\nabla f(x) \stackrel{?}{=} 0 \Leftrightarrow A^T A x = A^T b$$

Since  $A^T A$  is a **square** matrix and also **non singular** (which means invertible) we may find  $x$  by solving a linear system  $x = (A^T A)^{-1} (A^T b)$  via:

- Gaussian elimination;
- LU factorization;
- QR factorization;
- Cholesky factorization (specialized method for positive definite matrices) Idea:  $A^T A$  can be written as  $A^T A = R^T R$ , where  $R$  is a square, upper triangular matrix.

Why do we need factorization method? Let's compute complexity:

- $A^T A \rightarrow 2mn^2$ , where  $m > n$ ;
- $A^T b \rightarrow 2mn$ ;
- Solving  $A^T A x = A^T b$  with gaussian elimination has a computational complexity of  $\frac{2}{3}n^3$ ;
- Cholesky factorization  $A^T A = R^T R$  which has a cost of  $\frac{1}{3}n^3$

### 6.1.1 Method of normal equations

This method solves least squares problem and takes his name from the fact that “normal” means orthogonal.

The key idea is using symmetry to skip half of the entries of  $A^T A$

If  $Ax = b$  can't be solved, since  $A$  is tall and thin, we can multiply on both sides by  $A^T$  and try again, since the matrix is square now.

The residual  $Ax - b$  is orthogonal to any vector  $v \in \text{span}(A) : (Av)^T(Ax - b) = 0$

Why?

$v^T(A^T A x - A^T b) = 0$ , see Figure 6.1.




FIGURE 6.1: Geometric idea.

It's possible to find a close formula for solving this problem:  $\min \|Ax - b\|$  is given by  $x = (A^T A)^{-1} A^T b$ .

**Definition 6.1** (Moore-Penrose pseudoinverse). *Let  $A$  be a matrix in  $\mathcal{M}(n, m, \mathbb{R})$ , the Moore-Penrose pseudoinverse of  $A$  with full column rank is  $\mathbf{A}^\dagger := (A^T A)^{-1} A^T$*

So we can write  $x = A^\dagger b$  for the solution of a LS problem.

In particular, the solution of  $\|\min Ax - (b_1 + b_2)\|$  is the sum of two solutions of  $\min \|Ax_1 - b_1\|$  and  $\min \|Ax_2 - b_2\|$ .

**Obs:**  $AA^\dagger = A(A^T A)^{-1} A^T = AA^{-1} A^{T-1} A^T = I_{n \times m}$ , but this doesn't hold for  $A^\dagger A = (A^T A)^{-1}(A^T A)$ , which is not the identity matrix.

## 6.2 QR factorization

**Theorem 6.3.**  $\forall A \in \mathcal{M}(m, m, \mathbb{R})$ ,  $\exists Q \in \mathcal{O}(m, m, \mathbb{R})$  (space of orthogonal matrices of size  $m \times m$ ),  $\exists R \in \mathcal{M}(m, m, \mathbb{R})$  upper triangular such that  $A = QR$

QR factorization isn't as powerful as SVD factorization.

Why are we interested in studying factorizations?

- They reveal properties: singularity, rank, ...;
- They may be an intermediate step in algorithms.

**Example 6.1.** We would like to solve  $Ax = b$ , with  $A \in \mathcal{M}(m, m, \mathbb{R})$  we may:

1. first compute the QR factorization ( $A = QR$ ) and then obtain  $x = A^{-1}b = R^{-1}Q^{-1}b$
2. compute then  $c = Q^Tb$
3. and then  $x = R^{-1}c$

What's the computational cost?

1.  $QR \rightarrow O(m^3)$
2. compute  $c \rightarrow O(m^2)$
3. compute  $x \rightarrow O(m^2)$

Let's analyze the case in which  $A$  is a vector. Given  $x \in \mathbb{R}^m$ , we want to find an orthogonal

matrix  $Q$  such that  $Qx$  has the form  $\begin{pmatrix} s \\ 0 \\ \vdots \\ 0 \end{pmatrix} = se_1$ , where  $s = \pm\|x\|$ .

We denote  $e_i$  the i-th column of the identity matrix.

**Definition 6.2** (Householder reflector). Let  $U$  be a vector in  $\mathbb{R}^m$ . An **Householder reflector** is a matrix  $H$  such that  $H = I - \frac{2}{U^T U} \cdot UU^T$ .

We can observe that  $U^T U$  is a scalar.

Since  $U^T U = \|U\|^2$ , another way of seeing  $H$  may be  $H = I - \frac{2}{\|U\|^2} UU^T = I - 2vv^T$ , where  $v = \frac{1}{\|U\|}U$

**Lemma 6.4.** Matrices of this form are orthogonal.

*Proof.*

$$\begin{aligned}
HH^T &= \left(I - \frac{2}{\|U\|^2} \cdot UU^T\right) \cdot \left(I - \frac{2}{\|U\|^2} \cdot UU^T\right) \\
&= I \cdot I - \frac{2}{\|U\|^2} \cdot UU^T I - I \cdot \frac{2}{\|U\|^2} \cdot UU^T + \frac{2}{\|U\|^2} \cdot UU^T \cdot \frac{2}{\|U\|^2} \cdot UU^T \\
&= I - \frac{2}{\|U\|^2} \cdot UU^T - \frac{2}{\|U\|^2} \cdot UU^T + \frac{4}{\|U\|^4} \cdot UU^T UU^T \\
&= I - \frac{4}{\|U\|^2} UU^T + \frac{4}{\|U\|^4} U \|U\|^2 U^T \\
&= I
\end{aligned} \tag{6.2}$$

□

**Example 6.2.**  $Hx = \left(I - \frac{2}{\|U\|^2} UU^T\right)x = x - \frac{2}{\|U\|^2} U(U^T U)$

$y = \text{compute\_product}(U, x)$

$a = U' * x$

$b = U' * U$

$y = x \frac{2*a}{b} \cdot U$

All these operations are linear operations, so the complexity is  $O(m)$ , cheaper than generic matrix-vector product ( $O(m^2)$ ).

Can we find a matrix  $H$  in this family that maps  $x$  to  $y$  (equivalently  $Hx = y$ )? The answer is given by the following lemma

**Lemma 6.5.**  $\forall x, y \ s.t \ \|x\| = \|y\| \exists H \ s.t. \ Hx = y$ , choosing  $u = x - y$ .

A geometric idea is given by the following




FIGURE 6.2: On the left in  $\mathbb{R}^2$ , on the right in  $\mathbb{R}^3$ .

What happens if  $y = \begin{pmatrix} \|x\| \\ 0 \\ \cdot \\ \cdot \\ \cdot \\ 0 \end{pmatrix}$ ?

$y = H^T x$ , actually  $H^T = H^{-1} = H$  Let's map  $x$  to  $y$ :

$$U = x - y = \begin{pmatrix} x_1 \\ x_2 \\ \cdot \\ \cdot \\ \cdot \\ x_n \end{pmatrix} - \begin{pmatrix} s \\ 0 \\ \cdot \\ \cdot \\ \cdot \\ 0 \end{pmatrix} = \begin{pmatrix} x_1 - s \\ x_2 \\ \cdot \\ \cdot \\ \cdot \\ x_n \end{pmatrix}$$

where  $s = \|x\|$

### Matlab syntax for functions

`function[v,s] = householder_vector(x)`, where v and s are the returned values and x is the argument.

# 7 19th of October 2018 — A. Frangioni

## 7.1 Gradient method for quadratic functions

This is the simplest possible family of functions where a minimum exists.



Do you recall?

A quadratic function is defined as:  $f(x) = \frac{1}{2}x^T Qx + qx$ , so its gradient is the following  
 $\nabla f(x) = Qx + q$

We are interested in finding local minima of the function  $f$ .

**Fact 7.1.** A quadratic function  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ , s.t.  $f(x) = \frac{1}{2}x^T Qx + qx$  admits a minimum iff  $Q \succeq 0$  ( $Q$  is positive semidefinite).

The gradient method generates points that move along orthogonal directions. More formally,  $x^{i+1} = x^i + \alpha^i d^i$ , where  $d^i = -\nabla f(x^i) = -Qx^i - q$




FIGURE 7.1: Some iterations of the gradient method

**Fact 7.2.**  $\forall i < d^i, d^{i+1} \rangle = 0$ .

*Proof.* TODO □

$$i\alpha^i = \frac{\|d^i\|^2}{d^i Q d^i}, x^i \rightarrow \bar{x}$$

Our aim is to estimate how fast this converges.

$f(x^i) - f_* = -\frac{1}{2}q^T Q^{-1}q$  we know everything for this function, from last lecture ( $x^* = -Q^{-1}q$ ).

What can we say about  $f(x^{i+1}) - f_* = \frac{1}{2}(x^{i+1} - x_*)^T A(x^{i+1} - x_*)$ ?

I want it in terms of  $x^{i+1} = x^i - \frac{\|Qx^i - q\|^2}{(Qx^i - q)^T Q(Qx^i - q)}$

From this formula we can say that the error at the current step ( $i + 1$ ) is equal to the error of the step before ( $i$ ) divided by something. More precisely,  $f_*(x) = \frac{1}{2}(x - x_*)^T Q(x - x_*) = f(x) + \frac{1}{2}x_*^T Qx_* = f(x) - f_*$

If the quantity after the minus is positive but less than 1 the error at the next step will be less than the error at the previous step. This means not only **linear convergence**, but a bit more, because linear convergence only takes into consideration steps in proximity to the limit, while this formula holds also at the beginning.

**Fact 7.3.** If  $Q$  is positive semidefinite  $d^T Q d = \|d\|_Q^2$

**Fact 7.4.** If  $Q$  is positive definite we can say that the error goes like  $1 - \left( \frac{\|d_i\|^2}{(d^i)^T Q d^i) (d^i)^T Q^{-1} d^i} \right)$  or, equivalently,  $1 - \frac{\|d_i\|_I^2}{\|d_i\|_Q^2} \cdot \frac{\|d_i\|_I^2}{\|d_i\|_{Q^{-1}}^2}$ .

We are measuring a vector with three different norms.

We would like to estimate  $\frac{\langle d_i, d_i \rangle}{d_i^T Q d_i}$ .

Given  $\lambda_i$  the eigenvalues of  $Q$ ,  $\frac{1}{\lambda_i}$  are the eigenvalues of the matrix  $Q^{-1}$ .

We can say that  $\lambda^n \|x\|^2 \leq x^T Q x \leq \lambda^1 \|x\|^2$ , where  $\lambda^n$  is the smallest eigenvalue, while  $\lambda^1$  is the biggest.

We are looking for a close formula for calculating the convergence rate, since it depends recursively by the steps done. So we want to do a worst case analysis in order to find a faster way to calculate convergence rate.

We want to prove that  $R$  is smaller than 1 so we are looking for an upperbound. A coarse upperbound is  $(1 - \frac{\lambda^n}{\lambda^1})$ , but we can prove more:

$$\forall x \in \mathbb{R}^n \quad \frac{\|x\|^4}{(x^T Q x)(x^T Q^{-1} x)} \geq \frac{4\lambda^1 \lambda^n}{(\lambda^1 + \lambda^n)^2}$$

We won't see the proof of this fact.

$R$  close to 0 means that the algorithm is converging fast, so when the larger eigenvalue ( $\lambda^1$ ) and the smallest eigenvalue ( $\lambda^n$ ) are very close to each other the algorithm is converging fast. We can say that, since the eigenvalues are the axis of the ellipsoid, the algorithm is converging fast when the ellipsoid has a round shape.

We can provide an even better estimate, which is  $(\frac{\lambda^1 - \lambda^n}{\lambda^1 + \lambda^n})^2$

How can we understand if the number of iterations needed to converge is a good result? Well, it depends on how that number is obtained. If it's dimensional independent it's very good, because it scales well (when the size of the space — number of variables — increases). It only depends on the conditioning of the matrix  $Q$ .

Of course as  $n$  grows  $Q$  changes, so in practice it may happen that the conditioning of the problem is worsening as  $n$  grows.

If the balls are very rounded the zig zags needed to start converging are very few.

**Obs:** We found a bound for the convergence speed of the algorithm when  $Q$  is positive definite. What can we say when  $Q$  is positive semidefinite? The algorithm works, but we can't provide an upperbound for the convergence rate. We are even more restrictive when dealing with machine precision, since if there is an eigenvalue which is bigger than zero, but very close to zero, becomes 0 on the machine, so we can't give an upperbound.

We will see how to deal with this case.

## 7.2 MatLab implementation

Let's talk about the implementation. First of all we need to set a proper value for  $\epsilon$ . A good idea would be the norm of the gradient. We need to give also some performance bound, like maximum number of iterations or the maximum amount of time.

**MatLab call:**

```
1 function [x, status] = SDQ(Q, q, x, fStar, eps, MaxIter)
```

where  $f_{\text{Star}}$  is the optimal value, which should give us an idea of the convergence.

### Note

- Always check the coherence of input values (check if the user passed allowed parameters);
- Always give all possible information about your result (for example if the algorithm stopped because the maximum number of iterations was reached or because the epsilon value was reached);
- Always check in your code the accepted values of variables during calculations: for example if you need to divide by a quantity that may be smaller than the precision check it before computing the ratio;
- Design a good log, in order to understand what's happening at each step.

In the code of that function we also kept track of the actual ratio between the error at one step and the error at the next one. This information tells us how many orders of magnitude the error decrease. We can find starting points where the ratio is exactly  $R$ . We can observe that when the conditioning is quite good the error decrease faster than the  $R$  limitation, but as soon as we change the values for  $Q$  and  $q$  things may be worse.

We saw from some examples that if the conditioning grows the number of iterations needed to find the minimum increases as well.

Trying the algorithm on some examples shows that the theoretic results are reflected well in the practical case.

### 7.2.1 Error

When we are running an algorithm, starting from a point that will not lead to the optimum we would like to stop anyway, at a certain point, when we are close enough to the solution.

$$\varepsilon_A = f(x^i) - f_* \leq \varepsilon \quad (\text{absolute error})$$

In this context, we introduce the concept of **relative error**, since the error may be big or small if compared with the value of the function.

This relative error is invariant for scaling transformations. Notice that if  $f^*$  might be zero the formula should be changed.

$$\varepsilon_R = (f(x^i) - f_*) / |f_*| = \varepsilon_A / |f_*| \leq \varepsilon \quad (\text{relative error})$$

The problem is that often we don't know  $f^*$ , so it's impossible to compute this kind of error.

In this very common case a good lower bound  $\underline{f} \leq f^*$  for  $f^*$ . In this course we won't focus on finding  $\underline{f}$ .

Another stopping conditions may be the following:

- $\|\nabla f(x^i)\| \leq \varepsilon$  (“absolute version”)
- $\|\nabla f(x^i)\| / \|\nabla f(x^0)\| \leq \varepsilon$  (“relative version”)

The second stopping condition is expressed in relation to the first value for the norm of the gradient.

We usually chose the norm of the gradient as a threshold for precision, but we don't know how this quantity relates to  $\varepsilon_A$  or  $\varepsilon_B$ .

**Example 7.1.** for  $X = \mathcal{B}(0, r)$  and  $f$  convex, estimate  $\varepsilon_A$  when  $\|\nabla f(x^i)\| \leq \varepsilon$

I've got a convex function and I know the minimum is in a ball. We can minimize the linear function in the range of the ball and that minimum is surely a lower bound.

# 8 24th of October 2018 — A. Frangioni

## 8.1 Gradient method for non quadratic functions

We want to move from quadratic functions to wider families of functions.



Do you recall?

The step size  $\alpha$  in the quadratic case is defined as follows:  $\alpha = \frac{\|d\|^2}{d^T Q d}$

We would like to find a more general form for the step size, which doesn't depend on the fact that the function is quadratic.

The algorithm for finding local minima of non quadratic functions has the same structure of the one used for quadratic ones, i.e. first compute the direction of the step and then compute its size.

We will see that, differently from the quadratic case (where the gradient was  $\nabla f(x) = Qx + q$ ) computing the gradient in this more general case isn't easy at all.

We may recall from last lecture that the proof of the orthogonality of the gradient doesn't depend on the quadratic nature of our functions, so it works in this case too.

### 8.1.1 How fast does it converge?

Given Algorithm 8.1 for finding minima of quadratic functions (that differs from the one provided for quadratic ones for the step size) we would like to understand how fast the convergence is.

---

ALGORITHM 8.1 Pseudocode for non quadratic functions local minimum detection.

---

```

1: procedure SDQ( $f, x, \varepsilon$ )
2:   while ( $\|\nabla f(x)\| > \varepsilon$ ) do
3:      $d \leftarrow -\nabla f(x);$ 
4:      $\alpha \leftarrow \frac{\|d\|^2}{(d^T Q d)};$ 
5:      $x \leftarrow x + \alpha d;$ 
6:   end while
7: end procedure

```

---

**Theorem 8.1.** Let  $f$  be a function in  $C^2$  and let  $x_*$  be a local minimum for  $f$  s.t.  $\nabla^2 f(x_*) \succ 0$  (which means that the Hessian matrix is strictly positive definite):

$$\{x^i\} \rightarrow x_* \implies \{f(x^i)\} \rightarrow f(x_*)$$

linearly, with same  $R$  as the quadratic case, depending on  $\lambda_1$  and  $\lambda_n$  of  $\nabla^2 f(x_*)$

This theorem means that if the function is differentiable and the Hessian is strictly positive definite then when getting closer and closer to the minimum, the function is more and more similar to a quadratic function.

This similarity is a good news, since we can use the same methods of the quadratic case, but, as we recall from the previous lecture, we must pay attention to **conditioning**.

At this point we need to work on finding the local minimum of the one dimensional function  $\varphi^i$ , s.t.

$$\varphi^i(\alpha) = f(x^i + \alpha d^i),$$

where  $d^i = -\nabla f(x^i)$ .

Let's omit the  $i$ , since we are concentrating on a single iteration.

We are interested in finding a  $\alpha^*$  such that  $\varphi'(\alpha^*) = 0$

**Example 8.1.** Let's suppose we are in  $\mathbb{R}^2$  and  $f(x, y) = x^2 e^y$ . We can differentiate  $F$  and  $\nabla f(x, y) = (2xe^y, x^2 e^y)$ .

At the  $i$ -th iteration  $(x, y) = (1, 0)$ , so  $\nabla f(1, 0) = (2, 1)$ . Now  $x(\alpha) = (1, 0) - \alpha(2, 1) = (1 - 2\alpha, 0 - \alpha)$ .

At this point we obtain  $\varphi(\alpha) = f(x(\alpha)) = (1 - 2\alpha)^2 e^{-\alpha}$ .

It's hard to find the roots of this function. The points we can find are not  $\varphi'(\alpha) = 0$ , but instead  $|\varphi'(\alpha)| \leq \varepsilon'$ , where the meaning of  $\varepsilon$  is bounding the directional derivative to be small.

Chi è  
 $x(\alpha)$ ?

**Fact 8.2.** Let  $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ , such that  $\varphi(\alpha) = f(x^i + \alpha d^i)$ ,  $\varphi'(\alpha) = \langle \nabla f(x^i + \alpha d^i), d \rangle$ .

*Proof.* TODO: using the **chain rule**:  $f : \mathbb{R}^m \rightarrow \mathbb{R}^k$ ,  $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$ ,  $h : \mathbb{R}^n \rightarrow \mathbb{R}^k$  such that  $h(x) = f(g(x)) \Rightarrow \mathbf{J}h(\mathbf{x}) = \mathbf{J}f(\mathbf{g}(\mathbf{x})) \cdot \mathbf{J}g(\mathbf{x})$

Obs:  $Jf \in \mathbb{R}^{k \times m}$ ,  $Jg \in \mathbb{R}^{m \times n}$  and  $Jh \in \mathbb{R}^{k \times m} \cdot \mathbb{R}^{m \times n} = \mathbb{R}^{k \times n}$ . □

**Fact 8.3.** We claim that  $\varepsilon' = \varepsilon$ .

*Proof.* We want to find the relationship between the two parameters  $\varepsilon$  and  $\varepsilon'$ . Assuming that we've got a black box that finds  $\alpha$ , given  $\varepsilon'$ , we are interested in computing  $\varepsilon'$ .

**Key idea:** Normalization of the direction.

We may normalize the direction of movement  $d^i$  without perturbing the behaviour of the algorithm:  $d^i = -\frac{\nabla f(x^i)}{\|\nabla f(x^i)\|}$ . Note that we're not worried of dividing by the norm of the gradient, since if it gets 0 we have already stopped the procedure.

In this new context  $\|d^i\| = 1$  and

$$\begin{aligned}
 \varphi'(0) &= \frac{\partial f}{\partial d}(x) \\
 (\text{From Proposition 8.2}) &= \langle \nabla f(x), d \rangle \\
 &= \langle \nabla f(x), \frac{-\nabla f(x)}{\|\nabla f(x)\|} \rangle \\
 &= -\frac{\langle \nabla f(x), \nabla f(x) \rangle}{\|\nabla f(x)\|} \\
 &= -\frac{\|\nabla f(x)\|^2}{\|\nabla f(x)\|} \\
 &= -\|\nabla f(x^i)\|
 \end{aligned} \tag{8.1}$$

$$|\varphi'(\alpha^i)| = |\langle \nabla f(x^{i+1}), d^i \rangle| = \left| \langle \nabla f(x^{i+1}), -\frac{\nabla f(x^i)}{\|\nabla f(x^i)\|} \rangle \right|$$

Hence, if  $\{x^i\} \rightarrow x$

$$\lim_{i \rightarrow \infty} \left| \langle \frac{\nabla f(x^i)}{\|\nabla f(x^i)\|}, \nabla f(x^{i+1}) \rangle \right| = \langle \frac{\nabla f(x)}{\|\nabla f(x)\|}, \nabla f(x) \rangle = \|\nabla f(x)\| \leq \varepsilon \tag{8.2}$$

Since  $\|\nabla f(x^i)\| > \varepsilon$  we have the thesis.  $\square$

Per each phase the new epsilon is obtained  $\varepsilon \leftarrow \varepsilon \|\nabla f(x^i)\|$ .

If we can prove that the algorithm is converging we know when to stop.

This convergence isn't the perfect mathematical convergence, since  $\varepsilon \neq 0$ , because the line search will never terminate.

### 8.1.2 Exact line search, first orderd approach

We want to find the minimum points of  $\varphi$ , which corresponds to points where the first order derivative is zero and it goes from negative to positive. Since we are talking about numerical algorithms we are going to stop a little before the minimum is reached.

**Key idea:** We would like to reduce the range in which performing the search, at each step.

How can we be sure that in a given range there is a point where the derivative is 0? Rolle's theorem, as shown in Figure 8.1.

Since the gradient is continuous the directional derivative is continue, so  $\varphi$  is continuous (the scalar product is continuous).

Actually, we only need to find where the derivative is positive, because the 0 of the derivative is between the previous value and this point.

The algorithm is the following:




FIGURE 8.1: First, we restrict from  $\mathbb{R}$  to  $[x_1, x_2]$ , then double  $\alpha$  until the derivative is greater than 0

---

**ALGORITHM 8.2** First algorithm for exact line search

---

```

1:  $\bar{\alpha} \leftarrow x_1$ ; #or whatever value  $> 0$ 
2: while ( $\varphi'(\bar{\alpha}) < 0$ ) do
3:    $\bar{\alpha} \leftarrow 2\bar{\alpha}$ ; #or whatever factor  $> 1$ 
4: end while

```

---

We'll stop when  $\alpha < -10^{308}$ , which is the smallest value for a double, since we will get a numerical error and stop.

Works if  $\varphi$  **coercive**:  $\lim_{\alpha \rightarrow \infty} \varphi(\alpha) = \infty$  (ex.  $f$  strongly convex)

**Exercise 8.1.** Build an example where  $\bar{\alpha}$  exists but it is not found by this algorithm.

**Solution:** The function changes its derivative in a range between  $\alpha$  and  $2\alpha$ .

An alternative to the algorithm presented above may be the bisection algorithm, which follows.

---

**ALGORITHM 8.3** Bisection algorithm

---

```

1: procedure LSBM( $(\varphi' \bar{\alpha} \varepsilon)$ )
2:    $\alpha_- \leftarrow 0$ ;
3:    $\alpha \leftarrow \alpha_+$ ;
4:    $\alpha_+ \leftarrow \bar{\alpha}$ ;
5:   while ( $|\varphi'(\alpha)| > \varepsilon$ ) do
6:      $\alpha \leftarrow (\alpha_+ + \alpha_-)/2$ ;
7:     if ( $\varphi'(\alpha) < 0$ ) then
8:        $\alpha_- \leftarrow \alpha$ ;
9:     else
10:       $\alpha_+ \leftarrow \alpha$ ;
11:    end if
12:   end while
13: end procedure

```

---




FIGURE 8.2: The information we have about function  $\varphi$

We would like to improve this algorithm too, finding a better point in the middle than the middle point.

We may use the information we have about the function, since we know  $\varphi(\alpha^-)$ ,  $\varphi'(\alpha^-)$ ,  $\varphi(\alpha^+)$  and  $\varphi'(\alpha^+)$ .

At this point we can write a model ( $m(\alpha) = aa^2 + ba + c$ ) and specialize it with the information we have, via computing a linear system:

$$\begin{cases} a(\alpha^-)^2 + b(\alpha^-) + c = \varphi(\alpha^-) \\ a(\alpha^+)^2 + b(\alpha^+) + c = \varphi(\alpha^+) \\ 2a\alpha^- + b = \varphi'(\alpha^-) \\ 2a\alpha^+ + b = \varphi'(\alpha^+) \end{cases}$$

Then we look for the stationary point of this function and that's the ball where I'm likely to find the root of the derivative.

We can say something more, since the following fact holds:

**Fact 8.4.** Let  $\varphi \in C^3$ , then quadratic interpolation has convergence of order  $1 < p < 2$  (superlinear)

In Figure 8.3 we can observe a situation in which the hypothesis of Proposition 8.4 aren't satisfied.




FIGURE 8.3: If the function isn't  $C^3$  and one derivative is very big, then the range doesn't shrink much.

We would like to modify the formula to have at least linear convergence.

We can ensure to move not to close to one of the extremes, for example more than 10%.

Another idea is, since we have four equations, to build a cubic function, although the operation is very long. In this case the convergence get quadratic, which is better than linear.

# 9 25th of October 2018 — A. Frangioni

We need to choose a descending direction, we have more choices than the opposite of the gradient.

We want the derivative to be almost zero, given a tolerance value.

Let's see another variant of line search.

## 9.0.1 Line search: second order approaches

**Theorem 9.1.** Given  $f \in C^2$ ,  $\exists \varphi''(\alpha) = d^T \nabla^2 f(x + \alpha d)d$  and it's continuous.

*Proof.* Via chain rule.  $\square$

Since we are looking for a point where the derivative  $\varphi'(\alpha) = 0$ , we may use the second derivative to write a model and, assuming to trust the model, it can be studied.

**Definition 9.1** (Model–Newton tangent method). Our model, in this case is

$$\varphi'(\alpha) \approx \varphi'(\alpha^k) + \varphi''(\alpha^k)(\alpha - \alpha^k)$$

In this context, solving  $\varphi'(\alpha) = 0$  implies finding those  $\alpha$  such that  $\alpha = \alpha^k - \varphi'(\alpha^k)\varphi''(\alpha^k)$

---

ALGORITHM 9.1 Pseudocode for Newton method.

---

```

1: procedure LSNM( $\varphi'$ ,  $\varphi''$ ,  $\alpha$ ,  $\varepsilon$ )
2:   while ( $|\varphi'(\alpha)| > \varepsilon$ ) do
3:      $\alpha \leftarrow \alpha - \frac{\varphi'(\alpha)}{\varphi''(\alpha)}$ ;
4:   end while
5: end procedure

```

---

We need to understand when and why  $\varphi''(\alpha) \neq 0$  and when and why this method converges.

**Theorem 9.2.** Let  $\varphi \in C^3$  such that  $\varphi'(\alpha_*) = 0$  and  $\varphi''(\alpha_*) \neq 0$ .  $\exists \delta > 0$  s.t. if  $\alpha^0 \in [\alpha_* - \delta, \alpha_* + \delta]$  then  $\{\alpha^k\} \rightarrow \alpha_*$ , with  $p = 2$ .

*Proof.* We are in the hypothesis that the function  $\varphi$  is three times differentiable and we would like to prove that  $\alpha^{k+1} - \alpha_* \rightarrow 0$

We want to compute how much the error is, if compared to the error at the previous iteration. Since  $\varphi(\alpha_*) = 0$  we can use a dirty trick.

1. Since  $\alpha^{k+1} \stackrel{(1)}{=} \alpha^k - \frac{\varphi'(\alpha_*)}{\varphi''(\alpha^k)}$  and  $\varphi'(\alpha_*) \stackrel{(2)}{=} 0$ , we obtain:

$$\begin{aligned}
\alpha^{k+1} - \alpha_* &\stackrel{(1)}{=} \alpha^k - \alpha_* - \frac{(\varphi'(\alpha^k))}{\varphi''(\alpha^k)} \\
&\stackrel{(2)}{=} \alpha^k - \alpha_* - \frac{(\varphi'(\alpha^k) - \varphi'(\alpha_*))}{\varphi''(\alpha^k)} \\
&= \frac{[\varphi'(\alpha^k) - \varphi'(\alpha_*) + \varphi''(\alpha^k) \cdot (\alpha^k - \alpha_*)]}{\varphi''(\alpha^k)}
\end{aligned} \tag{9.1}$$

Where the term inside the square parenthesis is the first order model, centered in  $\alpha_*$  computed in  $\alpha^k$ .

2. Now we can use the first form of Taylor's formula, which says  $\exists \beta \in [\alpha^k, \alpha^*]$  s.t.  $\varphi'(\alpha_*) \stackrel{(3)}{=} \varphi'(\alpha^k) + \varphi''(\alpha^k)(\alpha^k - \alpha_*) + \varphi'''(\beta) \frac{(\alpha^k - \alpha_*)^2}{2}$ . Let's see what happens to  $\alpha^{k+1} - \alpha_*$ :

$$\begin{aligned}\alpha^{k+1} - \alpha_* &\stackrel{(5)}{=} \frac{[\varphi'(\alpha^k) - \varphi'(\alpha_*) + \varphi''(\alpha^k) \cdot (\alpha^k - \alpha_*)]}{\varphi''(\alpha^k)} \\ &\stackrel{(3)}{=} \frac{-\varphi'''(\beta)}{2\varphi''(\alpha^k)} \cdot (\alpha^k - \alpha_*)^2\end{aligned}\tag{9.2}$$

3. We can say that the quantity  $2\varphi''(\alpha^k)$  doesn't become too small and that the numerator  $\varphi'''(\beta)$  doesn't become too big. This is proved since  $\exists \delta > 0$  s.t.  $\varphi''(\alpha) \geq k_2 > 0$  and also  $|\varphi'''(\beta)| \leq k_1 < \infty$ . We can go on bounding the difference between  $\alpha^{k+1}$  and  $\alpha_*$  as follows: for  $\alpha, \beta \in [\alpha_* - \delta, \alpha_* + \delta]$

$$\begin{aligned}|\alpha^{k+1} - \alpha_*| &\stackrel{(6)}{=} \frac{-\varphi'''(\beta)}{2\varphi''(\alpha^k)} \cdot (\alpha^k - \alpha_*)^2 \\ &= |\alpha^{k+1} - \alpha_*| \leq \left[ \frac{k_1}{2k_2} \right] (\alpha^k - \alpha_*)^2\end{aligned}\tag{9.3}$$

We may notice that  $\left[ \frac{k_1}{2k_2} \right]$  may be very large, but it's multiplied for  $(\alpha^k - \alpha_*)^2$ , which means that if we start close enough to  $\alpha^*$  it's ok.

$$\begin{aligned}|\alpha^{k+1} - \alpha_*| &= |\alpha^{k+1} - \alpha_*| \leq \left[ \frac{k_1}{2k_2} \right] (\alpha^k - \alpha_*)^2 \\ &= |\alpha^{k+1} - \alpha_*| \leq \left[ \frac{k_1}{2k_2} \right] (\alpha^k - \alpha_*) \cdot (\alpha^k - \alpha_*)\end{aligned}\tag{9.4}$$

Where  $\left[ \frac{k_1}{2k_2} \right] (\alpha^k - \alpha_*) < 1$ , so  $\frac{k_1(\alpha^k - \alpha_*)}{2k_2} \leq 1 \implies |\alpha^{k+1} - \alpha_*| < |\alpha^k - \alpha_*|$  At this point, if we start from a point  $a^0$  close enough to  $\alpha^*$  (according to this formula) then  $\{\alpha^k\} \rightarrow \alpha_*$  and the convergence is quadratic.

□

In the end, we may conclude that if we start from the right point we converge with a quadratic speed.

**Problem:** This solution makes us compute all the derivatives until the third one. We will now see a solution to this issue.

### 9.0.2 Exact line search: zeroth-order approaches

Can we do line search without computing derivatives at all? Following this approach we can circumvent the problem of the existence of derivatives. In the case of derivatives definite, it's better if we don't have to compute them.

**Key idea:** The more derivatives we have, the smallest number of points we need (second derivative  $\rightarrow$  two points, third derivative  $\rightarrow$  zero points). The opposite holds as well.




FIGURE 9.1: The interesting interval is  $[x_1, x_2]$ , since  $\varphi(x_2) > \varphi(x_1)$  and we are allowed to exclude the interval  $[x_3, +\infty)$  since the value in  $x_3$  is bigger than  $\varphi(x_2)$ .

Obs: We have to minimize a function we know nothing about, except for its value in a point where we compute it. We would like to reduce to the interval which has as extremes the smallest point.

How can we choose these points? The idea is to choose the points that imply that the interval shrinks as fast as possible.

Obs: We have no guarantee that the interval that we are discarding doesn't contain a very deep minimum.

Elegant solution via golden ratio:

$$r = (\sqrt{5} - 1)/2 (\approx 0.618), \quad r : 1 = (1 - r) : r$$




FIGURE 9.2: The relationship between  $r$  and  $1 - r$  is  $r : 1 = (1 - r) : r$ .

---

ALGORITHM 9.2 Pseudocode for non differentiable functions for local minimum detection.

---

```

1: procedure LSGRM( $\varphi, \alpha, \varepsilon$ )
2:    $\alpha_{i-} \leftarrow 0; \alpha_+ \leftarrow \alpha;$ 
3:    $\alpha'_- \leftarrow (1 - \textcolor{blue}{r}) \alpha;$ 
4:    $\alpha'_{+} = \textcolor{blue}{r} \alpha;$ 
5:   while ( $\alpha_+ - \alpha_- \leq \varepsilon$ ) do // note: not the same  $\varepsilon$ 
6:     if  $\varphi(\alpha'_-) > \varphi(\alpha'_+)$  then
7:        $\alpha_- \leftarrow \alpha'_-;$ 
8:        $\alpha'_- \leftarrow \alpha \leftarrow \alpha'_+;$ 
9:        $\alpha'_+ \leftarrow \textcolor{blue}{r}(\alpha_+ - \alpha_-);$ 
10:    else
11:       $\alpha_+ \leftarrow \alpha'_+;$ 
12:       $\alpha'_+ \leftarrow \alpha \leftarrow \alpha'_-;$ 
13:       $\alpha'_- \leftarrow (1 - \textcolor{blue}{r})(\alpha_+ - \alpha_-);$ 
14:    end if
15:   end while
16: end procedure

```

---

### 9.0.3 Inexact line search: Armijo-Wolfe

**Key idea:** Take your favourite line search (it also suggest a simpler one), and run it, but you don't have to wait for the derivative to become zero.




FIGURE 9.3: The dotted line represents the line which has the derivative as slope.

**Definition 9.2** (Armijo condition). Let  $0 < m_1 < (\ll)1$

$$\varphi(\alpha) \leq \varphi(0) + m_1 \alpha \varphi'(0) \quad (A)$$

Problem of Armijo condition: small steps satisfy the armijo condition, but make convergence very slow.




FIGURE 9.4: In the left picture Armijo condition chooses a new line which slope is still negative, but less steep than the original one. In the right one, the ranges where to search are highlighted.

We need another condition, in order to have a lower bound for the step size:

**Definition 9.3** (Goldstein condition). *Let  $m_1 < m_2 < 1$*

$$\varphi(\alpha) \geq \varphi(0) + m_2 \alpha \varphi'(0) \quad (G)$$




FIGURE 9.5: Goldstein condition chooses a new line which slope is still negative, less steep than the original one, but steeper than the one obtained by Armijo.

Problem: the point that satisfies both Goldstein and Armijo may not contain a local minimum.

To circumvent this problem another condition comes to help us.




FIGURE 9.6: Here are the intervals that satisfy both Armijo and Goldstein conditions.

**Definition 9.4** (Wolfe condition). Let  $m_1 < m_3 < 1$

$$\varphi'(\alpha) \geq m_3 \varphi'(0) \quad (W)$$




FIGURE 9.7: On the left Wolfe condition (which chooses derivatives that are substantially zero). On the right part the intervals selected by Wolfe.

Another issue of this conditions is that the derivative in the interval is quite big on the right side.

**Definition 9.5** (Strong Wolfe condition).

$$|\varphi'(\alpha)| \leq m_3 |\varphi'(0)| = -m_3 \varphi'(0)$$

**Fact 9.3.** If  $\varphi'(\alpha) \gg 0$  and  $(A) \cap (W) / (W')$  then all local minima (& maxima) are captured unless  $m_1$  too close to 1 (that's why usually  $m_1 \approx 0.0001$ )




FIGURE 9.8: Strong Wolfe condition.




FIGURE 9.9: If the line is too close to the original one only a few points will satisfy Armijo condition which means that the intersection between Armijo and Wolfe will almost be empty.

The  $m_i$  are like the hyperparameters of machine learning. Less formally, if we choose an  $m_1$  far enough from 1 everything works fine.

**Theorem 9.4.** *Let  $\varphi \in C^1$  and  $\varphi(\alpha)$  bounded below for  $\alpha \geq 0$  then  $\exists \alpha$  s.t.  $(A) \cap (W')$  holds.*

*Proof.*  $l(\alpha) = \varphi(0) + m_1\alpha\varphi'(0)$ ,  $d(\alpha) = l(\alpha) - \varphi(\alpha) \implies d(0) = 0$ ,  $d'(0) = (m_1 - 1)\varphi'(0) > 0$  ( $m_1 < 1$ ) □

0 and  $\bar{\alpha}$  are the two roots of the function  $d$ , so we can use Rolle's theorem, in order to prove that the function  $d$  has a stationary point in the interval  $[0, \bar{\alpha}]$ .




FIGURE 9.10: If the function isn't going to  $-\infty$  the blue line and the function will meet again and we denote the value along the  $x$  axis  $\bar{\alpha}$ .

$$d(\alpha) = \varphi(0) + \alpha(m_1\varphi'(0)) - \varphi(\alpha)$$

$$d'(\alpha) = m_1\varphi'(0) + \varphi'(\alpha)$$

So  $d'(\alpha^*)$  iif  $\varphi'(\alpha^*) = m_1\varphi'(0)$ . Then strong Wolfe requests that  $|\varphi'(\alpha^*)| \leq m_1|\varphi'(0)|$ . How can we find such a point?

---

ALGORITHM 9.3 Pseudocode for backtracking line search.

---

```

1: procedure BLS( $\varphi, \varphi', \alpha, m_1, \tau$ )
2:   while ( $\varphi(\alpha) > \varphi(0) + m_1\alpha\varphi'(0)$ ) do
3:      $\alpha \leftarrow \tau\alpha; \tau < 1;$ 
4:   end while
5: end procedure

```

---

- Fundamental assumption:  $\nabla f$  Lipschitz  $\implies \varphi'$  Lipschitz and  $L$  does not depend on  $x^i$  (**check**)
- Recall:  $\exists \bar{\alpha}$  s.t. (A) holds  $\forall \alpha \in ]0, \bar{\alpha}]$  and  $\varphi'(\bar{\alpha}) > m_1\varphi'(0) > \varphi'(0)$ ;
- $\varphi'$  Lipschitz  $\implies \bar{\alpha}$  is “large” if  $\|\nabla f(x^i)\|$  is:

$$L(\bar{\alpha} - 0) \geq \varphi'(\bar{\alpha}) - \varphi'(0) > (1 - m_1)(-\varphi'(0)) \implies \bar{\alpha} > (1 - m_1) \frac{\|\nabla f(x^i)\|}{L}$$

(recall  $-\varphi'(0) = \|\nabla f(x^i)\|$ );

- Fundamental trick:  $\bar{\alpha}$  can  $\searrow 0$ , but only as fast as  $\|\nabla f(x^i)\|$  does;
- Enough to prove that  $\alpha^i \geq \bar{\alpha}$ , or “not too smaller”.

Now we can prove the following

**Theorem 9.5.** *If  $(A) \cap (W)$  holds  $\forall i$  then either  $\{f(x^i)\} \rightarrow -\infty$  or  $\{\|\nabla f(x^i)\|\} \rightarrow 0$ .*

*Proof.* By **contradiction**, we assume  $-\varphi'(0) = \|\nabla f(x^i)\| \geq \varepsilon > 0 \forall i$ . Then

1.  $(W) \implies \alpha^i \geq \bar{\alpha} > (1 - m_1) \frac{\|\nabla f(x^i)\|}{L} \implies \alpha^i \geq \delta > 0$ ;
2.  $(A) \implies f(x^{i+1}) \leq f(x^i) - m_1 \alpha^i \|\nabla f(x^i)\| \leq f(x^i) - m_1 \delta \varepsilon$ ;
3. So  $\{f(x^i)\} \rightarrow -\infty$  (or  $\{\|\nabla f(x^i)\|\} \rightarrow 0$ ).

□

Backtracking is similar: for simplicity,  $\alpha = 1$  (input)

$\|\nabla f(x^i)\| > \varepsilon \forall i \implies \bar{\alpha} > \delta > 0 \forall i$   
 $h = \min\{k : \tau^{-k} \leq \delta\} \implies \alpha^i \geq \tau^{-h} > 0 \forall i \implies f(x^{i+1}) \leq f(x^i) - m_1 \tau^{-h} \varepsilon \implies \{f(x^i)\} \rightarrow -\infty$  or  $\pmb{\lhd}$ .

# 10 26th of October 2018 — F. Poloni

## 10.1 How to construct a QR factorization

In the previous lecture we introduced the  $QR$  factorization and we defined what an Householder reflector is.

At the end of the lecture we gave a first MatLab implementation of `householder_vector`:

---

ALGORITHM 10.1 Householder vector Matlab implementation.

---

```

1      function [v,s] = householdervector(x)
2          s = norm(x);
3          v = x;
4          v(1) = v(1) - s;
5          v = v / norm(v);

```

---

What's the problem of this algorithm? That the subtraction may create a problem with machine numbers, if  $s$  and  $\|x\|$  are very close. If we take  $\|x\| = -s$  the subtraction becomes an addition, and everything works well.

In the end, we would like to obtain this behaviour for every possible value for  $x$  and  $s$ , so line 2 may be modified as  $s = -\text{sign}(x(1)) * \text{norm}(x)$ .




FIGURE 10.1: If  $x$  is oriented as in the plot it's better if we choose  $-\|x\|e_1$  verse, since it's opposite to  $x$ .

**Example 10.1.** Given  $A = \begin{pmatrix} \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times \end{pmatrix} \in \mathcal{M}(m, m, \mathbb{R})$ , where  $m = 5$ , we would like to calculate the  $QR$  factorization of  $A$ .

STEP 1 : construct a Householder matrix that sends  $A(:, 1)$  (first column of  $A$ ) to a multiple

$$\text{of } e_1. \text{ Then we have } H_1 A = \begin{pmatrix} \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \end{pmatrix}$$

STEP 2 : take  $H_2 \in \mathcal{M}(m-1, m-1, \mathbb{R})$  such that  $H_2 A(2 : \text{end}, 2) = \begin{pmatrix} \times \\ 0 \\ 0 \\ 0 \end{pmatrix}$  and compute:

$$\begin{aligned} \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & & & & \\ 0 & & H_2 & & \\ 0 & & & & \\ 0 & & & & \end{pmatrix} \cdot H_1 A &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & & & & \\ 0 & & H_2 & & \\ 0 & & & & \\ 0 & & & & \\ 0 & & & & \end{pmatrix} \cdot \begin{pmatrix} \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \end{pmatrix} \\ &= \begin{pmatrix} \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & \times & \times & \times \end{pmatrix} \end{aligned} \tag{10.1}$$

And we denote  $Q_2 = \begin{pmatrix} I_{1 \times 1} & 0 \\ 0 & H_2 \end{pmatrix}$ ,  $Q_1 = H_1$ ;

STEP 3 : take  $H_3 \in \mathcal{M}(m-2, m-2, \mathbb{R})$  such that  $H_3 A(3 : \text{end}, 3) = \begin{pmatrix} \times \\ 0 \\ 0 \end{pmatrix}$  and we compute:

$$\begin{aligned} Q_3 \cdot (Q_2 Q_1 A) &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & & & \\ 0 & 0 & & H_3 & \\ 0 & 0 & & & \end{pmatrix} \cdot \begin{pmatrix} \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & \times & \times & \times \end{pmatrix} \\ &= \begin{pmatrix} \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & 0 & \times & \times \\ 0 & 0 & 0 & \times & \times \end{pmatrix} \end{aligned} \tag{10.2}$$

So,  $Q_3 = \begin{pmatrix} I_{2 \times 2} & 0 \\ 0 & H_3 \end{pmatrix}$ ;

STEP 4 : take  $H_4 \in \mathcal{M}(m-3, m-3, \mathbb{R})$  such that  $H_4 A(4 : end, 4) = \begin{pmatrix} \times \\ 0 \end{pmatrix}$  and we compute:

$$\begin{aligned} Q_4 \cdot (Q_3 Q_2 Q_1 A) &= \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & & \\ 0 & 0 & & H_4 & \\ 0 & 0 & & & \end{pmatrix} \cdot \begin{pmatrix} \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & 0 & \times & \times \\ 0 & 0 & 0 & \times & \times \end{pmatrix} \\ &= \begin{pmatrix} \times & \times & \times & \times & \times \\ 0 & \times & \times & \times & \times \\ 0 & 0 & \times & \times & \times \\ 0 & 0 & 0 & \times & \times \\ 0 & 0 & 0 & 0 & \times \end{pmatrix} \end{aligned} \quad (10.3)$$

Where,  $Q_4 = \begin{pmatrix} I_{3 \times 3} & 0 \\ 0 & H_4 \end{pmatrix}$ .

In the end, since  $Q_i$  is an orthogonal matrix and the product of orthogonal matrices is orthogonal,  $Q_1 \cdot Q_2 \cdot Q_3 \cdot Q_4 A = T$ , which is an upper triangular matrix.

**Theorem 10.1** (Product of block matrices). Let  $I \in \mathcal{M}(k, k, \mathbb{R})$ , let  $H_i \in \mathcal{M}(m-k, m-k, \mathbb{R})$  and let  $B_i \in \mathcal{M}(k, k, \mathbb{R})$ ,  $C_i \in \mathcal{M}(k, m-k, \mathbb{R})$  and  $A_i \in \mathcal{M}(m-k, m-k, \mathbb{R})$ , then the product between the two following block matrices is exactly the one showed below.

$$\begin{pmatrix} I & 0 \\ 0 & H_i \end{pmatrix} \cdot \begin{pmatrix} B_i & C_i \\ 0 & A_i \end{pmatrix} = \begin{pmatrix} B_i I & C_i \\ 0 & H_i \cdot A_i \end{pmatrix}$$

*Proof.* It's trivial computation, using the definition of matrix product.  $\square$

### 10.1.1 Matlab implementation

---

ALGORITHM 10.2 First implementation of QR factorization.

---

```

1 function [Q, R] = myqr(A)
2 [m, n] = size(A);
3 Q = eye(m);
4 for j = 1:n
5     v = householder_vector(A(j:end, j));
6     H = eye(length(v)) - 2*v*v';
7     A(j:end,j:end) = H * A(j:end,j:end);
8     Q(:, j:end) = Q(:, j:end) * H;
9 end
10 R = A;

```

---

**Fact 10.2.** The cost of this implementation when  $A$  is a square matrix is  $O(n^3 + (n-1)^3 + \dots + 1^3)$ . If  $A$  is a rectangular matrix, then the computational complexity is  $O(m \cdot n^2 + (m-1) \cdot (n-1)^2 + \dots + (m-n+1)^3)$ .

*Proof.* Line 7 does a matrix product between matrices of size  $n, n-1, \dots, 1$ , so the resulting cost is  $O(m \cdot n^2 + (m-1) \cdot (n-1)^2 + \dots + (m-n+1)^3)$ .  $\square$

We may design a faster algorithm, since  $HA_j = A_j - 2v(v^T A_j)$ .

---

ALGORITHM 10.3 More efficient implementation of QR factorization.

---

```

1  function [Q, A] = myqr(A)
2  [m, n] = size(A);
3  Q = eye(m);
4  for j = 1:n-1
5      [v, s] = householder_vector(A(j:end, j));
6      A(j,j) = s; A(j+1:end, j) = 0;
7      A(j:end, j+1:end) = A(j:end, j+1:end) - ...
8          2*v*(v'*A(j:end, j+1:end));
9      Q(:, j:end) = Q(:, j:end) - Q(:, j:end)*v*2*v';
10 end

```

---

Let's suppose that  $A$  is square matrix, partitioned as:  $A = \begin{pmatrix} A_1 & A_2 \end{pmatrix} = \begin{pmatrix} Q_1 & Q_2 \end{pmatrix} \cdot \begin{pmatrix} R_{1,1} & R_{1,2} \\ 0 & R_{2,2} \end{pmatrix}$

Then we can recover the factorization of  $A_1$  from the factorization of  $A$ , since  $A_1 = Q \cdot R_{11}$ .

**Fact 10.3** (Thin QR factorization). we may replace  $Q \in \mathcal{M}(m, m, \mathbb{R})$  and  $R \in \mathcal{M}(m, m, \mathbb{R})$  with  $Q_1 \in \mathcal{M}(m, n, \mathbb{R})$  and  $R_1 \in \mathcal{M}(n, n, \mathbb{R})$  and the same factorization holds:  $A = QR = Q_1 R_1$ . This is called **thin QR factorization**.

*Proof.*  $A_1 \in \mathcal{M}(m, n, \mathbb{R})$ ,  $A = \begin{pmatrix} Q_1 & Q_2 \end{pmatrix} \cdot \begin{pmatrix} R_1 \\ 0 \end{pmatrix} = Q \cdot R = \begin{pmatrix} Q_1 \\ Q_2 \end{pmatrix} \cdot \begin{pmatrix} R_1 \\ 0 \end{pmatrix} = Q_1 \cdot R_1$ .  $\square$

In order to save space we may work in the following way:

$$Q \cdot B = \begin{pmatrix} 1 & \times & \cdots & \times \\ 0 & & & \\ \vdots & I - 2V_1 V_1^T & & \\ 0 & & & \\ 0 & & & \end{pmatrix} \cdot \begin{pmatrix} 1 & \times & \times & \cdots & \times \\ 0 & 1 & \times & \cdots & \times \\ \vdots & 0 & I - 2V_2 V_2^T & & \\ 0 & \vdots & & & \\ 0 & 0 & & & \end{pmatrix} \cdot \dots \cdot \begin{pmatrix} 1 & \times & \times & \cdots & \times \\ 0 & 1 & \times & \cdots & \times \\ \vdots & 0 & 1 & & \\ 0 & \vdots & 0 & I - 2V_n V_n^T & \\ 0 & 0 & 0 & & \end{pmatrix} \cdot B.$$

### Fun fact

There are some libraries that store the  $v_i$  vectors in the lower part of matrix  $R$  which is upper triangular and has only zeros below the main diagonal.

## 10.2 How to use the thin QR factorization to solve a least squares problem

We would like to solve  $\|Ax - b\| \forall A \in \mathcal{M}(m, n, \mathbb{R})$  and  $\forall B \in \mathbb{R}^n$  where  $m > n$  (a.k.a  $A$  is a tall, thin matrix), through the  $QR$  factorization. We would like to solve  $\min \|Ax - b\|$  through the  $QR$  factorization.

We may write first the  $QR$  factorization of  $A$ , so  $\forall A \in \mathcal{M}(m, n, \mathbb{R})$ ,  $\exists Q \in \mathcal{M}(m, m, \mathbb{R})$ ,  $\exists R \in \mathcal{M}(m, n, \mathbb{R})$  such that  $A = QR$ , where  $Q = \begin{pmatrix} Q_1 & Q_2 \end{pmatrix}$  and  $R = \begin{pmatrix} R_1 \\ 0 \end{pmatrix}$ .

Then,

$$\begin{aligned} \|Ax - b\| &= \|Q^T(Ax - b)\| = \|Q^T Q R x - Q^T b\| \\ &= \|R x - Q^T b\| = \left\| \begin{pmatrix} R_1 \\ 0 \end{pmatrix} x - \begin{pmatrix} Q_1^T \\ Q_2^T \end{pmatrix} b \right\| \\ &= \left\| \begin{pmatrix} R_1 x - Q_1^T b \\ Q_2^T b \end{pmatrix} \right\| \end{aligned} \tag{10.4}$$

How can we pick  $x$  to minimize the norm of  $Ax - b$ ?

Can we choose  $x$  such that  $R_1 x - Q_1^T b = 0$ ? Yes, we can, since this is a linear square system, so  $x = R_1^{-1} Q_1^T b$

$$\|Ax - b\| = \|Q_2^T b\|$$

We used the fact that  $R_1$  is invertible, but is it always true that  $R_1$  is invertible?

**Lemma 10.4.**  $R_1$  is invertible  $\Leftrightarrow A$  has full column rank.

*Proof.*  $A$  has full column rank  $\Leftrightarrow A^T A$  is positive definite  $\Leftrightarrow A^T A$  is positive semidefinite and invertible, but  $A^T A$  is positive semidefinite, so we only need to prove its invertibility.

Let's compute  $QR^T QR = R^T Q^T QR = R^T R = \begin{pmatrix} R_1^T & 0 \end{pmatrix} \cdot \begin{pmatrix} R_1 \\ 0 \end{pmatrix} = R_1^T \cdot R_1$ .

So,  $A^T A$  is invertible  $\Leftrightarrow R_1$  is invertible. □

### Note

$R_1^T R_1$  is the Cholesky factorization of  $A^T A$ .

The computational complexity is asymptotically equal to the one of computing the  $QR$  factorization, since the other operations are cheaper (the product  $Q_1^T \mathbf{b}$  costs  $O(mn)$  and solving the triangular linear system by back-substitution costs  $O(n^2)$ ).

# 11 7th of November 2018 — F.Poloni

## 11.1 Least squares problem with SVD

 Do you recall?

**Tall thin SVD:** A matrix  $A$  can be written as  $A = USV^T$ , where  $U$  is orthogonal,  $S$  is a diagonal matrix and  $V$  is orthogonal as well. In the case of a tall, thin  $A$  the decomposition has the following shape:

$$\begin{pmatrix} & & & \\ & & & \\ U^1 & U^2 & \dots & U^m \\ & & & \end{pmatrix} \cdot \Sigma \cdot \begin{pmatrix} V^1 & V^2 & \dots & V^n \end{pmatrix}^T$$

where

$$\Sigma = \begin{pmatrix} \sigma_1 & & & \\ & \sigma_2 & & \\ & & \ddots & \\ & & & \sigma_n \end{pmatrix}$$

And if we denote  $U_1$  the matrix obtained as the first  $n$  columns of  $U$  we have the tall, thin SVD:  $A = U_1 \cdot \Sigma \cdot V^T$

We would like to see how we can solve a least squares problem through *SVD*:

$$\begin{aligned}
\|Ax - b\| &= \|USV^T x - b\| \quad \leftarrow \text{Def. of SVD} \\
&= \|U^T(USV^T x - b)\| \quad \leftarrow U^T \text{ is orthogonal} \\
&= \|SV^T x - U^T b\| \quad \leftarrow \text{Distributivity + orthogonality} \\
&= \|Sy - U^T b\| \quad \leftarrow y = V^T x \\
&= \left\| \begin{pmatrix} \sigma_1 & & & \\ & \sigma_2 & & \\ & & \ddots & \\ & & & \sigma_n \\ & & & 0 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} - \begin{pmatrix} U^{1T} b \\ U^{2T} b \\ \vdots \\ U^{nT} b \\ U^{n+1T} b \\ \vdots \\ U^{mT} b \end{pmatrix} \right\| \tag{11.1} \\
&= \left\| \begin{pmatrix} \sigma_1 y_1 - U^{1T} b \\ \sigma_2 y_2 - U^{2T} b \\ \vdots \\ \sigma_n y_n - U^{nT} b \\ \sigma_{n+1} y_{n+1} - U^{n+1T} b \\ \vdots \\ \sigma_m y_m - U^{mT} b \end{pmatrix} \right\|
\end{aligned}$$

Where the first  $n$  rows may be assigned to 0 iff  $y_i = -\frac{U^{iT} b}{\sigma_i}$  (if  $\sigma_i \neq 0 \forall i$ ), while the latter  $m - n$  do not depend on  $y$ . This process produces a solution  $y$ , but the variable change may

be inverted, so

$$\begin{aligned}
x &= Vy && \leftarrow \text{Orthogonality of } V \\
&= V^1 y_1 + V^2 y_2 + \cdots + V^n y_n \\
&= V^1 \frac{1}{\sigma_1} U^{1T} b + V^2 \frac{1}{\sigma_2} U^{2T} b + \cdots + V^n \frac{1}{\sigma_n} U^{nT} b \\
&= V \cdot \begin{pmatrix} \frac{1}{\sigma_1} & & & & 0 \\ & \frac{1}{\sigma_2} & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \frac{1}{\sigma_n} \end{pmatrix} \cdot U^T \cdot b \tag{11.2} \\
&= V \cdot \begin{pmatrix} \frac{1}{\sigma_1} & & & & \\ & \frac{1}{\sigma_2} & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \frac{1}{\sigma_n} \end{pmatrix} \cdot U_1^T \cdot b
\end{aligned}$$

Which depends only on the tall, thin SVD.

**Fact 11.1.** *The  $\sigma_i$  are different from 0 iff  $A$  has full column rank.*

*Proof.*  $A$  has full column rank

$$\begin{aligned}
&\Updownarrow \\
A^T A \text{ is invertible} &\Updownarrow \\
&\Updownarrow \\
&\left( \sigma_1^2 \quad \sigma_2^2 \quad \cdots \quad \sigma_n^2 \right) \cdot V^T \text{ is invertible} \\
(USV^T)^T (USV^T) = VS^T U^T USV^T &= V \cdot \begin{pmatrix} \sigma_1^2 & & & & \\ & \sigma_2^2 & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \sigma_n^2 \end{pmatrix} \cdot V^T
\end{aligned}$$

□

**Observation 11.1.** *This lemma also proves that the factorization is also a QR factorization.*

#### Note on Matlab syntax

`svd(A, 0)` and `qr(A, 0)` express that we are only interested in the parts of the factorization without zeros, in case of a tall, thin matrix  $A$ .

We may observe that the computational complexity is  $O(15 \cdot n^3)$  for square matrices, while it's  $O(m \cdot n^2)$  in the tall, thin case.

### 11.1.1 Behaviour in case of zeros as singular values

What happens when there are some zeros as singular values?

 Do you recall?

We may recall that the singular values are ordered on the diagonal in decreasing order (the largest in top left position). From this assumption, we may say that if there are some  $\sigma_i = 0$  then they are in the bottom right part of the matrix.

$$\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r > \sigma_{r+1} = \cdots = \sigma_n = 0$$

$$\text{We also recall the following } \|Ax - b\| = \left\| \begin{pmatrix} \sigma_1 y_1 - U^{1T} b \\ \sigma_2 y_2 - U^{2T} b \\ \vdots \\ \sigma_n y_n - U^{nT} b \\ 0 \cdot y_{n+1} - U^{n+1T} b \\ \vdots \\ 0 \cdot y_m - U^{mT} b \end{pmatrix} \right\|.$$

No matter what value we choose for  $y_{r+1} \cdots y_n$ , the value doesn't change since it's

$$\text{multiplied by 0. Therefore we get infinite solutions of the form } y = \begin{pmatrix} -\frac{U^{1T} b}{\sigma_1} \\ -\frac{U^{2T} b}{\sigma_2} \\ \vdots \\ -\frac{U^{rT} b}{\sigma_r} \\ * \end{pmatrix}$$

We would like to make the solution unique, so we can modify the problem:

- taking the value that minimize the norm:

$$\min_{x \in \arg \min(\|Ax - b\|)} \|x\|. \quad (\text{P2})$$

Note that  $\|x\| = \|y\|$ , because  $x = Vy$ . It follows from the expression of  $y$  that the choice that minimizes its norm is  $y_{r+1} = \cdots = y_n = 0$ .

-

The solution of  $P2$  is given by

$$V y = \begin{pmatrix} V^1 & V^2 & \cdots & V^n \end{pmatrix} \cdot \begin{pmatrix} -\frac{U^{1T} b}{\sigma_1} \\ -\frac{U^{2T} b}{\sigma_2} \\ \vdots \\ -\frac{U^{rT} b}{\sigma_r} \\ 0 \end{pmatrix} = V^{1T} \frac{1}{\sigma_1} U^{1T} b + \cdots + V^{rT} \frac{1}{\sigma_r} U^{rT} b$$

What happens when working with machine precision? Let's make an example where  $r = n - 1$ , so only the last singular value is 0. If the check of  $\sigma_n = 0$  fails,  $\frac{1}{\sigma_n}$  becomes very big. A way to circumvent this problem is to find the linear dependencies between the columns, so that the algorithm works correctly.

## 11.2 Truncated SVD

In many real world setups first singular components correspond to the most prominent features of the dataset, while the smallest ones are fine details and noise. Note, though, that in the sum  $\sum_{i=1}^n V^i \frac{U^{iT} b}{\sigma_i}$  the small singular values may have a large impact, because  $\sigma_i$  is in the denominator.

We can modify the solution to cope with real world data problems:

$$x = \sum_{i=1}^n V^i \frac{U^{iT} b}{\sigma_i} \quad \rightarrow \quad x_{trunc} = \sum_{i=1}^k V^i \frac{U^{iT} b}{\sigma_i}$$

for a certain  $k$ , ignoring small singular values.

Another way to modify the problem is the following.

## 11.3 Tikhonov regularization / ridge regression

The Tikhonov regularization is a smoother version of truncated SVD.

$$x_{Tik} = \arg \min_{x \in \mathbb{R}^n} \|Ax - b\|^2 + \alpha^2 \|x\|^2$$

**Fact 11.2.** *The Tikhonov regularization is equivalent to*

$$x_{Tik} = \arg \min_{x \in \mathbb{R}^n} \left\| \begin{pmatrix} A \\ \alpha I \end{pmatrix} x - \begin{pmatrix} b \\ 0 \end{pmatrix} \right\|^2 \quad (11.3)$$

We show that the two objective functions coincide.

*Proof.*

$$\begin{aligned}
\left\| \begin{pmatrix} A \\ \alpha I \end{pmatrix} x - \begin{pmatrix} b \\ 0 \end{pmatrix} \right\|^2 &= \left\| \begin{pmatrix} Ax \\ \alpha x \end{pmatrix} - \begin{pmatrix} b \\ 0 \end{pmatrix} \right\|^2 \\
&= \left\| \begin{pmatrix} Ax - b \\ \alpha x \end{pmatrix} \right\|^2 \\
&= \|Ax - b\|^2 + \|\alpha x\|^2 \\
&= \|Ax - b\|^2 + \alpha \cdot \|x\|^2
\end{aligned} \tag{11.4}$$

□

**Fact 11.3.** *The solution of the Tikhonov regularization is given by the formula  $x_{Tik} = (A^T A + \alpha^2 I)^{-1} A^T b$ .*

*Proof.* We start by writing the explicit solution of Equation (11.3) using the pseudoinverse

$$\begin{aligned}
\begin{pmatrix} A \\ \alpha I \end{pmatrix}^+ \begin{pmatrix} b \\ 0 \end{pmatrix} &= \left( \begin{pmatrix} A \\ \alpha I \end{pmatrix}^T \begin{pmatrix} A \\ \alpha I \end{pmatrix} \right)^{-1} \begin{pmatrix} A \\ \alpha I \end{pmatrix}^T \begin{pmatrix} b \\ 0 \end{pmatrix} \\
&= \left( \begin{pmatrix} A^T & \alpha I \end{pmatrix} \begin{pmatrix} A \\ \alpha I \end{pmatrix} \right)^{-1} \begin{pmatrix} A^T & \alpha I \end{pmatrix} \begin{pmatrix} b \\ 0 \end{pmatrix} \\
&= (A^T A + \alpha^2 I)^{-1} A^T b.
\end{aligned} \tag{11.5}$$

□

**Fact 11.4.** *We can observe that  $(A^T A + \alpha^2 I)$  is positive definite.*

*Proof.*  $z^T \cdot (A^T A + \alpha^2 I) \cdot z = z^T A^T A z + \alpha^2 z^T z \stackrel{(1)}{=} \alpha^2 z^T z = \alpha^2 \|x\|^2 > 0$ . The equality (1) is obtained because  $z^T A^T A z \geq 0$ , since  $A^T A$  is positive semidefinite. □

**Exercise 11.1.** *Show using the SVD of  $A$  that the Tikhonov / Ridge solution can be written as*

$$x = \sum_{i=1}^n V^i \frac{\sigma_i}{\sigma_i^2 + \alpha^2} U^{i^T} b.$$

When  $\sigma_i \gg \alpha$ ,  $\frac{\sigma_i}{\sigma_i^2 + \alpha^2} \approx \frac{1}{\sigma_i}$ : similar to the ‘true’ solution.

When  $\sigma_i \ll \alpha$ ,  $\frac{\sigma_i}{\sigma_i^2 + \alpha^2} \approx \frac{\sigma_i}{\alpha^2} \approx 0$ : approximately ignoring small singular values.

**Per casa:** fare esercizio.

How can we choose  $k$ ? We don’t know.




FIGURE 11.1: Here is the shape of the formula for the singular values.

## 12 8th of November 2018 — A. Frangioni

### 12.1 Good practice of designing a project

In this lecture we focused on how to implement the gradient method inn Matlab and some good hints were underlined:

- When we have to implement a function the first question we need to ask ourselves is: “how many parameters should the function take?”;
- The interface needs to be designed, i.e. in the case of the gradient method we would like the user to be able to run it although he might not know what a good starting point could be;
- There might be some parameters with the same sematic of hyperparameters (using machine learning terminology), so they need to be adjusted in order to specify the algorithm;
- Another thing that should be taken into consideration is that sometimes, when measuring the number of function calls we need to count also the calls to functions inside a single step;
- It’s important to check the conditions that should be satisfied by the input data for the theoretical coherence. In case of not valid input an error message should be returned;
- Another important thing to be done is building a set of test functions, possibly limit case for the problem, to see how the algorithm works;
- `diff(f, x)` calculates the gradient of the function  $f$  in the variable  $x$ . If we want to get a simplified version, we may run `simplify(diff(f,x))`;
- A very nice tool to compute derivatives for arbitrary functions is **Adigator** (click here).

# 13 9th of November 2018 — F. Poloni

## 13.1 Conditioning

Two lectures ago we introduced the QR factorization to solve least squares problems and we noticed that it has a computational complexity which is much worse than the normal equations method.

Why did we introduce the QR factorization to solve the least squares problem, then? Although it's more complex computationally speaking, it's much better than the normal equation for what concerns accuracy. Let's see why through an example:

**Example 13.1.** Let  $A \in \mathcal{M}(4, 3, \mathbb{R})$  s.t.

$$A = \begin{pmatrix} 1 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 4 \\ 1 & 2 & 3 + 10^{-8} \end{pmatrix}$$

In this case the normal equations method doesn't even find the order of magnitude correctly.

At this point we may introduce the problem of **sensitivity**:

**Definition 13.1** (Sensitivity). We call **sensitivity** the measure of how much the output of a problem changes when we perturb its input.

As an example, let  $f(x, y) = x + 2y$ . If we perturb the second parameter of  $f$  as follows  $\tilde{y} = y + \delta$ , we can compute the variation in the output value of the function  $v$  as

$$v = f(x, \tilde{y}) - f(x, y) = x + 2(y + \delta) - (x + 2y) = 2\delta.$$

A good example of this behaviour is the temperature of water coming from the shower: in particular, when we rotate little the knob the water becomes too cold or too hot very fast. This function is plotted in Figure 13.1.

**Definition 13.2** (Absolute condition measure). The **absolute condition number** of a function  $f$  is the **maximum** possible output change / input change ratio in the limit for a **small** change of the input.

$$\kappa_{abs}(f, x) = \lim_{\varepsilon \rightarrow 0} \sup_{|\tilde{x}-x| \leq \varepsilon} \frac{|f(\tilde{x}) - f(x)|}{|\tilde{x} - x|}.$$

We would like to focus on what this definition means.

Why are we interested in the limit of a very small change?

If we zoom-in a continuous function it gets basically linear (key idea of derivative) and then the ratio between the difference on the outputs and the one of the inputs is approximatively the derivative, as shown in Figure 13.2.

We take a point  $x$  and we consider a ball of radius  $\varepsilon$  and we compute the change in the output over the change in the input, then we take the maximum.




FIGURE 13.1: Geometric idea of temperature of the water in the shower




FIGURE 13.2: Geometric idea behind the derivative, as “zoom” of the function in a certain point.




FIGURE 13.3: Geometric idea behind absolute condition number.

**Example 13.2.** Let  $f(x) = x^2$ .

If we perturb the input  $x$  to  $x + \delta$ , then  $f(\tilde{x}) = (x + \delta)^2 = x^2 + 2x\delta + \delta^2$ , then we obtain

the ratio

$$r = \frac{|f(\tilde{x}) - f(x)|}{|\tilde{x} - x|} = \frac{|2x\delta + \delta^2|}{|\delta|} = |2x + \delta|$$

If we denote  $\varepsilon$  the radius of the ball, we obtain the following

$$\max_{\substack{|\delta| < |\varepsilon| \\ \tilde{x} \in B(x, \varepsilon)}} r = |2x| + |\varepsilon|$$

then

$$\lim_{\varepsilon \rightarrow 0} \max_{|\delta| < |\varepsilon|} r = \lim_{\varepsilon \rightarrow 0} |2x| + |\varepsilon| = |2x|$$

**Example 13.3.** It's more interesting to see a multivariate function:

Let  $f(x) = x^T Q x$ , for instance for  $x \in \mathbb{R}^2$  so that we can plot its graph in  $\mathbb{R}^3$ .




FIGURE 13.4: Paraboloid

We shall take a general example where the cross-section are ellipses, so that there is a direction of faster and slower ascent; this is not just a circular "cup" seen in perspective. Note that these directions of faster ascent and lower ascent correspond to the eigenvectors of the matrix  $Q$ .

In this case the absolute condition number is  $\lim_{\varepsilon \rightarrow 0} \max_{\tilde{x} \in B(x, \varepsilon)} \frac{\|f(\tilde{x}) - f(x)\|}{\|\tilde{x} - x\|}$ , and one can see that the output/input ratio varies with the direction in which  $\tilde{x}$  is, so we have to take a maximum in the whole ball  $B(x, \varepsilon)$ .

At this point, an observation is mandatory: **any** absolute measure doesn't take into account the values of the function in other points, so we want to define the following

**Definition 13.3** (Relative error). The **relative error** of an approximation  $\tilde{x}$  to a quantity  $x$  is  $\frac{\|\tilde{x} - x\|}{\|x\|}$ .

Here are some examples of good and bad accuracy:




FIGURE 13.5: Level curves of a quadratic function (“seen from above”).

- $\frac{|\tilde{x}-x|}{|x|} \approx 1$ : **very bad** accuracy; it's just a number with the same order of magnitude.
- $\frac{|\tilde{x}-x|}{|x|} \approx 10^{-3}$ : about 3 correct significant digits.
- $\frac{|\tilde{x}-x|}{|x|} \approx 10^{-16}$ : about 16 correct digits; we **can't do better** typically (with double precision arithmetic).

**Definition 13.4** (Relative condition number). *The relative condition number of a function  $f$  is defined as*

$$\kappa_{rel}(f, \mathbf{x}) = \lim_{\delta \rightarrow 0} \sup_{\frac{\|\tilde{\mathbf{x}}-\mathbf{x}\|}{\|\mathbf{x}\|} \leq \delta} \frac{\frac{\|f(\tilde{\mathbf{x}})-f(\mathbf{x})\|}{\|f(\mathbf{x})\|}}{\frac{\|\tilde{\mathbf{x}}-\mathbf{x}\|}{\|\mathbf{x}\|}} = \kappa_{abs}(f, \mathbf{x}) \frac{\|\mathbf{x}\|}{\|f(\mathbf{x})\|},$$

i.e., we replace the absolute error  $\|\tilde{\mathbf{x}} - \mathbf{x}\|$  with the relative error.

### 13.1.1 Conditioning of linear systems

At this point we would like to compute the condition number of solving a linear system, i.e., the condition number of the function  $f(A, b) = A^{-1}b$ , perturbing the inputs  $A$  and  $b$ , one at a time.

**PERTURBING  $b$**  We want to compute the limit of the relative error  $\frac{\|f(A, \tilde{b}) - f(A, b)\|}{\|f(A, b)\|}$ , so we set  $x = A^{-1}b$  and  $\tilde{x} = A^{-1}\tilde{b}$ , and we estimate *output error*  $= \frac{\|\tilde{x} - x\|}{\|x\|} = ?$

1.

$$\begin{aligned}
\|\tilde{x} - x\| &= \|A^{-1}\tilde{b} - A^{-1}b\| \\
&= \|A^{-1}(\tilde{b} - b)\| \\
&\leq \|A^{-1}\| \cdot \|\tilde{b} - b\|
\end{aligned} \tag{13.1}$$

2. Since  $\|b\| = \|Ax\| \leq \|A\| \cdot \|x\|$  we have  $\frac{\|\tilde{x}-x\|}{\|x\|} \leq \|A^{-1}\| \cdot \|A\| \cdot \frac{\|\tilde{b}-b\|}{\|b\|}$

In the end, since  $input\ error = \frac{\|\tilde{b}-b\|}{\|b\|}$  we obtain

$$\kappa_{rel}(f, x) = \lim_{\varepsilon \rightarrow 0} \frac{output\ error}{input\ error} \leq \lim_{\varepsilon \rightarrow 0} \|A^{-1}\| \cdot \|A\| = \|A^{-1}\| \cdot \|A\|$$

We denote  $\kappa(A) = \|A^{-1}\| \cdot \|A\|$  the **condition number of  $A$** ;

**PERTURBING  $A$**  Given  $Ax = b$  we obtain  $(A + \Delta_A) \cdot (x + \Delta_x) = b$ , where  $\tilde{A} = A + \Delta_A$  and  $\tilde{x} = x + \Delta_x$ . Then we can expand as follows

$$A\tilde{x} + \Delta_A \cdot x + A \cdot \Delta_x + \Delta_A \cdot \Delta_x = b$$

We can stop taking into account  $\Delta_A \cdot \Delta_x$ , since it's a sort of second order term ( $\Delta_A \cdot \Delta_x = o(\|\Delta_A\| \cdot \|\Delta_x\|)$ ), so we get the following

$$\Delta_A \cdot x + A \cdot \Delta_x = 0$$

$$\Delta_x = -A^{-1} \cdot \Delta_A \cdot x$$

then  $\|\Delta_x\| \leq \|A^{-1}\| \cdot \|\Delta_A\| \cdot \|x\|$ , which implies  $\frac{\|\Delta_x\|}{\|x\|} \leq \|A^{-1}\| \cdot \frac{\|\Delta_A\|}{\|A\|}$ .

We obtain that *relative output error  $\leq \kappa(A) \cdot relative\ input\ error$* .

We only proved an inequality, but it turns out that it is tight: for every  $A$  and  $b$  there is a possible choice of the perturbation  $\tilde{x}$  that attains equality.

In the end, in both cases, the error in the output is the error in the input (namely  $b$  or  $A$ ) times the condition number.

## 13.2 Condition number, SVD, and distance to singularity

**Fact 13.1.**  $\kappa(A) = \frac{\sigma_1}{\sigma_n}$ , i.e.,  $\kappa(A)$  is the ratio between the smallest and the largest singular value.

So we can say that if a matrix is close to a singular matrix, then its condition number is going to be large.

*Proof.* Let  $A = USV^T$ , then  $\|A\| = \|U \cdot S \cdot V^T\| = \|S\| = \sigma_1$ , since

$$S = \begin{pmatrix} \sigma_1 & & & \\ & \sigma_2 & & \\ & & \ddots & \\ & & & \sigma_n \end{pmatrix} \text{ and } \sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n.$$

It's also true that  $\|A^{-1}\| = \|(USV)^{-1}\| = \|VS^{-1}U^T\| = \|S^{-1}\| = \sigma_n$ , since

$$S^{-1} = \begin{pmatrix} \frac{1}{\sigma_1} & & & \\ & \frac{1}{\sigma_2} & & \\ & & \ddots & \\ & & & \frac{1}{\sigma_n} \end{pmatrix} \text{ and } \sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n.$$

In the end  $\kappa(A) = \frac{\|A\|}{\|A^{-1}\|} = \frac{\sigma_1}{\sigma_n}$

□

**Fact 13.2.** *The relative distance between  $A$  and the closest singular matrix is  $\frac{1}{\kappa(A)}$ .*

 Do you recall?

**Eckart-Young theorem:** the closest matrix to  $A$  that has rank  $\leq n-1$  is

$$\hat{A} = U \begin{pmatrix} \sigma_1 & & & \\ & \sigma_2 & & \\ & & \ddots & \\ & & & \sigma_n \\ & & & 0 \end{pmatrix} \cdot V^T$$

*Proof.*

$$bau \tag{13.2}$$

$$\begin{aligned}
\|A - \hat{A}\| &= \left\| U \cdot \left( \begin{pmatrix} \sigma_1 & & & \\ & \sigma_2 & & \\ & & \ddots & \\ & & & \sigma_n \end{pmatrix} - \begin{pmatrix} \sigma_1 & & & \\ & \sigma_2 & & \\ & & \ddots & \\ & & & \sigma_{n-1} & 0 \end{pmatrix} \right) \cdot V^T \right\| \\
&= \left\| U \cdot \begin{pmatrix} 0 & & & \\ & \ddots & & \\ & & \ddots & \\ & & & 0 & \sigma_n \end{pmatrix} \cdot V^T \right\| \\
&= \sigma_n
\end{aligned} \tag{13.3}$$

Thus,  $\|A - \hat{A}\| = \sigma_n$ . We know already that  $\|A\| = \sigma_1$ , so we just need to take the ratio.  $\square$

We analyzed the conditioning of linear systems, but the main problem we want to study in this course is least squares problem.

### 13.3 Conditioning of least squares problem

We need two quantities to be able to measure the conditioning of least squares problem:

★  $\kappa(A)$  Let  $A \in \mathcal{M}(m, n, \mathbb{R})$ , with  $m > n$  (tall, thin  $A$ ). We define  $\kappa(A) = \frac{\sigma_1}{\sigma_n}$ . We take this as a definition of  $\kappa(A)$ . Note that we cannot use the other definition  $\kappa(A) = \|A\| \|A^{-1}\|$ , since  $A^{-1}$  does not exist for a non-square  $A$ . However, one can verify that  $\|A\| \cdot \|A^\dagger\| = \frac{\sigma_1}{\sigma_n} = \kappa(A)$ , where  $A^\dagger$  is the pseudoinverse.

**Observation 13.1.** Note that  $\frac{1}{\kappa(A)}$  is the relative distance to the closest  $\hat{A}$  without full column rank.

★  $\theta$  The second quantity needed is the angle between  $Ax$  and  $b$ , see Figure 13.6.  $\theta = \arccos \frac{\|Ax\|}{\|b\|}$

Now we can express the theorem:

**Theorem 13.3** (Trefethen, Bau, Theorem 18.1). Consider the linear least squares problem  $\min \|Ax - b\|$ , with  $A \in \mathbb{R}^{m \times n}$  with full column rank. Its relative condition number with respect to the input  $b$  is

$$\kappa_{rel,b \rightarrow x} \leq \frac{\kappa(A)}{\cos \theta},$$




FIGURE 13.6: The triangle in the figure (the one whose cathets are  $Ax$  and  $b - Ax$ ) is a square triangle.

and with respect to  $A$  it is

$$\kappa_{rel, A \rightarrow x} \leq \kappa(A) + \kappa(A)^2 \tan \theta,$$

where  $\theta$  is the angle such that  $\cos \theta = \frac{\|Ax\|}{\|b\|}$ .

At this point we have two condition numbers and they both depend on  $\kappa(A)$  and  $\theta$ .

### Observation 13.2.

SPECIAL CASE 1:  $\theta \approx 90^\circ$  We can see from the figure that a big change of  $b$  induces a small perturbation of  $Ax$ . No matter what the conditioning of  $A$  is, a small (relative) perturbation in  $b$  can change a large (relative) perturbation in  $x$  and  $Ax$ , see Figure 13.7.




FIGURE 13.7: Special case 1

SPECIAL CASE 2:  $\theta \approx 0^\circ$  When  $b$  is almost in plane with  $Im(x)$ . In this case  $cond \approx \kappa(A)$ , see Figure 13.8.




FIGURE 13.8: Special case 2.

GENERAL CASE:  $\theta$  FAR FROM  $0^\circ$  AND  $90^\circ$  In the more general case,  $cond \approx \kappa(A)^2$ .

## 14 14th of November 2018 — A. Frangioni

### ★ Mantra

If you want better convergence, use a better model.

So far we chose the direction for the step as  $d^i = -\nabla f(x^i) \cdot \|\nabla f(x^i)\|$ , in particular this is the direction where the decrease of the function is maximum.

We introduced the **convergence argument** which says that in proximity of the stationary point of the linear model the norm of the gradient goes to 0. Formally,  $\varphi'(0) = \frac{\partial f}{\partial d^i}(x^i) = \langle d^i, \nabla f(x^i) \rangle = -\langle \nabla f(x^i), \nabla f(x^i) \rangle = -\|\nabla f(x^i)\|^2$ , which implies that  $\|\nabla f(x)\| \rightarrow 0$  when  $\varphi'(0) \rightarrow 0$ .

Can we take another direction which isn't the opposite of the gradient and have that the same argument holds? Yes, for example if we choose as direction a rotation of the opposite of the gradient, the value of  $\varphi'$  is then the cosine of the angle of the rotation times the opposite of the norm of the gradient. It's trivial to observe that there are infinite angles that could be chosen, so we have a lot of flexibility.

Note that the angle between  $d^i$  and the gradient shouldn't be too close to 90°, otherwise the cosine would get approximately 0..

**Theorem 14.1** (Zoutendijk). *Let  $f \in C^1$ ,  $\nabla f$  Lipschitz and  $f$  bounded below.*

*If  $(A) \cap (W')$  then  $\sum_{i=1}^{\infty} \cos^2(\theta^i) \|\nabla f(x^i)\|^2 < \infty$ .*

If we have a positive infinite sequence and we have that the corresponding series converges to a number, than the limit of the sequence is going to 0 reasonably fast.

If we choose an angle which is bounded below then the norm of the gradient has to converge very fast. More formally,

**Fact 14.2.**  $\cos(\theta^i) \geq \varepsilon > 0 \implies \|\nabla f(x^i)\| \rightarrow 0$

*Proof.* From the observation above, we may recall that the  $n$ -th term of the Zoutendijk sum should be approximately 0  $\forall n > \tilde{n}$ , which means that one between  $\cos(\theta^i)$  and  $\|\nabla f(x^i)\|^2$  should be zero.

The proof is obtained from the fact that it can't be the cosine, because it's bounded below.  $\square$

### 14.1 Taylor method

At this point we assume that the function is perfectly convex (Hessian strictly positive definite).

**Theorem 14.3.** *Let  $f$  be a function such that  $\nabla^2 f(x^i) \succ 0$ . Then the second order model  $Q_{x^i}(y)$  admits a minimum.*

*Proof.* For simplicity of notation let us forget about the index  $i$  of  $x^i$ . The Taylor expansion is the following:

$$\begin{aligned}
Q_x(y) &= f(y) + \langle \nabla f(x), y - x \rangle + \frac{1}{2}(y - x)^T \nabla^2 f(x)(y - x) \\
&= f(x) + \langle \nabla f(x), y \rangle - \langle \nabla f(x), x \rangle + \\
&\quad + \frac{1}{2} \left( y^T \nabla^2 f(x) y - 2x^T \nabla^2 f(x) y + x^T \nabla^2 f(x) y \right) \\
&\stackrel{*}{=} \langle \nabla f(x) + x^T \nabla^2 f(x), y \rangle + \frac{1}{2} y^T \nabla^2 f(x) y
\end{aligned} \tag{14.1}$$

Where  $\stackrel{*}{=}$  is given by the fact that there are some constant terms.  
 $qy + \frac{1}{2}y^T Qy \Leftrightarrow q + Qy = 0$  so

$$\nabla f(x) - x^T \nabla^2 f(x) + \nabla^2 f(x)y = 0 \nabla^2 f(x)y = \nabla f^2(x)x - \nabla f(x)y = x - [\nabla^2 f(x)]^{-1} \nabla f(x)$$

□

---

**Corollary 14.4.** Newton's direction is  $d^i \leftarrow -[\nabla^2 f(x^i)]^{-1} \nabla f(x^i)$  (just  $\mathbb{R}^n$  version).

Scrivere  
meglio la  
dim

The direction  $d$  is obtained taking the opposite of the inverse of the Hessian times the gradient of the function

**Observation 14.1.** We need the Hessian to be invertible, which isn't true in general.

We need to solve a non linear equation, namely a system of equations. A way to circumvent this problem is putting the gradient to 0, so we can write the Taylor form of the gradient and solve a linear equation which is  $\nabla f(x) \approx \nabla f(x^i) + \nabla^2 f(x^i)(x - x^i)$ .

**Theorem 14.5.** Let  $f$  be a function s.t.  $f \in C^3$ ,  $\nabla f(x_*) = 0$  and  $\nabla^2 f(x_*) \succ 0$ . Then  $\exists \mathcal{B}(x_*, r)$  s.t.  $x^1 \in \mathcal{B}$ , which implies  $\{x^i\} \rightarrow x_*$  quadratically.

We may observe that the direction of the Newton method is good not only when we are close to the minimum, but also when we are far.

The scalar product should be negative and it is so, but we also need to ensure that the scalar product isn't too close to 0. When is it that this condition is satisfied? When the function is "reasonable".

Let us now introduce what we mean by "reasonable": a function  $f$  such that  $uI \preceq \nabla^2 f \preceq LI$ , which implies that the function is strongly convex, in other words that the eigenvalues of the Hessian don't get too close to zero (numerically speaking).

**Theorem 14.6.** Let  $f$  be a function that satisfies  $uI \preceq \nabla^2 f \preceq LI$  and  $\cos(\theta^i) \leq -\lambda^n / \lambda^1 \leq -u/L$ . Then the method converges.

*Proof.*

STEP 1 From the definition of  $d^i$  we have that

$$d^i = -[\nabla^2 f(x^i)]^{-1} \nabla f(x^i)$$

or, equivalently,

$$\nabla^2 f(x^i) d^i = -\nabla f(x^i)$$

which implies

$$d^i \nabla f(x^i) = -(d^i)^T \nabla^2 f(x^i) d^i \leq -\lambda^n \|d^i\|^2$$

STEP 2 Since we have that  $\cos(\theta^i) = d^i \nabla f(x^i) \cdot (\|x^i\| \cdot \|\nabla f(x^i)\|) \leq \delta < 0$ , we want to bound the norm of the gradient:

$$\|\nabla f(x^i)\| = \|\nabla^2 f(x^i) d^i\| \leq \|\nabla^2 f(x^i)\| \cdot \|d^i\| = \lambda^1 \|d^i\|$$

which implies

$$\cos(\theta^i) \leq -\lambda^n / \lambda^1 \leq -u/L$$

□

**Theorem 14.7.** Let  $f$  be a function that satisfies  $uI \preceq \nabla^2 f \preceq LI$  and  $\cos(\theta^i) \leq -\lambda^n / \lambda^1 \leq -u/L$ . Then the method not only converges, but we also have that for some iteration onwards,  $\alpha^i = 1$  always satisfy (A).

*Proof.*

$$\begin{aligned} f(x^i + d^i) &= f(x^i) + d^i \nabla f(x^i) + \frac{1}{2} \cdot (d^i)^T [\nabla^2 f(x^i)] d^i + R(\|d^i\|) \\ &= f(x^i) - \nabla f(x^i)^T [\nabla^2 f(x^i)]^{-1} \nabla f(x^i) + \\ &\quad + \frac{1}{2} \nabla f(x^i)^T [\nabla^2 f(x^i)]^{-1} \nabla f(x^i) + R(\|d^i\|) \\ &= f(x^i) - \frac{1}{2} \cdot \nabla f(x^i)^T \cdot [\nabla^2 f(x^i)]^{-1} \nabla f(x^i) + R(\|d^i\|) \\ &= f(x^i) + \frac{1}{2} \langle \nabla f(x^i), d^i \rangle + R(\|d^i\|) \end{aligned} \tag{14.2}$$

□

It can be proved that the convergence is superlinear.

If we start with a step size of 1, we end up in a situation in which the line search isn't computed when we are close to the minimum.

This works under the assumption that the eigenvalues are bounded both above and below (deriving from the bounds on the Hessian).

### 14.1.1 Interpretation of Newton method

Let us consider the Newton method from a different point of view. We can construct a different space where the gradient method coincides with the Newton method. Given  $R$  which is not singular, we make a variable change ( $y = Rx$ ) — which is possible given that  $R$  is non singular — and we get  $f_y(y) = \frac{1}{2}y^T I y + qR^{-1}y$ , which has as an Hessian the identity matrix, which is the optimal matrix for convergence in Newton method.

Formally, the descending direction  $d_y$  is computed as follows:  $d_y = -\nabla f_y(y) = -y - R^{-1}q$ . Since we chose 1 as step size we obtain that  $\nabla f_y(y+d_y) = \nabla f_y(y-y-R^{-1}q) = \nabla f_y(-R^{-1}q) = 0$ .

It takes only one iteration, because all the eigenvalues are 1 so the ratio between the greatest and the smallest is 1 and the subtraction is 0.

If we do the inverse operation ( $x = R^{-1}y$ ) to the direction we get the direction in  $x$  variable.

#### Problem:

We made a lot of assumptions on the Hessian, without the ones there's no guarantee that we are moving on a descending direction. How can we relax these constraints?

### 14.1.2 Non convex case

If the Hessian isn't positive definite we may take something which is close to the Hessian, which has the bounding property we used before (eigenvalues bounded below and above). In truth, there is no reason to choose exactly the opposite of the inverse of the Hessian times the gradient for the direction, we may use another matrix which is strictly positive definite and has more or less the same properties of the Hessian.

In particular, given an Hessian that has some negative eigenvalues we can sum to it a multiple of the identity matrix:  $H^i = \nabla^2 f(x^i) + \varepsilon^i I \succ 0$ . We iterate this procedure until the resulting matrix is strictly positive definite.

How to compute  $\varepsilon$  numerically? How much larger should  $\varepsilon$  be? We also want the smallest eigenvalue to be not too close to 0.

$\varepsilon = \max\{0, \delta - \lambda^n\}$  for “appropriately chosen smallish  $\delta$ ”, in other words we want all the eigenvalues to be at least  $\delta$ .

The reasons behind the choice of  $\delta$  (not too small) are both numerical (any double  $\leq 1e-16$  “is 0”) and algorithmical (if  $\lambda^n(\nabla^2 f(x^i) + \varepsilon I)$  “very small” then the axes of  $S(Q_{x^i}, \cdot)$  are “very elongated”).

It can be proved that the  $\varepsilon$  we chose is the solution to an optimization problem:  $\min\{\|H - \nabla^2 f(x^i)\| \mid H \succeq \delta I\}$ . The choice for  $\delta$  in the MatLab code is  $10^{-6}$ .

**Observation 14.2.** Note that these constraints are important and we will get back to them later on in the course.

Could we use a different norm instead of the 2-norm? Yes, for example we can use Frobenius norm, changing a bit the algorithm, but we still get convergence. We would need

to solve  $\min\{\|H - \nabla^2 f(x^i)\|_{\textcolor{red}{F}} \mid H \succeq \delta I\}$ , which is performed in two steps:

1. compute spectral decomposition  $\nabla^2 f(x^i) = H \Lambda H^T$
2.  $H^i = H \bar{\Lambda} H^T$  with  $\bar{\gamma}^i = \max\{\lambda^i, \delta\}$

In both cases,  $\{x^i\} \rightarrow x_*$  with  $\nabla^2 f(x_*) \succeq \delta I$ , which implies  $varepsilon^i = 0$  and  $H^i = \nabla^2 f(x^i)$  eventually. It holds also that we have superlinear convergence if “ $H^i$  looks like  $\nabla^2 f(x^i)$  along  $d^i$ ”, formally  $\lim_{i \rightarrow \infty} \|(H^i - \nabla^2 f(x^i))d^i\| \|d^i\| = 0$ .

### Computational complexity

We still need to compute eigenvalues, which takes  $O(n^3)$ , which is too much if we are in multidimensional spaces.

As a closing observation we may notice that Newton method is very fast to go to a local minimum and this may represent a problem, because it misses global minima.

# 15 15th of November 2018 — F. Poloni

## 15.1 Stability of algorithms

In this lecture we will try to answer the question: “Is our algorithm (using floating point) going to compute a good approximation of the answer?”

It is related to sensitivity / conditioning but different. Depends on how we perform the computation.



Do you recall?

Computers work with IEEE arithmetic and the basic idea is the following.

TL;DR: floating point numbers are numbers in base-2 scientific (exponential) notation.

`double` (64-bit numbers):

$$\pm 1. \underbrace{01001011101\dots101}_{52 \text{ binary digits}} \cdot 2^{\pm \underbrace{101\dots01}_{10 \text{ binary digits}}}.$$

We use 1 bit for the sign, 52 bits for the “mantissa” and 11 bits for the exponent and its sign.

Some of these combinations of bits are reserved for special numbers, e.g. Inf and NaN, -0.

This system is subject to approximation errors, exactly like the “usual” decimal arithmetic: for example, if we do  $\frac{1}{3} = 0.33333\dots$  and if we do  $\frac{1}{3} + \frac{1}{3} + \frac{1}{3} = 0.99999\dots \neq 1$ .

Whenever we store a number on a computer we need to approximate it, using the “representable numbers”, as shown in Figure 15.1.




FIGURE 15.1: We have  $2^{52}$  equispaced numbers between  $\frac{1}{2}$  and 1 and between 1 and 2, and also  $2^{52}$  between 2 and 4 and so on and so forth, so we have the same number of integers, although the space is enlarging.

Not all numbers are exactly representable, take 0.1 (decimal). It’s a periodic number when written in binary, hence we can’t represent it exactly as a machine number.

**Definition 15.1** (Error bound). *For each  $x \in \pm[10^{-308}, 10^{308}]$ , there is an exactly representable number  $\tilde{x}$  such that  $\frac{|\tilde{x}-x|}{|x|} \leq u$ , with  $u = 2^{-52} \approx 2 \cdot 10^{-16}$ .*

Let us assume that we have the best possible algorithm that returns  $\tilde{y} = f(\tilde{x})$  which is the best representation of  $f(\tilde{x})$ , then

$$\begin{aligned}\frac{|\tilde{y} - y|}{|y|} &\leq \kappa_{rel}(f, x) \frac{|\tilde{x} - x|}{|x|} + o\left(\frac{|\tilde{x} - x|}{|x|}\right) \\ &\leq \kappa_{rel}(f, x) u + o(u).\end{aligned}$$

In practice we may ignore  $o(u)$ , since it's small o of  $2^{-16}$ .

When we are approximating a result, we are approximating it up to a relative error of order of magnitude  $10^{-16}$ .

$$a \oplus b = (a + b)(1 + \delta), \quad |\delta| \leq u.$$

$\Updownarrow$

$$\frac{|(a \oplus b) - (a + b)|}{|a + b|} = \delta$$

We would like to compute the error on the function  $f(a, b) = a^T b$ .

$$a = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}, b = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}, \text{ so } f(a, b) = a_1 \cdot b_1 + a_2 \cdot b_2 + a_3 \cdot b_3$$

Denoting  $\odot$  the approximated product and  $\oplus$  the approximated sum on a computer, we can write the following:

$$\begin{aligned}\text{computer result} &= a_1 \odot b_1 \oplus a_2 \odot b_2 \oplus a_3 \odot b_3 \\ &= \left[ [a_1 b_1 \cdot (1 + \delta_1) + a_2 b_2 \cdot (1 + \delta_2)] \cdot (1 + \delta_3) + a_3 b_3 \cdot (1 + \delta_4) \right] \cdot (1 + \delta_5) \\ &= a_1 b_1 \cdot (1 + \delta_1) \cdot (1 + \delta_3) \cdot (1 + \delta_5) + a_2 b_2 \cdot (1 + \delta_2) \cdot (1 + \delta_3) \cdot (1 + \delta_5) \\ &\quad + a_3 b_3 \cdot (1 + \delta_4) \cdot (1 + \delta_5) \\ &= a_1 b_1 (1 + \delta_1 + \delta_3 + \delta_5 + O(u^2)) \\ &\quad + a_2 b_2 (1 + \delta_2 + \delta_3 + \delta_5 + O(u^2)) \\ &\quad + a_3 b_3 (1 + \delta_4 + \delta_5 + O(u^2)) \\ &\approx a_1 b_1 (1 + \delta_1 + \delta_3 + \delta_5) + a_2 b_2 (1 + \delta_2 + \delta_3 + \delta_5) + a_3 b_3 (1 + \delta_4 + \delta_5)\end{aligned}\tag{15.1}$$

Where  $O(u^2)$  comes from the summation of  $\delta_i \delta_j$ , for some  $i, j$  and allows us to do an approximation up to second order terms of precision.

The absolute error then is

$$\begin{aligned}
\text{Err}_a &= |a_1 b_1 (\cancel{1} + \delta_1 + \delta_3 + \delta_5) + a_2 b_2 \cdot (\cancel{1} + \delta_1 + \delta_3 + \delta_5) + a_3 b_3 \cdot (\cancel{1} + \delta_4 + \delta_5) + \cancel{a_1 b_1} + \cancel{a_2 b_2} + \cancel{a_3 b_3}| \\
&\stackrel{(1)}{\leq} |a_1 b_1| \cdot 3u + |a_2 b_2| \cdot 3u + |a_3 b_3| \cdot 2u \\
&\leq (|a_1 b_1| + |a_2 b_2| + |a_3 b_3|) \cdot 3u
\end{aligned} \tag{15.2}$$

Where  $\stackrel{(1)}{\leq}$  follows from the observation that  $|\delta_i| \leq u$ .

The result that we can obtain is weaker than we would have expected: if  $a_i b_i \geq 0, \forall i = 1, 2, 3$ , then

$$\frac{|\text{computer result} - (a_1 b_1 + a_2 b_2 + a_3 b_3)|}{a_1 b_1 + a_2 b_2 + a_3 b_3} \leq 3u,$$

which means that the algorithm is stable.

However, if  $a_1 b_1, a_2 b_2$  and  $a_3 b_3$  have different signs, then we can bound the error not with  $a_1 b_1 + a_2 b_2 + a_3 b_3$ , but only with  $|a_1 b_1| + |a_2 b_2| + |a_3 b_3|$ . This might be a lot larger than what we want to compute.

**Example 15.1.** Take  $\varepsilon = 10^{-16}$ . Compute  $\begin{pmatrix} 1 & -1 & 0 \end{pmatrix} \cdot \begin{pmatrix} 1 + \varepsilon \\ 1 \\ 1 \end{pmatrix} = (1 + \varepsilon) - 1 + 0 = \varepsilon$ .

In this case we have a subtraction between two very close numbers.

$|\text{computer result}| \leq u \cdot (i + |1 \cdot (1 + \varepsilon)| + |-1 \cdot 1| + |0 \cdot 1|) = (2 + \varepsilon) \cdot u$ , which means that we have 10 correct digits.

The problem is in the fact that we have a very small result ( $\varepsilon = O(10^{-6})$ ) and a very small error (of the same order of the result), which implies a large relative error.

An attentive reader might have noticed that it's very long to make this computation. Therefore, since we computed it on easy examples, we have to observe that we can't afford it for SVD or other complicated methods.

Wilkinson trick (from the 60's) comes to help us, to simplify this computation for more complicated problems.

**Idea:** see the computer result as the exact output of an algorithm run on a slightly perturbed input.

$$\begin{aligned}
\tilde{y} &= \dots \\
&= ((a_1 b_1 (1 + \delta_1) + a_2 b_2 (1 + \delta_2)) (1 + \delta_4) + a_3 b_3 (1 + \delta_3)) (1 + \delta_5) \\
&= a_1 \tilde{b}_1 + a_2 \tilde{b}_2 + a_3 \tilde{b}_3
\end{aligned} \tag{15.3}$$

where

$$\begin{aligned}
\tilde{b}_1 &= b_1 (1 + \delta_1) (1 + \delta_4) (1 + \delta_5) = b_1 + 3u b_1 + o(u), \\
\tilde{b}_2 &= b_2 (1 + \delta_2) (1 + \delta_4) (1 + \delta_5) = b_2 + 3u b_2 + o(u), \\
\tilde{b}_3 &= b_3 (1 + \delta_3) (1 + \delta_5) = b_3 + 2u b_3 + o(u),
\end{aligned} \tag{15.4}$$

$$\tilde{a}_i = a_i \quad i = 1, 2, 3.$$

And the relative error is  $\frac{\|\tilde{b} - b\|}{\|b\|} \leq 3u + o(u)$ .  
Hence,

$$\frac{\|\tilde{y} - y\|}{\|y\|} \leq \kappa_{rel,b} \frac{\|\tilde{b} - b\|}{\|b\|} \leq k_{rel,b} \cdot 3u$$

This isn't bad, because it's within a factor 3 of the optimal error for a perfect algorithm.  
Computing the conditioning is much easier than making all the calculations of  $\delta$ s.

**Definition 15.2** (Backward stability of an algorithm). *An algorithm to compute  $\mathbf{y} = f(\mathbf{x})$  is called **backward stable** if the computed output  $\tilde{\mathbf{y}}$  can be written as  $\tilde{\mathbf{y}} = f(\tilde{\mathbf{x}})$ , where  $\tilde{\mathbf{x}} = \mathbf{x} + O(\mathbf{u} \|\mathbf{x}\|)$  (exact function, perturbed input).*

**Observation 15.1.** *In real-life usage, this  $O()$  notation often hides polynomial factors in the dimension  $n$ . Although this may look an illicit simplification, we observe that these factors are much more harmless than the error that we could make otherwise.*

**Theorem 15.1.** *Backward stable algorithms are as accurate as theoretically possible (given the condition number of a problem), up to some factor that depends only on the dimension (e.g.,  $n$ ,  $2n^2 + 18n$ , ...).*

*Proof.*

$$\frac{\|\tilde{\mathbf{y}} - \mathbf{y}\|}{\|\mathbf{y}\|} \leq \kappa_{rel}(f, \mathbf{x}) \frac{\|\tilde{\mathbf{x}} - \mathbf{x}\|}{\|\mathbf{x}\|} = \kappa_{rel}(f, \mathbf{x}) O(\mathbf{u}),$$

while the best attainable accuracy is  $\kappa_{rel}(f, \mathbf{x}) \mathbf{u}$ . □

We may ask ourselves if it's possible to perturb the input in order to get  $\tilde{y}$  for every possible algorithm and the answer is no. Let us see a counterexample, where  $f(a, b) = a^T b$  (vector-vector product in the order that produces a matrix).

We observe that in general it's not true that the computed approximation of  $f(a, b)$  has rank 1.

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \cdot \begin{bmatrix} y_1 & y_2 & y_3 \end{bmatrix} = \begin{bmatrix} x_1 y_1 & x_1 y_2 & x_1 y_3 \\ x_2 y_1 & x_2 y_2 & x_2 y_3 \\ x_3 y_1 & x_3 y_2 & x_3 y_3 \end{bmatrix} = M$$

While

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \odot \begin{bmatrix} y_1 & y_2 & y_3 \end{bmatrix} = \begin{bmatrix} x_1 \odot y_1 & x_1 \odot y_2 & x_1 \odot y_3 \\ x_2 \odot y_1 & x_2 \odot y_2 & x_2 \odot y_3 \\ x_3 \odot y_1 & x_3 \odot y_2 & x_3 \odot y_3 \end{bmatrix} = \widetilde{M}$$

And we are looking for  $\tilde{x}$  and  $\tilde{y}$  such that  $\widetilde{M} = \tilde{x}^T \cdot \tilde{y}$

**Example 15.2.** *Example (with exaggerated errors):*

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \cdot \begin{bmatrix} 4 & 5 & 6 \end{bmatrix} = \begin{bmatrix} 4 & 5 & 6 \\ 8 & 10 & 12 \\ 12 & 15 & 18 \end{bmatrix}$$

While

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \odot \begin{bmatrix} 4 & 5 & 6 \end{bmatrix} = \begin{bmatrix} 4.01 & 4.99 & 6.01 \\ 7.99 & 10.01 & 12.02 \\ 11.98 & 15.02 & 17.97 \end{bmatrix}$$

and this second matrix doesn't have rank 1.

## 15.2 Backward stability of QR factorization

### 💡 Do you recall?

A generic step of the computation of the QR factorization of a matrix has the following shape:

$$\begin{pmatrix} & & & & \\ & I & & & \\ & & H & u_k & \end{pmatrix} \begin{pmatrix} * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & * & * & * \end{pmatrix} = \begin{pmatrix} * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & * & * \end{pmatrix}$$

Each step of the QR factorization is backward stable.

Let us compute the backward stability of QR factorization:

With some tedious computations like the ones above, one can show that *one* step of the QR factorization is backward stable, i.e.,  $\tilde{R}_{k-1} = R_{k-1} + \Delta_{R_{k-1}}$ , where  $\|\Delta_{R_{k-1}}\| \leq O(u) \|R_{k-1}\|$  and this allows us to write the following

$$\frac{\|\tilde{R}_{k-1} - R_{k-1}\|}{\|R_{k-1}\|} \leq O(u)$$

At a generic step  $k$  we have that

$$R_k = Q_k \cdot R_{k-1}$$

$\Updownarrow$

$$\widetilde{R}_k = Q_k(R_{k-1} + \Delta_{R_{k-1}})$$

The idea is doing this procedure one step after the other, starting from step 1, where  $R_0 = A$ .

$$\widetilde{R}_1 = Q_1 \widetilde{R}_0 = Q_1 \cdot (R_0 + \Delta_{R_0}) = Q_1 \cdot (A + \Delta_{R_0})$$

$$\widetilde{R}_2 = Q_2 \widetilde{R}_1 = Q_2(R_1 + \Delta_{R_1}) = Q_2Q_1(A + \Delta_{R_0}) + Q_2\Delta_{R_1} = Q_2Q_1(A + \Delta_{R_0} + Q_1^T\Delta_{R_1})$$

We go on combining errors like this and we see that all the quantities  $(A + \Delta_{R_j})$  are perturbations of the original matrix  $A$ .

In the end we may observe that the final computed  $R_n$  is the exact result obtained from  $A$  plus  $n$  perturbations.

**Observation 15.2.** *We should notice that the norm of each perturbation is small with respect to the norm of  $A$ .*

*In formal terms,  $\|\Delta_{R_0}\| \leq u \cdot \|R_0\| = u \cdot \|A\|$  and  $\|Q_1^T\Delta_{R_1}\| \stackrel{(1)}{=} \|\Delta_{R_1}\| \leq u \cdot \|R_1\| \stackrel{(2)}{=} u \cdot \|A\|$ .*

*Where  $\stackrel{(1)}{=}$  and  $\stackrel{(2)}{=}$  hold because  $Q_i$  are orthogonal.*

### ★ Mantra

Orthogonal transformations are the key for stability.

#### 15.2.1 Stability of algorithms for least-squares problems

Let us see how various algorithms to solve LS problems (implemented in Matlab) behave in relation to backward stability.

##### Least squares problem via QR

STEP 1: Computing a thin QR (`qr(A, 0)`) → backward stable;

STEP 2: (`Q1'*b`) → backward stable;

STEP 3: (`R1 * c`) → backward stable;

##### Least squares problem via SVD

Scrivere meglio

STEP 1: Computing a SVD (`svd(A, 0)`) → backward stable;

STEP 2: (`U'*b`) → backward stable;

STEP 3: (`c ./ diag(S)`) → backward stable;

STEP 4: (`V*d`) → backward stable;

##### Least squares problem via Normal Equations

STEP 1: `C = A' * A;`

STEP 2:  $d = A' * b;$

STEP 3:  $x = C \cdot d;$

We can also prove that some algorithms aren't backward stable, like normal equations. The issue here is that the same input is needed in more than one operation, so it should satisfy more than one equation and this may lead to a solution set which is empty. Also, in the last equation we solve a linear system with matrix  $C$ , and this gives an error of the order of  $\kappa(A)^2$ , never of  $\kappa(A)$ :  $\kappa(C) = \kappa(A^T A) = \kappa(A)^2$ .

Let us take the example of two lectures ago: ??.

**Example 15.3.** Let  $A \in \mathcal{M}(4, 3, \mathbb{R})$  s.t.

$$A = \begin{pmatrix} 1 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 4 \\ 1 & 2 & 3 + 10^{-8} \end{pmatrix}$$

We may observe that  $A$  is at distance  $10^{-8}$  from a matrix without full column rank, hence  $\kappa \approx 10^8$ .

What is the condition number of solving this least squares problem? It's about  $10^8$ , since in that problem we generated  $b = A \cdot \begin{pmatrix} 3 & 4 & 5 \end{pmatrix}$ , so  $b$  lies in the image of  $A$  ( $\text{Im}(A)$ ). The geometric idea in Figure 15.2.




FIGURE 15.2:  $b$  lies (at least almost, because of numerical error) on the plane of  $\text{Im}(A)$ . In this case  $\kappa_{\text{rel}}(\text{solving LS}) \approx \kappa(A)$ .

A brief recap may be found in Table 1.

### 15.2.2 A posteriori checks

Let's assume that Matlab gives us a solution of a problem. We want to be able to check how good this result is.

**Example 15.4.** Suppose we have solved a linear system...

```
» A = randn(4, 4); b = randn(4, 1);
» x = A \ b;
» A*x - b
ans =
```

	Normal eqns	QR	SVD
$m \approx n$	$\frac{4}{3}n^3$	$\frac{4}{3}n^3$	$\approx 13n^3$
$m \gg n$	$mn^2$	$2mn^2$	$2mn^2$
Unstable when $cond \approx \kappa(A)$	Backward stable	Backward stable	stable; reveals info on sensitivity, allows regularization

TABLE 1: Brief recap of the complexities of the algorithms we studied for solving the least squares problem. The last row takes into account the stability.

$$\begin{array}{r} 0 \\ -1.3878e-17 \\ 0 \\ 2.2204e-16 \end{array}$$

**Definition 15.3** (Residual). Let  $A \in \mathcal{M}(m, \mathbb{R})$  and  $b \in \mathbb{R}^m$ , and  $x$  be the solution of  $Ax = b$ . For a given  $\tilde{x}$  we define **residual** the following  $\mathbf{r} = A\tilde{x} - b$ .

assume that  $\|\mathbf{r}\|$  is small; does this mean that  $\tilde{x}$  is close to the exact solution  $x$ ?

**Theorem 15.2.** Let  $A \in \mathbb{R}^{m \times m}, b \in \mathbb{R}^m$ , and  $x$  be the solution of  $Ax = b$ .

For a given  $\tilde{x}$ , the relative error of  $\tilde{x}$  is bounded by the condition of matrix  $A$  times the ratio between the norm of the residual and the norm of  $b$ .

$$\frac{\|\mathbf{r}\|}{\|b\|} \leq \kappa(A) \frac{\|\mathbf{r}\|}{\|b\|}.$$

This theorem tells us that  $\tilde{x}$  is “close to the solution” apart from a factor which is the conditioning of matrix  $A$ .

*Proof.* Follows from the perturbation results for linear systems. The idea is that  $\tilde{x}$  is the exact solution of the perturbed system

$$A\tilde{x} = b + \mathbf{r} = \tilde{b}$$

Where  $=^*$  follows from the definition of  $r$  and  $\frac{\|\tilde{b}-b\|}{\|b\|} = \frac{\|\mathbf{r}\|}{\|b\|}$ .  
A relative perturbation of size  $\frac{\mathbf{r}}{b}$  is amplified by  $\kappa(A)$ .  $\square$

It’s important to notice that also computing  $A \odot x \ominus b$  is an approximated operation. We choose to simplify things and ignore this error.

### 15.3 A posteriori check for Least Squares Problems

Can we make an analogous check for the Least Squares Problem?

Take a LSP  $\min \|Ax - b\|$ , with  $A$  tall thin, with full column rank.

The problem is that  $\|Ax - b\|$  isn't small at all, indeed it could be as large as  $b$ , as you can see from Figure 15.3.




FIGURE 15.3:  $A$  can be as large as  $b$  if  $b$  is perfectly orthogonal.

**Observation 15.3.** If you solve LSP via QR  $\|Ax - b\| = \left\| \begin{pmatrix} R_1 x - Q_1^T b \\ -Q_2^T b \end{pmatrix} \right\|$ . We said that the entries in the second block are fixed irrespective of  $x$ , but we could make the entries in the first block zero, by choosing  $x = R_1^{-1} Q^T b$ . This information let us infer something about the values of the vectors in Figure 15.3, in particular the minimum of the value that we can get is  $\|Q_2^T b\| = \|Ax - b\|$ .

With some algebra we may also check that  $\|Q_1^T b\| = \|Ax\|$ .

Since this is a minimum problem, we know that the gradient of the function is small near the optimum value:  $\min \|Ax - b\|^2 = \min x^T A^T Ax - 2b^T Ax + b^T b$ .

$$\nabla_{\tilde{x}} f = 2(A^T Ax - A^T b) \rightarrow 0$$

**Theorem 15.3.**  $\frac{\|\tilde{x} - x\|}{\|x\|} \leq (\kappa(A))^2 \cdot \frac{\|A^T Ax - A^T b\|}{\|A^T b\|}$ . Although we might have wanted to have the condition number of the problem, instead of the condition number of  $A$  and this could lead to underestimating the error.

Another idea could be using as error the first entry of the vector obtained via QR (namely  $R_1^T x - Q_1^T b$ ), by imposing  $R_1 x = Q_1^T b$

We may observe that this is a truly backward stable measure:

given  $r = \|R_1^T \tilde{x} - Q_1^T b\|$ , there exists  $\tilde{b}$  with  $\|\tilde{b} - b\| = \|r\|$  such that  $\tilde{x}$  is the exact solution of  $\min \|Ax - \tilde{b}\|$ .

We have proved the following

**Fact 15.4.**  $\frac{\|\tilde{x} - x\|}{\|x\|} \leq \kappa_{rel, LS} \cdot \frac{\|\tilde{b} - b\|}{\|b\|} = \kappa_{rel, LS} \cdot \frac{\|r\|}{\|b\|}$ .

**Theorem 15.5.** Let  $A = Q_1 R_1$  be a thin QR factorization. Let  $\mathbf{r}_1 = Q_1^T(A\tilde{\mathbf{x}} - \mathbf{b})$ . Then,  $\tilde{\mathbf{x}}$  is the exact solution of the LS problem

$$\min \|A\mathbf{x} - (\mathbf{b} + Q_1 \mathbf{r}_1)\|,$$

so the backward error of  $\tilde{\mathbf{x}}$  is  $\|Q_1 \mathbf{r}_1\| = \|\mathbf{r}_1\|$ .

*Proof.* Idea: replay the solution of a LS problem with QR factorization, and use  $Q_1^T T Q_1 = I$ . You will get in the first block  $R_1 x = Q_1^T b + \mathbf{r}_1$ , i.e.,  $Q_1^T(Ax - b) = \mathbf{r}_1$ , which is verified by  $\tilde{x}$ .  $\square$

# 16 16th of November 2018 — A. Frangioni

What happens when the Hessian isn't strictly positive definite? If there are some negative eigenvalues, I can deduce that there are some directions in which the function goes to  $-\infty$ . The model has no minimum, unless we restrict to a **compact set**.

In particular, we may decide to restrict to a part of the space where we can trust our model, which is called **trust region**.

Finding such a region is a NP-hard problem, if we don't restrict to a ball.

**Definition 16.1** (Karush-Kuhn-Tucker conditions). *Any optimal solution of the problem  $x^{i+1}$  must satisfy that  $\exists \lambda \geq 0$  s.t.*

KARUSH:  $[H^i + \lambda I]x^{i+1} = -\nabla f(x^i)$  [linear];

KHUN:  $H^i + \lambda I \succeq 0$  [semidefinite];

TUCKER:  $\lambda(r - \|x^{i+1}\|) = 0$  [nonlinear].

Where the last condition means that  $\lambda$  has to be 0, unless the solution we get is exactly on the border of the ball.

What's the difference between this approach and what we used to do before? We have two different cases:

- $\|x^{i+1}\| < r \implies \lambda = 0 \implies$  normal Newton step ( $\mathcal{T}$  has no effect);
- $\lambda > 0 (= \text{small radius } r) \implies$  like in line search with  $\varepsilon^i = \lambda$ .

The problem is computing these systems of equations taking less than  $O(n^3)$ .

## Key idea

We don't need to compute the Hessian, we can use the first order information to infer things on the second order matrix, although we don't really need to compute the Hessian.

## 16.1 Quasi-newton methods

At a step we have:  $m^i(x) = \nabla f(x^i)(x - x^i) + \frac{1}{2}(x - x^i)^T H^i(x - x^i)$ ,  $x^{i+1} = x^i + \alpha^i d^i$

At the next step we recompute the gradient in  $x^{i+1}$  and the matrix  $H^{i+1}$ :  $m^{i+1}(x) = \nabla f(x^{i+1})(x - x^{i+1}) + \frac{1}{2}(x - x^{i+1})^T H^{i+1}(x - x^{i+1})$

How should  $H^i$  be chosen? It should satisfy the following:

1. positive definite ( $H^{i+1} \succ 0$ );
2. we know the gradient in the previous point, we know the gradient in the current point, we construct  $H^{i+1}$  such that the model works:  $\nabla m^{i+1}(x^i) = \nabla f(x^i)$ ;

3.  $\|H^{i+1} - H^i\|$  is “small”.

This new model isn’t too different from the previous one, because of the third preerty. Or equivalently  $H^{i+1}(x^{i+1} - x^i) = \nabla f(x^{i+1}) - \nabla f(x^i)$ , which we call **secant equation** and we denote (S).

To ease notation we define  $s^i$  such that:  $s^i = x^{i+1} - x^i = \alpha^i d^i$  and  $y^i = \nabla f(x^{i+1}) - \nabla f(x^i)$ .  $s^i$  is chosen, while  $y^i$  is decided by the function.

In order to have a matrix  $H^i$  that satisfies the first two condition we could check that  $H^{i+1}s^i = y^i$ , because this implies  $s^i y^i = (s^i)^T H^{i+1} s^i$  and this implies 1. and 2., hence we obtain the **curvature condition**  $s^i y^i > 0$ .

**Theorem 16.1.** *Wolf condition implies  $s^i y^i > 0$ , using the notation we introduced:  $(W) \implies (C)$ .*

*Proof.*

$$\begin{aligned}\varphi'(\alpha^i) &= \nabla f(x^{i+1})d^i \geq m_3 \varphi'(0) = m_3 \nabla f(x^i)d^i \\ &\Downarrow \\ (\nabla f(x^{i+1}) - \nabla f(x^i))d^i &\geq (m_3 - 1)\varphi'(0) > 0\end{aligned}$$

□

**Observation 16.1.** *We may observe that this theorem implies that if we perform Armijo Wolf exact line search condition (C) can always be satisfied.*

### 16.1.1 Davidon-Fletcher-Powell

How can we choose a  $H^i$  that satisfies the three conditions enumerated above? Taking  $H^{i+1} = \operatorname{argmin}\{\|H - H^i\| : H \in (S) \text{ and } H \succeq 0\}$  is a good idea and for this minimum problem holds the following:

**Theorem 16.2** (Davidson-Feltcher-Powell). *The new matrix is obtained at each step constructing a rank two matrix, obtained from  $H^i$  as a rank to correction, as follows:  $H^{i+1} = (I - \rho^i y^i (s^i)^T) H^i (I - \rho^i s^i (y^i)^T) + \rho^i y^i (y^i)^T$*

Let us denote  $B^i = H^{i-1}$ . At any step we need to compute  $B^{i+1} = (H^{i+1})^{-1}$ , because we need to solve the system. We have some formulas that give us a way to compute  $(H^{i+1})^{-1}$  from  $H^{i-1}$ .

**Theorem 16.3** (Sherman-Morrison-Woodbury). *The inverse of a matrix of the form  $A + ab^T$  has the following shape:  $[A + ab^T]^{-1} = \frac{A^{-1} - A^{-1}ab^TA^{-1}}{1 - b^TA^{-1}a}$ .*

**Observation 16.2.** *From Theorem 16.3 we can conclude that  $B^{i+1} = \frac{B^i + \rho^i s^i (s^i)^T - B^i y^i (y^i)^T B^i}{(y^i)^T B^i y^i}$ .*

It’s important to notice that this operation has a cost of  $O(n^2)$ .

We can do better, in terms of computational complexity.

### 16.1.2 Broyden-Fletcher-Goldfarb-Shanno

We can use directly  $B^i$ , the inverse of  $H^i$ . Write (S) for  $B^{i+1}$ :  $s^i = B^{i+1}y^i \implies B^{i+1} = \operatorname{argmin}_{\{ \|B - B^i\| : \dots \}}$

$$B^{i+1} = B^i + \rho^i [(1 + \rho^i(y^i)^T B^i y^i) s^i (s^i)^T - (B^i y^i (s^i)^T + s^i (y^i)^T B^i)]$$

This formula proves to be more stable than the other one.

This method takes  $O(n^2)$ .

The two  $B^i$ 's, obtained from DFP and BFGS, are different although both sensible, but we can use a convex combination of the two.

**Observation 16.3.** *How can we choose  $H^1$ ? The value we choose will make a difference in the results, at least for the first steps.*

Let us see a couple of choices for  $B^1$ :

- Scalar multiples of identity, but how to choose the scalar?
- Compute the gradient in every direction and approximate  $H$ . This will cost  $O(n^3)$ , but it should be done only once.

Let us compute the space needed to store the  $B^i$ 's: order of  $n^2$  is still a lot. What happens if we restrict to working with information of the last  $k$  operations?

### 16.1.3 Poorman's approach

At each step we only consider  $B^{i-k}$  and  $k$  rank one operations. This operations cost  $n$  each, and we have  $k$  lines. The problem is that  $B^{i-k}$  takes  $O(n^2)$  space. We can optimize if we choose  $B^{i-k}$  to be simpler, say a multiple of the identity, or finite difference of the gradient. Then the space complexity is  $O(kn)$ .

I need to tune the algorithm to find the right  $k$  which gives me enough precision and also keeps the computational cost low.

#### Final observation of quasi Newton methods

We may notice that this variation of Newton method doesn't get trapped in local minima, as Newton method did. In the end, the fact that quasi Newton isn't that precise at the beginning may be a good feature.

## 16.2 Conjugate gradient method

### 💡 Do you recall?

In the gradient method, the angle between two consecutive directions is exactly  $90^\circ$ , as can be seen in Figure 16.1.




FIGURE 16.1: Geometric idea on how the new direction is chosen.

We would like to take into account not only the subspace spanned by  $d^{i+1}$  but we would like to optimize over larger and larger subspaces (spanned by  $d^i$  and  $d^{i+1}$ ).

**Definition 16.2** (Q-conjugate). *Let  $v$  and  $w$  be vectors in  $\mathbb{R}$ . We say that  $v$  ad  $w$  are Q-conjugate if  $(v)^T Q w = 0$ .*

We would like pick a direction to be  $Q$ -conjugate with all the previous iterations. The point is that we can't take into account all the previous directions, but we will see that we only need the previous direction to obtain all the information we need.

---

ALGORITHM 16.1 Pseudocode for conjugate gradient method for quadratic functions.

---

```

1: procedure CGQ( $Q, q, x, \varepsilon$ )
2:    $d^- \leftarrow 0;$ 
3:   while ( $\|\nabla f(x)\| > \varepsilon$ ) do
4:     if ( $d^- = 0$ ) then
5:        $d \leftarrow -\nabla f(x);$ 
6:     else
7:        $\beta = (\nabla f(x)^T Q d^-) / ((d^-)^T Q d^-);$ 
8:        $d \leftarrow -\nabla f(x) + \beta d^-;$ 
9:     end if
10:     $\alpha \leftarrow (\nabla f(x)^T d) / (d^T Q d);$ 
11:     $x \leftarrow x + \alpha d;$ 
12:     $d^- \leftarrow d;$ 
13:   end while
14: end procedure

```

---

The number of iterations needed to converge is proportional to the clusterization of the eigenvalues of the matrix  $Q$ .

The algorithm that was presented is for quadratic functions, but the same algorithm works for non quadratic function as well, as long as we change the formula for  $\beta$ .

The pseudocode of Algorithm 16.2 is referred to Fletcher-Reeves definition of  $\beta^i$   $\beta^i = \|\nabla f(x^i)\| / \|\nabla f(x^{i-1})\|^2$ .

This algorithm converges in at most  $n$  iterations.

---

ALGORITHM 16.2 Pseudocode for conjugate gradient method for arbitrary functions

---

```

1: procedure CGA( $Q, q, x, \varepsilon$ )
2:    $\nabla f^- = 0;$ 
3:   while ( $\|\nabla f(x)\| > \varepsilon$ ) do
4:     if ( $\nabla f^- = 0$ ) then
5:        $d \leftarrow -\nabla f(x);$ 
6:     else
7:        $\beta = \|\nabla f(x^i)\|^2 / \|\nabla f^-\|^2;$ 
8:        $d \leftarrow -\nabla f(x) + \beta d^-;$ 
9:     end if
10:     $\alpha \leftarrow \text{AWLS}(f(x + \alpha d));$ 
11:     $x \leftarrow x + \alpha d;$ 
12:     $d^- \leftarrow d;$ 
13:     $\nabla f^- \leftarrow \nabla f(x);$ 
14:   end while
15: end procedure

```

---

We have three different formulas for  $\beta_i$ , which coincide in the quadratic case.

1. Polak-Ribière:  $\beta^i = \frac{[\nabla f(x^i)^T (\nabla f(x^i) - \nabla f(x^{i-1}))]}{\|\nabla f(x^{i-1})\|^2}$
2. Hestenes-Stiefel:  $\beta^i = \frac{\nabla f(x^i)^T (\nabla f(x^i) - \nabla f(x^{i-1}))}{(\nabla f(x^i) - \nabla f(x^{i-1}))^T d^{i-1}}$
3. Dai-Yuan:  $\beta^i = \frac{\|\nabla f(x^i)\|^2}{(\nabla f(x^i) - \nabla f(x^{i-1}))^T d^{i-1}}$

Some of these algorithms require some hypothesis on the function in order for the conjugate method to converge.

1. Fletcher-Reeves requires  $m_1 < m_2 < \frac{1}{2}$  for (A)  $\cap$  (W') to work;
2. (A)  $\cap$  (W')  $\not\Rightarrow d^i$  of Polak-Ribière is of descent, unless  $\beta_{PR}^i = \max\{\beta^i, 0\}$ .

Sometimes it's important to restart from scratches if the algorithm isn't converging, because many bad choices may lead to a bad result.

The idea of taking the gradient and modify it instead of multiplying by a factor, adding the previous direction.

It's possible to design hybrids between quasi-Newton and conjugate method.

## 17 21st of November 2018 — F. Poloni

In this lecture we address the problem of solving linear systems exactly.

Someone could observe that this subject has already been studied in linear algebra course, but we are interested in computing fast the solution to this problem when the dimensions are a lot and the matrix  $A$  is sparse.

Since the complexity of Gauss method is cubic, this algorithm is unfeasible for large inputs.

Let us see some real life examples, where the matrices are large and sparse.

**LOCAL FUNCTION ON GRAPHS:** A **local function on graphs** is a function that depend on few nearby vertices. This kind of functions lead to a sparse adjacency matrix  $A$ , as can be observed in Figure 17.1;




FIGURE 17.1: A local function on graph.

**IMAGES:** Take an  $m \times m$  image and blur it (each pixel is obtained as the average of its neighbours).  $T : \mathcal{M}(m, \mathbb{R}) \rightarrow \mathcal{M}(m, \mathbb{R})$  such that  $T(A)_{ij} = \frac{1}{9} \cdot (A_{i-1j} + A_{i-1j-1} + A_{i-1j+1} + A_{ij} + A_{ij-1} + A_{ij+1} + A_{i+1j} + A_{i+1j-1} + A_{i+1j+1})$ .  $T$  may be written as a matrix that maps all the  $m$  images to a set of  $m$  blurred images and has the following shape  $T \in \mathcal{M}(m^2, \mathbb{R})$  such that the  $(i, j)$ -th row of  $T$  has exactly 9 entries with value  $\frac{1}{9}$  and all the others are 0. The non zero entries correspond to  $A_{i-1j}, \dots, A_{i+1j+1}$ ;

**KKT SYSTEMS** constrained optimization;

**ENGINEERING PROBLEM:** To check stability of a bridge, it gets splitted into small cells. It can be proven tha the stress on each of these cells corresponds to the force applied by the neighbours. In the end, this local phenomenon may be represented by a sparse matrix.




FIGURE 17.2: Graphic idea of a bridge partitioned into small blocks

## 17.1 Gaussian elimination and LU factorization

Gaussian elimination can be seen as a factorization:  $A = LU$ . The intuition is to proceed iteratively, multiplying each time for a new matrix, just like QR factorization.

Since the idea of Gauss elimination is to add multiples of row 1 to all the rows from 2 to  $n$  to kill off  $A_{2:end,1}$  we have that:

STEP 1:

$$\begin{pmatrix} 1 & & & & \\ * & 1 & & & \\ * & & 1 & & \\ * & & & 1 & \\ * & & & & 1 \end{pmatrix} \begin{pmatrix} \textcircled{\ast} & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \end{pmatrix} = \begin{pmatrix} \textcircled{\ast} & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \end{pmatrix},$$

Where the  $\textcircled{\ast}$  is called **pivot**.

$$L_1 A = A_1.$$

$$(L_1)_{k1} = -\frac{A_{k1}}{A_{11}}, \quad k = 2, 3, \dots, m.$$

STEP 2: we multiply for a matrix that has an “identity frame” and inside does the same  $L_1$  was doing before.

$$\begin{pmatrix} 1 & & & & \\ & 1 & & & \\ * & 1 & & & \\ * & & 1 & & \\ * & & & 1 & \end{pmatrix} \begin{pmatrix} * & \textcircled{\ast} & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \end{pmatrix} = \begin{pmatrix} * & \textcircled{\ast} & * & * & * \\ 0 & * & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & * & * & * \end{pmatrix}$$

$$L_2 A_1 = A_2,$$

$$(L_2)_{k2} = \frac{(A_1)_{k2}}{(A_1)_{22}}, \quad k = 3, \dots, m.$$

STEP 3: we go on and

$$\begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & * & 1 & & \\ & * & & 1 & \end{pmatrix} \begin{pmatrix} * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & 0 & \textcircled{\times} & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & * & * & * \end{pmatrix} = \begin{pmatrix} * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & 0 & \textcircled{\times} & * & * \\ 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & * & * \end{pmatrix}$$

$$L_3 A_2 = A_3,$$

$$(L_3)_{k3} = \frac{(A_2)_{k3}}{(A_2)_{33}}, \quad k = 4, \dots, m.$$

STEP 4: one more operation

$$\begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & * & 1 & & \end{pmatrix} \begin{pmatrix} * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & 0 & \textcircled{\times} & * \\ 0 & 0 & 0 & * & * \end{pmatrix} = \begin{pmatrix} * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & 0 & \textcircled{\times} & * \\ 0 & 0 & 0 & 0 & * \end{pmatrix}$$

$$L_4 A_3 = A_4,$$

$$(L_4)_{k4} = \frac{(A_3)_{k4}}{(A_3)_{44}}, \quad k = 5, \dots, m.$$

In the generic case we have  $L_{m-1} L_{m-2} \dots L_1 A = U$ , where  $U$  is upper triangular, or  $A = \underbrace{L_1^{-1} L_2^{-1} \dots L_{m-1}^{-1}}_{=L} U$ , with  $U$  upper triangular and  $L$  lower triangular.

**Theorem 17.1.** Let  $A \in \mathcal{M}(m, \mathbb{R})$  such that we do not encounter zero pivots in the algorithm.  $A$  admits a factorization  $A = LU$ , where  $L$  is lower triangular with ones on its diagonal, and  $U$  is upper triangular.

**Observation 17.1** (Stroke of luck). The product of the  $L_i^{-1}$ 's (denoted  $L$ ) can be computed for free, since the following holds:

$$\begin{bmatrix} 1 & & & & \\ -a_2 & 1 & & & \\ -a_3 & & 1 & & \\ -a_4 & & & 1 & \\ -a_5 & & & & 1 \end{bmatrix}^{-1} \cdot \begin{bmatrix} 1 & & & & \\ & 1 & & & \\ -b_3 & 1 & & & \\ -b_4 & & 1 & & \\ -b_5 & & & 1 & \end{bmatrix}^{-1} \cdot \begin{bmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & -c_4 & 1 & \\ & & -c_5 & & 1 \end{bmatrix}^{-1} \cdot \begin{bmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & & & & \\ a_2 & 1 & & & \\ a_3 & b_3 & 1 & & \\ a_4 & b_4 & c_4 & 1 & \\ a_5 & b_5 & c_5 & d_5 & 1 \end{bmatrix}$$

---

ALGORITHM 17.1 LU factorization, Matlab implementation.

---

```
1 function [L, U] = lu_factorization(A)
2 m = size(A, 1);
3 L = eye(m);
4 U = A;
5 for k = 1 : m - 1
6     % compute "multipliers"
7     L(k+1:end, k) = U(k+1:end, k) / U(k, k);
8     % update U
9     U(k+1:end, k) = 0;
10    U(k+1:end, k+1:end) = U(k+1:end, k+1:end) ...
11        - L(k+1:end, k) * U(k, k+1:end);
12 end
```

---

The idea behind the implementation of Algorithm 17.1 is shown in Figure 17.3.




FIGURE 17.3: Assuming that we are at step  $k$ , we have that the  $h$ -th multiplier is expressed as  $\frac{A_{hk}}{A_{kk}}$  and this multiplier goes to the right position in  $L$ .

**Observation 17.2.** *The computational complexity of this algorithm is concentrated at lines 10, 11 and the cost of thi operation is  $O((m - 1)^2 + (m - 2)^2 + \dots + 2^2 + 1)$ . So we have that for a dense matrix the computational complexity is  $\frac{2}{3}m^3 + O(m^2)$ , in other words half as much as QR factorization.*



## Something on Matlab ...

**Implementation of \ in Matlab:** We consider important to remark how the operator  $\backslash$  is implemented in Matlab. It works using the LU factorization, making some checks and changing the operations as follows:

- upper/lower triangular systems: back-substitution ( $O(n^2)$ );
- non-triangular linear systems: LU (then throw away the factors);
- symmetric and/or sparse systems: uses appropriate LU variants (will see in the following).

Why isn't there any check on the orthogonality of the matrix? Because in that case the inverse is equal to the transpose and the computational complexity would decrease. Well, in order to find out that a matrix is orthogonal we first need to compute its transpose, which is too costly.

**Obs:** LU isn't stable, as shown in Section 17.1.1, so Matlab uses a slightly different implementation.

missing

### 17.1.1 Stability of LU

A downside of this approach is that it's not numerically stable. The intuition is that the condition is bad whenever the matrix  $A$  has a very small pivot.

Let us see an example:

#### Example 17.1.

$$A = \begin{bmatrix} 10^{-30} & 1 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 10^{30} & 1 \end{bmatrix} \begin{bmatrix} 10^{-30} & 1 \\ 0 & 1 - 10^{30} \end{bmatrix}.$$

In this case the LU factorization produces  $L, U$  with norm much larger than  $\|A\|$ .

Luckily, it's easy to circumvent this issue multiplying  $L_i$ s by some permutation matrices (which swap rows in order to keep "large" pivots), as follows

$$L_{m-1}P_{m-1} \dots L_2P_2L_1P_1A = U.$$

**Observation 17.3.** Thanks to another "stroke of luck" we can reorder those factors:

$$L_{m-1}P_{m-1} \dots L_2P_2L_1P_1 = \hat{L}_{m-1}\hat{L}_{m-2} \dots \hat{L}_1P_{m-1}P_{m-2} \dots P_1$$

where  $\hat{L}_i$  have the same structure as the  $L_i$ .

We can now introduce the following

**Theorem 17.2.** Let  $A \in \mathcal{M}(m, \mathbb{R})$ .  $A$  admits a factorization  $A = PLU$ , where  $P$  is a permutation matrix,  $L$  is lower triangular with ones on its diagonal, and  $U$  is upper triangular.

What about the stability of this improvement to LU factorization, called **LU with partial pivoting**?

It's not stable at all, in the worst case it may happen that  $\|U\|/\|A\|$  may grow as  $\approx 2^m$ , although on average it never happens.

### 17.1.2 Gaussian elimination on sparse matrices

Given a sparse matrix

$$A = \begin{pmatrix} * & * & & * & * & * \\ & * & * & * & * & * \\ * & * & * & * & * & * \\ * & & * & & * & * \\ & * & & * & * & * \\ * & & * & * & * & * \\ * & * & & * & & * \\ * & * & * & * & & * \\ * & * & * & * & & * \end{pmatrix}$$

Gaussian elimination causes some fill-in, due to the sum of a multiple of the first row, which has non zero entries in different positions:

$$\begin{pmatrix} * & * & & * & * & * \\ & * & * & * & * & * \\ * & * & * & * & * & * \\ * & & * & & * & * \\ * & & & * & * & * \\ * & * & * & * & * & * \\ * & * & & * & * & * \\ * & * & * & * & * & * \\ * & * & * & * & * & * \end{pmatrix} \rightarrow \begin{pmatrix} * & * & & * & * & * \\ & * & * & * & * & * \\ 0 & * & * & * & * & * \\ 0 & * & * & * & * & * \\ * & & * & * & * & * \\ 0 & * & * & * & * & * \\ 0 & * & * & * & * & * \\ 0 & * & * & * & * & * \\ 0 & * & * & * & * & * \end{pmatrix}$$

We may observe that the computational complexity of sparse LU is linear in the number of non zero entries of the final matrix, obtained by the algorithm, which is possibly much larger than the number of non zeros in  $A$ .

How to circumvent this problem? At each step we may use as pivot row the most sparse one. This computation may be done in a more sofisticate way, considering the “relative” position of non zeros between couples of rows.

Because of this trade-off the choice is made in relation to the needs of the implementation. We won't study any algorithm that deals with sparse matrices, since they are very complicated and make use of heuristics.

There are some lucky cases in which the fill-in is almost none, for example a matrix that only has 5 diagonals which entries are different from 0 (called **tridiagonal**). In this particular case  $L$  is tridiagonal and lower triangular and  $U$  is tridiagonal and upper triangular, as shown below.

$$A = \begin{pmatrix} * & * & * & 0 & 0 & 0 & \cdots & 0 \\ * & * & * & * & 0 & 0 & \cdots & 0 \\ * & * & * & * & * & 0 & \cdots & 0 \\ 0 & * & * & * & * & * & & \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & \cdots & * & * & * & * \\ 0 & 0 & 0 & \cdots & * & * & * & * \\ 0 & 0 & 0 & \cdots & 0 & * & * & * \end{pmatrix}$$

$$L = \begin{pmatrix} * & & & & & \\ * & * & & & & \\ * & * & * & & & \\ & \ddots & \ddots & \ddots & & \\ & & * & * & * & \\ & & & * & * & * \\ & & & & * & * \end{pmatrix} U = \begin{pmatrix} * & * & * & & & \\ * & * & * & & & \\ * & * & * & & & \\ & \ddots & \ddots & \ddots & & \\ & & * & * & * & \\ & & & * & * & * \\ & & & & * & * \end{pmatrix}$$

**Observation 17.4.** We should remark that if we are interested into parallel computing we need to pay attention to the blocking, because we go from vector-vector operation to matrix-matrix operation and some of these operations may be performed efficiently via vectorization. Parallel implementations are available by means of parallel libraries for Matlab.

# 18 22nd of November 2018 — A. Frangioni

## 18.1 Deflected gradient methods

The idea behind this family of algorithms, is to determine the next position  $x^{i+1}$  using the gradient and summing to it something else that gives us more information.

This kind of algorithms work also in cases in which the gradient isn't continuous.

This methods use the information about the previous iterations without exploiting properties about the second order derivative.

### 18.1.1 Heavy ball gradient method

The intuition behind this algorithm may be expressed through the following metaphor: an object is moving in the space and it's subject to a force. We can observe that the heavier the object, the stronger should be the force imposed in order to make it describe a certain trajectory.

In this interpretation, we may define the  $(i + 1)$ -th iteration as

$$x^{i+1} \leftarrow x^i - \alpha^i \nabla f(x^i) + \beta^i (x^i - x^{i-1}),$$

where  $\beta^i$  is called **momentum**,  $x^i$  **heavy** and  $\nabla f(x^i)$  **force**.

This isn't a descent algorithm: we are not choosing a direction and doing line search. We have no guarantee that the value of the function after one iteration will be smaller than the previous one.

First thing to do is to choose  $\alpha^i$  and  $\beta^i$  properly.

We can prove for some cases that these methods are better than gradient method, although they aren't as good as Newton or quasi Newton. Their strength resides in their simplicity though.

Notice that if the smaller eigenvalue isn't zero (i.e. quadratic case) we have a close formula to choose  $\alpha$  and  $\beta$  independently from the iteration:

$$\alpha = \frac{4}{(\sqrt{\lambda^1} + \sqrt{\lambda^n})^2}, \quad \beta = \max \left\{ \left| 1 - \sqrt{\alpha \lambda^n} \right|, \left| 1 - \sqrt{\alpha \lambda^1} \right| \right\}^2$$

We may observe that the step we take is something that goes like  $\frac{1}{L}$ , where  $L$  is the Lipschitz constant, since  $\lambda_n$  is very small. With these choices the rate is the following. We observe that in the gradient the rate is the same, although there aren't the square roots

$$\|x^{i+1} - x_*\| \leq \left( \frac{\sqrt{\lambda^1} - \sqrt{\lambda^n}}{\sqrt{\lambda^1} + \sqrt{\lambda^n}} \right) \cdot \|x^i - x_*\|$$

An alternative idea could be choosing  $\beta^i$  and finding  $\alpha^i$  using line search. A possible issue is that we don't know if we are moving along a descending direction, but in this method it is perfectly acceptable not to make any movement at a single step (notice that in gradient method if one step has size 0 then we will not move anymore).

$\beta^i$  is seen as an hyperparameter, hence its value is tuned rerunning the algorithm several times.




FIGURE 18.1: On the left side Newton method and on the right side the heavy ball method.

The plot of the convergence of the heavy ball method gives a graphical idea of the fact that the direction isn't orthogonal to the one at the previous iteration, since we have the gradient plus some quantity.

In particular, the bigger  $\beta^i$  the “less orthogonal” the steps are. This feature allows the algorithm to have good performances on elongated functions.

### 18.1.2 Accelerated gradient

This method works only on convex functions, it has some similarities with heavy ball, but it's slightly different.

---

ALGORITHM 18.1 Pseudocode for accelerated gradient method.

---

```

1: procedure ACCG( $f, \nabla f, x, \varepsilon$ )
2:    $x_- \leftarrow x;$ 
3:    $\gamma \leftarrow 1;$ 
4:   repeat
5:      $\gamma_- \leftarrow \gamma;$ 
6:      $\gamma \leftarrow (\sqrt{4\gamma^2 + \gamma^4} - \gamma^2)/2;$ 
7:      $\beta \leftarrow \gamma(1/\gamma_- - 1);$ 
8:      $y \leftarrow x + \beta(x - x_-);$ 
9:      $g \leftarrow \nabla f(y);$ 
10:     $x_- \leftarrow x;$ 
11:     $x \leftarrow y - (1/L)g;$ 
12:   until ( $\|g\| > \varepsilon$ )
13: end procedure
```

---

The rational behind this algorithm is the following: When we are in a certain point at a certain iteration, we go on a little bit  $\beta^i$  and we end up in a point  $y$ . The gradient is computed in that point and used to choose the next point.




FIGURE 18.2: We build a linear model, which is a lowerbound for the function, then we build a quadratic model, which is above our function.

If we choose  $\gamma$  optimally then the quadratic approximation is very close to the function. We want to find the value for  $\gamma_s$  that gives best results in the worst case.

We can prove that the error  $\|f(x^i) - f_*\| \leq \sigma^i(f(x^i) - f_*) \searrow 0$  is multiplied by this factor  $\delta_i$ , that goes like the inverse of  $i^2$ .

Notice that if we choose  $\alpha$  small, it will always be small.

The convergence is sublinear.

**Theorem 18.1** (Optimality of accelerated gradient). *If the function isn't strongly convex no algorithm has better convergence than  $|f(x^i) - f_*| = 3L \frac{\|x^1 - x_*\|^2}{32(i+1)^2}$ .*

**Observation 18.1.** *This theorem tells us that this algorithm never gets worse than  $|f(x^i) - f_*| = 3L \frac{\|x^1 - x_*\|^2}{32(i+1)^2}$ , but this doesn't imply that this method is fast on average. The state of the art provides a lot of different formulas for  $\beta$ , which of the ones leads to some theoretical results.*

From now on we will move towards a different family of functions, that aren't even differentiable, hence we can't compute the gradient.

## 18.2 Incremental gradient methods

This method has good performances in real world machine learning cases, where the function is differentiable but we do not want to compute the gradient.

Let  $I = \{1, \dots, m\}$  be the set of observations,  $X = [X^i \subset \mathcal{X}]_{i \in I}$  the set of inputs and  $y = [y^i]_{i \in I}$  the set of outputs. Our goal is to explain  $y$  from  $X$ .

Since these vectors are uniformly distributed over the space (at least this is our hypothesis) when they get summed we expect some of them to cancel out.

The idea is to rewrite the problem as learning a linear function in the feature space, formally  $\min \{ \sum_{i \in I} l(y^i, \langle \Phi(X^i), w \rangle) : w \in \mathbb{R}^n \}$ , where  $l(\cdot, \cdot)$  is called **loss function**.

We are now interested in computing the gradient, which is not hard to compute since the function is linear:  $\nabla f(w) = \sum_{i \in I} \nabla f^i(w) = \sum_{i \in I} -A^i(y^i - A^i w)$ .

The issue here is that computing the gradient, although it's the gradient of a very simple function, takes too long to be computed (since there are too many vectors in machine learning datasets).

To overcome this problem we choose to restrict to only a subset of observations. How can we choose such set? Randomly, of course.

At this point the algorithm isn't deterministic anymore, but it's completely **stochastic**.

The intuition behind this algorithm is to take only one observation, compute what is needed on this observation and make a small step.

An online application may be a sensor that produces hundreds of outputs per second and it's not possible to store each of them. They should be used to infer some information and then thrown away.

We study the converge of this kind of algorithms from a stochastic point of view.

In machine learning we always need some regularization, because the tuning of hyperparameters clearly takes into account only the error in the samples that have been seen. Let us regularize the model as follows

$$\min \left\{ \sum_{i \in I} l(y^i, \langle \Phi(X^i), w \rangle) + \mu \Omega(w) : w \in \mathbb{R}^n \right\}$$

The usage of regularization may be useful, since we want to keep close to the minimum, but not reach it. It's enough to change slightly the problem and then solve it.

The regularization hyperparameter  $\Omega(w)$  may be chosen as follows:

1. Lasso regularizer (best known):  $\Omega(w) = \|w\|_1$ ;
2. In order to increase sparsity:  $\Omega(w) = \|w\|^2$ ;
3. Leading to feature selection: many  $w_j = 0$  as possible.

$\Omega$  function is not differentiable, so the function gets non differentiable, as an example look at Figure 18.3, which represents the plot of  $f(w_1, w_2) = (3w_1 + 2w_2 - 2)^2 + 10(|w_1| + |w_2|)$ .

### 18.3 Subgradient methods

These methods are thought for convex functions that are not differentiable.

Let us consider a kinky point: how can we choose between all the subgradients of that point? We assume to be able to compute some subgradients; since the function is convex we may recall




FIGURE 18.3: This function has a lot of kinky points.

**Property 18.2.** Let  $f$  be a convex function,  $\forall g \in \partial f(x), \forall y \in \mathbb{R}^n g(y) \equiv f(y) \geq f(x) + g(y - x)$ .

Let  $x^*$  be the optimum,  $\langle x^* - x^i, g \rangle$  is smaller than 0. This means that the angle between  $x$  and  $x^*$  is acute. If we knew the exact direction  $x - x^*$  the line search would land on  $x^*$ . For a pictorial representation see Figure 18.4




FIGURE 18.4: Since we know that  $\langle g, x^* \rangle$  is negative we get that the angle between  $g$  and  $x^*$  is larger than  $90^\circ$ . Moreover, we know where the optimum is not (red region), hence we restrict to the green region. At this point we will perform a line search on  $g$  direction. Since the function is not smooth, the line search may not succeed since the value of the function along the half line from  $x$  in  $g$  direction may remain the same, but there may be some points there that are closer to  $x^*$ .

The intuition behind this algorithm is to move in the direction of  $-g$ , but with a small  $\alpha^i$ , because if the step is too large we may end up in a point which is actually further from  $x^*$  than the previous step. In that point we may find a point where perform line search, because

that point isn't kinky. In this context we won't try to minimize  $\|f(x) - f(x^*)\|$ , but we will minimize  $\|x - x^*\|$ , because the function may zigzag near that point. It goes without saying that choosing a too small value for  $\alpha$  leads to a too slow convergence speed.

# 19 23rd of November 2018 — F. Poloni

## 19.1 Gaussian elimination on symmetric matrices



Do you recall?

In Gaussian elimination we had  $A$  and we multiplied it by  $L_1$  in order to get a big chunk of 0s in the first column

$$\begin{pmatrix} 1 & & & & & \\ * & 1 & & & & \\ * & & 1 & & & \\ * & & & 1 & & \\ * & & & & 1 & \\ * & & & & & 1 \end{pmatrix} \begin{pmatrix} \odot & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \end{pmatrix} = \begin{pmatrix} \odot & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \end{pmatrix},$$

Let us consider an upgrade of Gaussian elimination in the case of  $A \in S(m, \mathbb{R})$ .

Let us see what happens if we multiply  $L_1 A$  on the right by the transpose of  $L_1$ :

STEP 1:

$$\begin{pmatrix} 1 & & & & & \\ * & 1 & & & & \\ * & & 1 & & & \\ * & & & 1 & & \\ * & & & & 1 & \\ * & & & & & 1 \end{pmatrix} \begin{pmatrix} * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \\ * & * & * & * & * \end{pmatrix} \begin{pmatrix} 1 & * & * & * & * \\ 1 & 1 & & & \\ & 1 & 1 & & \\ & & 1 & 1 & \\ & & & 1 & \end{pmatrix} = \begin{pmatrix} * & 0 & 0 & 0 & 0 \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \end{pmatrix}$$

STEP 2:

$$\begin{pmatrix} 1 & & & & & \\ & 1 & & & & \\ * & 1 & & & & \\ * & & 1 & & & \\ * & & & 1 & & \\ * & & & & 1 & \end{pmatrix} \begin{pmatrix} * & 0 & 0 & 0 & 0 \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \end{pmatrix} \begin{pmatrix} 1 & & & & \\ 1 & 1 & * & * & * \\ & 1 & 1 & & \\ & & 1 & 1 & \\ & & & 1 & \end{pmatrix} = \begin{pmatrix} * & 0 & 0 & 0 & 0 \\ 0 & * & 0 & 0 & 0 \\ 0 & 0 & * & * & * \\ 0 & 0 & * & * & * \\ 0 & 0 & * & * & * \end{pmatrix}$$

STEP  $m$ :

$$L_{m-1} L_{m-2} \dots L_1 A L_1^T \dots L_{m-2}^T L_{m-1}^T = D,$$

where  $D$  is diagonal, or

$$A = L_1 L_2 \dots L_{m-1} D L_{m-1}^T \dots L_2^T L_1^T = LDL^T.$$

**Observation 19.1** (Stroke of luck). Notice that the stroke of luck of Observation 17.1 holds in this case too, hence we pay nothing to compute matrix  $L$ .

$$\begin{bmatrix} 1 & & & & \\ -a_2 & 1 & & & \\ -a_3 & & 1 & & \\ -a_4 & & & 1 & \\ -a_5 & & & & 1 \end{bmatrix}^{-1} \cdot \begin{bmatrix} 1 & & & & \\ & 1 & & & \\ -b_3 & 1 & & & \\ -b_4 & & 1 & & \\ -b_5 & & & 1 & \end{bmatrix}^{-1} \cdot \begin{bmatrix} 1 & 1 & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ -c_5 & & & & 1 \end{bmatrix}^{-1} \cdot \begin{bmatrix} 1 & 1 & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & & & & \\ a_2 & 1 & & & \\ a_3 & b_3 & 1 & & \\ a_4 & b_4 & c_4 & 1 & \\ a_5 & b_5 & c_5 & d_5 & 1 \end{bmatrix}$$

**Theorem 19.1** (Symmetric Gaussian elimination). Let  $A \in S(m, \mathbb{R})$  such that during Gaussian elimination we don't encounter any 0 pivot.  $A$  admits a factorization  $A = LDL^T$ , where  $L$  is lower triangular with ones on its diagonal, and  $D$  is diagonal.

A Matlab implementation of symmetric Gaussian elimination is shown in Algorithm 19.1.

---

ALGORITHM 19.1 Symmetric Gaussian factorization, Matlab implementation.

---

```

1  function [L, D] = ldl_factorization(A)
2      m = size(A, 1);
3      L = eye(m); D = zeros(m);
4      for k = 1:m-1
5          D(k, k) = A(k, k);
6          L(k+1:end, k) = A(k+1:end, k) / A(k, k);
7          A(k+1:end, k+1:end) = A(k+1:end, k+1:end) ...
8              - L(k+1:end, k) * A(k, k+1:end);
9      end
10     D(m, m) = A(m, m);

```

---

It is possible to make an optimization of this algorithm: since  $A$  is supposed to be symmetric, we only need to update the lower triangular part of  $A$ , since the rest is mirrored by symmetry, hence the computational complexity is half the one of Gaussian elimination.

This algorithm is not stable, exactly like the one on non symmetric matrices. Pivoting may be performed in order to augment stability. It comes without saying that the row swap should be done consistently on the columns to preserve symmetry.

Of course there are some matrices (like the ones with all 0s on the diagonal) that cannot be “pivoted”. There are some libraries that permute the rows and the columns in order to get better pivots. As an example, Matlab's  $[L, D, P] = \text{ldl}(A)$  produces matrices such that  $P^T AP = LDL^T$ , where  $D$  may have  $2 \times 2$  diagonal blocks.



Do you recall?

We recall the definition of **positive definite matrix**  $A \in M(m, \mathbb{R})$ : all its eigenvalues are strictly positive. In other words,  $A$  is positive definite if  $\forall z \neq 0 \in \mathbb{R}^m z^T A z > 0$ .

**Lemma 19.2.** In the context of positive definite matrices the following holds:

1. Let  $A$  be a symmetric matrix.  $A$  is positive definite if and only if  $MAM^T$  is so, for some invertible  $M \in M(m, \mathbb{R})$ . Formally,  $\forall A \in S(m, \mathbb{R})$  s.t.  $A \succ 0 \Leftrightarrow \exists M \in GL(m, \mathbb{R})$  s.t.  $MAM^T \succ 0$ ;

2. Let  $A$  a symmetric positive definite matrix such that  $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$ , then  $A_{11}$  and  $A_{22}$  are, too. Formally,  $\forall A \in S(m, \mathbb{R})$  s.t.  $A \succ 0$  and  $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \Rightarrow A_{11} \succ 0$  and  $A_{22} \succ 0$ .

*Proof.*

1.

$$\Rightarrow) A \in S(m, \mathbb{R}) \text{ and } A \succ 0 \Rightarrow MAM^T \in S(m, \mathbb{R}) \text{ and } MAM^T \succ 0.$$

Take  $z \in \mathbb{R}^m$ ,  $z \neq 0$   $z^T MAM^T z = y^T A y > 0$ , where we performed a variable change  $y = M^T z$ . Notice that  $y \neq 0$  because  $M$  is invertible (and  $\ker(M) = \{0\}$ ). The symmetry of the matrix  $MAM^T$  follows from  $(MAM^T)^T = M^{T^T} A^T M^T = MAM^T$ ;

$$\Leftarrow) MAM^T \in S(m, \mathbb{R}) \text{ and } MAM^T \succ 0 \Rightarrow A \in S(m, \mathbb{R}) \text{ and } A \succ 0.$$

This proof follows from the previous arrow, where the substitution is  $z = M^{-1}y$ .

2.  $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$  positive definite  $\Rightarrow A_{11}$  and  $A_{22}$  are positive definite too.

Since  $A$  is positive definite, its scalar product is greater than zero with all the vectors in  $\mathbb{R}^m$ .

$A_{11}$ ) Let us take  $\mathbf{z} = \begin{bmatrix} \mathbf{z}_1 \\ 0 \end{bmatrix}$ .

$$[\mathbf{z}_1^T \ 0] \cdot \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \cdot \begin{bmatrix} \mathbf{z}_1 \\ 0 \end{bmatrix} = \mathbf{z}_1^T A_{11} \mathbf{z}_1 > 0, \forall \mathbf{z}_1 \in \mathbb{R}^{\text{sizeof} A_{11}}$$

$A_{22}$ ) Let us take  $\mathbf{z} = \begin{bmatrix} 0 \\ \mathbf{z}_2 \end{bmatrix}$ .

$$[0 \ \mathbf{z}_2^T] \cdot \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \cdot \begin{bmatrix} 0 \\ \mathbf{z}_2 \end{bmatrix} = \mathbf{z}_2^T A_{22} \mathbf{z}_2 > 0, \forall \mathbf{z}_2 \in \mathbb{R}^{\text{sizeof} A_{22}}$$

□

**Corollary 19.3.** Let  $A \in M(n, \mathbb{R})$  such that  $A$  is positive definite. When computing the  $LDL^T$  factorization of  $A$ , at each step we have  $D_{kk} > 0$ , hence we need no pivoting technique.

*Proof.* From the first point of Lemma 19.2 we have that, since  $A \succ 0$ ,  $L_1 A L_1^T$  is positive

definite. Thanks to the second point of the same lemma we have  $L_1 A L_1^T = \begin{pmatrix} * & 0 & 0 & 0 & 0 \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \end{pmatrix}$

and so the first and the second diagonal blocks are positive definite ( $D_{11} > 0$  and  $D_{22} \succ 0$ ).

Notice that, since  $A$  is positive definite  $A_{11} \succ 0$ , but it's a scalar, hence  $A_{11} > 0$  and this implies no breakdown case. □

We may introduce another kind of factorization.

## 19.2 Cholesky factorization

The key idea is to write the diagonal matrix of the Gaussian elimination  $D$  as product of  $D^{1/2}$  times itself:

$$D = \begin{pmatrix} d_{11} & & & \\ & d_{22} & & \\ & & \ddots & \\ & & & d_{mm} \end{pmatrix} = \begin{pmatrix} \sqrt{d_{11}} & & & \\ & \sqrt{d_{22}} & & \\ & & \ddots & \\ & & & \sqrt{d_{mm}} \end{pmatrix} \cdot \begin{pmatrix} \sqrt{d_{11}} & & & \\ & \sqrt{d_{22}} & & \\ & & \ddots & \\ & & & \sqrt{d_{mm}} \end{pmatrix}$$

The Gaussian elimination may be rewritten as follows

$$A = LDL^T = LD^{1/2}(D^{1/2}L^T) = CC^T,$$

where  $D^{1/2} = \text{diag}(D_{11}^{1/2}, D_{22}^{1/2}, \dots, D_{mm}^{1/2})$ , and  $C$  is lower triangular (but not anymore with ones on the diagonal).

In Matlab the Cholesky factorization of a positive definite matrix is performed by the function `chol(A)`; and returns  $C^T$ .

**Observation 19.2.** *We will not discuss stability further, but Cholesky is always backward stable even without pivoting ( $\|C\| = \|A\|^{1/2}$ ).*

**Observation 19.3.** *In a sparse matrix, we can choose the (symmetric) permutation with the only goal of reducing fill-in. The same considerations about LU factorization hold in this case too.*

## 19.3 Krylov subspace methods

In this part we will discuss how to combine optimization methods to transform a linear system into a minimum problem.

**Example 19.1.** Let us assume we are interested in solving a linear system  $Ax = b$ .

STEP 1:  $x_0 = 0$ ,  $\nabla f(x_0) = -b$ ;

STEP 2:  $x_1 = \text{some multiple of } b \in \text{Span}(b)$

$$f(x_1) = Ax_1 - b$$

$$\nabla f(x_1) = \text{some multiple of } Ab + \text{some multiple of } b \in \text{span}(b);$$

STEP 3:  $x_2 = \text{mult. of } x_1 + \text{mul. of } \nabla f(x_1) + \text{mult. of } x_0 + \dots \in \text{span}(Ab_1, b)$ ;

STEP 4:  $x_3 = \text{mult. of } x_2 + \dots \nabla f(x_2) + \text{previous iterates} \dots$

$$Ax_2 - b = A \cdot (sAb + tb) - b = sA^2b + tAb - b \in \text{span}(b, Ab, A^2b), \text{ where } s \text{ and } t \text{ are scalars};$$

STEP 5:  $x_4 \in \text{span}(b, Ab, A^2b, A^3b)$ .

Notice that we can make some linear combinations of the vectors we have available and  $A(A(sb + tAb) + uAb + vb) + e(sb + tAb) + fAb + gb \in \text{span}(b, Ab, A^2b, A^3b)$ .

**Idea:** first compute the basis of  $\text{span}(b, Ab, A^2b, A^3b)$ , then look for the best solution inside this subspace.

**Observation 19.4.** *The cost of multiplying a sparse matrix  $A$  with a vector  $z$  is  $O(\text{nnz}(A))$ , where  $\text{nnz}(A)$  is the number of non zero entries of  $A$ .*

*Proof.* Let us assume the matrix  $A$  is stored as a vector, whose entries are  $(i, j, A_{ij})$ .

The matrix-vector product would then be

1. `x=zeros`
2. `for (i, j, Aij) such that Aij ≠ 0`
3.  $x_i = x_i + A_{ij} * z_j$
4. `end`

□

From now on we will consider to have a function `compute_product_with_A` that we use to compute matrix-vector products with  $A$ . In particular,  $x = \text{compute\_product\_with\_A}(z)$  computes  $x = Az$ , given  $z$ .

If we somehow have matrices not really sparse, but allow to compute a clever matrix-vector product this class of algorithms will give good results.

**Definition 19.1** (Krylov subspace). *Let  $A \in M(m, \mathbb{R})$  and let  $b \in \mathbb{R}^m$ . The **Krilev subspace** of index  $n$  is  $\mathbf{K}_n(\mathbf{A}, \mathbf{b}) = \text{span}(b, Ab, A^2b, \dots, A^{n-1}b)$ .*

*Equivalently,  $k \in K_n(A, b) \iff \exists \alpha_1, \dots, \alpha_{n-1} \in \mathbb{R}^m$  s.t.  $v = \alpha_0b + \alpha_1Ab + \alpha_2A^2b + \dots + \alpha_{n-1}A^{n-1}b$ .*

*Equivalently,  $(\alpha_0 + \alpha_1A + \alpha_2A^2 + \dots + \alpha_{n-1}A^{n-1})b = p(A)b$  for a polynomial  $p$  of degree such that  $\deg(p) \leq n - 1$ .*

**Observation 19.5** (Properties).

1.  $v, w \in K_n(A, b) \Rightarrow \alpha + \beta W \in K_n(A, b);$
2.  $v \in K_n(A, b) \Rightarrow Av \in K_n(A, b)$ . *Proof.* Let us take  $v = \alpha_0b + \dots + \alpha_{n-1}b$ , then  $Av = A(\alpha_0b + \dots + \alpha_{n-1}b) = \alpha_0Ab + \dots + \alpha_{n-1}A^n b;$
3.  $\dim(K_n(A, b)) \leq n$ . It is exactly  $n$  if  $b, Ab, A^2b, \dots, A^{n-1}b$  are linearly independent.
4. Let us assume  $\dim(K_n(A, b)) \leq n$ . In the second point, if  $A^{n-1}$  was really necessary  $\alpha_{n-1} \neq 0$  or  $v \in K_n(A, b)$ ,  $v \notin K_{n-1}(A, b)$ , equivalently then  $A^n b$  is really necessary to write  $Av$ , i.e.  $Av \in K_{n+1}(A, b)$  but  $Av \notin K_n(A, b)$ .
5. We may observe that  $\dim(K_1(A, b)) < \dim(K_2(A, b)) < \dots < \dim(K_{n_{\max}}(A, b)) = \dim(K_{n_{\max}+1}(A, b)) = \dots$

# 20 28th of November 2018 — A. Frangioni

## 20.1 Subgradient methods

We are still in the hypothesis of convex objective functions that are not differentiable in the whole domain.



Property 18.2 of last lecture: for  $f$  convex function,  $\forall g \in \partial f(x), \forall y \in \mathbb{R}^n g(y) \equiv f(y) \geq f(x) + \langle g, y - x \rangle$ . This is a characterization of the function with respect to the model.

Let us assume we know the minimum point  $x_*$ . We observe that the scalar product between the subgradient and the direction we should choose is negative. Formally,  $f(x_*) \geq f(x) + \langle g, x_* - x \rangle$ , hence  $\langle g, x_* - x \rangle \leq f(x_*) - f(x) \leq 0$ .

We want to bound the distance between the point at the “next step” and the optimum:

$$\begin{aligned} \|x^{i+1} - x_*\|^2 &\stackrel{(1)}{=} \|x^i - \alpha^i d^i - x_*\|^2 \\ &= \|x^i - x_*\|^2 + 2\alpha^i g^i \frac{(x_* - x^i)}{\|g^i\|} + (\alpha^i)^2 \\ &\stackrel{(2)}{\leq} \|x^i - x_*\|^2 - 2\alpha^i \frac{(f(x^i) - f(x_*))}{\|g^i\|} + (\alpha^i)^2 \end{aligned} \quad (20.1)$$

Where  $\stackrel{(1)}{=}$  follows from the definition of a normalized step:  $x^{i+1} = \frac{x^i - \alpha^i g^i}{\|g^i\|}$  and  $\stackrel{(2)}{\leq}$  follows from the inequality we stated above.

**Observation 20.1.** *The distance from the optimum is bounded by a square function in  $\alpha_i$  (the step size), where the linear part is negative and the quadratic part is positive.*

Hence, if the steps are short enough we get close to the optimum fast enough, because the linear part dominates the quadratic one.

Formally,  $-2 \frac{(f(x^i) - f(x_*))}{\|g^i\|} < 0 + \alpha^i \searrow$ , hence  $(\alpha^i)^2 \searrow 0_+$ .

An attentive reader may notice that from Equation (20.1) follows  $\|x^{i+1} - x_*\|^2 < \|x^i - x_*\|^2 + (\alpha_i)^2$  and  $\|x^{i+1} - x_*\|^2 \leq \|x^1 - x_*\|^2 + \sum_{k=1}^i (\alpha^k)^2$ .

The point here is that we cannot choose a too small  $\alpha$ , recall Armijo conditions.

If the series of the squares of step sizes does not diverge, then the sequence does not diverge as well, hence it converges somewhere, say  $\bar{x}$ .

The convergence of the series of the squares may be obtained using the following

**Definition 20.1** (Diminishing-Square Summable). *We term a series that diverges, while the series of the squares of the terms diverges, as **diminishing-square summable**.*

Formally,

$$(DSS) \sum_{i=1}^{\infty} \alpha^i = \infty \wedge \sum_{i=1}^{\infty} (\alpha^i)^2 < \infty.$$

Assumptions: function convex, definite in all its domain, hence the norm of the gradient will never become very large.

**Fact 20.1.** *We claim that the sequence of the stepsizes is a diminishing-square sequence.*

*Proof by contraddiction.* Let us assume  $f(x^i) - f_* \geq \varepsilon > 0$ ,  $\forall i$ . Then,

$$\|x^{i+1} - x_*\|^2 \leq \|x^i - x_*\|^2 - \delta \cdot \sum_{k=1}^i \alpha^k + \sum_{k=1}^i (\alpha^k)^2$$

The contraddiction is due to the fact that as  $i \rightarrow \infty$  the right-hand side goes to  $-\infty$ .  $\square$

This family of algorithms is clearly incredibly robust in theory, but in practice it does not work very well.

Let us make an experiment: let us suppose we know the minimum of the function  $f$ .

**Definition 20.2** (Polyak stepsize). *Let  $f$  be our convex objective function and let  $x_*$  be the optimum for  $f$ . We term **Polyak stepsize**:*

$$(PSS) \quad \alpha^i = \beta^i \frac{(f(x^i) - f(x_*))}{\|g^i\|}$$

where  $\beta \in (0, 2)$ .

If we pick a Polyak stepsize, then  $\|x^{i+1} - x_*\|^2 < \|x^i - x_*\|^2$ , so  $\{x^i\} \rightarrow x_*$ . The best value for  $\beta_i$  is 1. In this case, we obtain what follows by substituting  $\beta_i = 1$  in Equation (20.1)

$$\frac{(f(x^i) - f(x_*))^2}{\|g^i\|^2} \leq \|x^i - x_*\|^2 - \|x^{i+1} - x_*\|^2$$

The problem is that we don't know the optimum.

Let us assume that we know it and compute the efficiency:

Since we know that the sequence is bounded, we know that the objective function is globally Lipschitz (the norm of the gradient is bounded above).

The point is that the sequence  $\{x_1\}$  is not necessarily monotone, so we pick the so-called record value of best value ( $\underline{f}^i = \min\{f(x^h) : h = 1, \dots, i\}$ )

$$\frac{(\underline{f}^i - f(x_*))^2}{L^2} \leq \frac{(f^i - f(x_*))^2}{\|g_i\|^2} \leq \|x^i - x_*\|^2 - \|x^{i+1} - x_*\|^2$$

Summing for  $i = 1, \dots, k$  we obtain a telescopic series:

$$\|x^1 - x_*\| - \|x^2 - x_*\| + \|x^3 - x_*\| - \|x^4 - x_*\| + \dots + \|x^k - x_*\| - \|x^{k+1} - x_*\|$$

Hence resulting in

$$k \cdot \frac{(\underline{f}^k - f(x_*))^2}{L^2} \leq \|x^1 - x_*\|^2 - \|x^{k+1} - x_*\|^2 \leq \|x^1 - x_*\|^2 = R$$

which is equivalent to  $(\underline{f}^k - f(x^*))^2 \leq \frac{R^2 L^2}{k}$ , which is again equivalent to  $\underline{f}^k - f(x^*) \leq \sqrt{\frac{RL}{k}}$ , where  $L$  is the Lipschitz constant.

The issue here is that the convergence is sublinear:  $k \geq \frac{1}{\varepsilon^2}$ .

**Theorem 20.2.** *Take an algorithm that uses only the subgradient. It's possible to construct a function that makes the algorithm converge with sublinear speed. Hence, we cannot do better.*

It comes without saying that although this algorithm is not very good it is the “less bad” it can be.

There are some lucky cases in which we do know the optimal value, but this is not the case usually.

### 20.1.1 Target level stepsize

Let us assume  $f(x_*)$  is unknown. The only information available is that this value is below any value in any iteration.

The rationale behind this algorithm is to assume to know the optimal value and as soon as we realize it is not correct we change it.

Let us first give an informal description of the algorithm:

- $\delta$  is the displacement: how much below the function is with respect to the best value obtained so far;
- reference value  $f_{rec} = \underline{f}$ . At the beginning is the value at the first iterate and then we define the target value as the difference between the reference value and some  $\delta$  (at the beginning  $\delta_0$ ).

---

ALGORITHM 20.1 Pseudocode for target level stepsize.

---

```
1: procedure SGPTL( $f, g, x, i_{max}, \beta, \delta_0, R, \rho$ )
2:    $r \leftarrow 0;$ 
3:    $\delta \leftarrow \delta_0;$ 
4:    $f_{ref} \leftarrow f_{rec} \leftarrow f(x);$ 
5:    $i \leftarrow 1;$ 
6:   while ( $i < i_{max}$ ) do
7:      $g = g(x);$ 
8:      $\alpha = \beta(f(x) - (f_{ref} - \delta)) / \|g\|^2;$ 
9:      $x \leftarrow x - \alpha g;$ 
10:    if ( $f(x) \leq f_{ref} - \delta/2$ ) then
11:       $f_{ref} \leftarrow f_{rec};$ 
12:       $r \leftarrow 0;$ 
13:    else
14:      if ( $r > R$ ) then
15:         $\delta \leftarrow \delta\rho;$ 
16:         $r \leftarrow 0;$ 
17:      else
18:         $r \leftarrow r + \alpha \|g\|;$ 
19:      end if
20:    end if
21:   end while
22:    $f_{rec} \leftarrow \min\{f_{rec}, f(x)\};$ 
23:    $i \leftarrow i + 1;$ 
24: end procedure
```

---

At this point we defined the algorithm and we are ready to implement it, except for the fact that we need to choose of a lot of parameters. A way to choose them is to use the ML approach: try many possible values.

Two big issues of this algorithm are that it does not provide a good stopping criterion and it is very sensitive to many parameters.

## 20.2 Deflected subgradient

The idea is to use the same trick of ball-step (also called primal-dual).

Let us assume that our function was differentiable. The subgradient method collapses to the gradient method and we know that the gradient method does not provide a good convergence. Yet, deflection is possible:  $d^i = \gamma^i g^i + (1 - \gamma^i)d^{i-1}$ ,  $x^{i+1} = x^i - \alpha^i d^i$ . We can prove that  $d^i$  approximates the subgradient. We can also prove that the algorithm converges in the end. The parameters of this algorithm are two:  $\beta$  (stepsize) and  $\gamma$  (deflection). In order to choose them we have two different approaches:




FIGURE 20.1: There are two cases: either the function value is significantly below the reference (for example  $\frac{\delta}{2}$  below the reference) or it's not. If we are in the “happy case” (we just move the reference to the best value). For what concerns the “unhappy case”, if we are unhappy for 1 iteration, no problem. Two iterations? no problem. Many iterations? Problem: after some iterations in which we are not improving it means that we have to decrease the reference values. How?  $r$  is updated at each bad step and reset when a good step occurs. When  $r$  gets too large we decrease  $\delta$  and reset everything.

STEP SIZE-RESTRICTED  $\equiv$  deflection-first. We first choose  $\beta$  and when choosing  $\alpha$  we need to take into account  $\beta$ :  $\alpha^i = \frac{\beta^i(f(x^i) - f_*)}{\|d^i\|} \wedge \beta^i \leq \gamma^i$  “as deflection  $\nearrow$ , stepsize has to  $\searrow$ ”;

DEFLECTION-RESTRICTED  $\equiv$  stepsize-first. We first choose  $\gamma$ , then we pick a step size that depends on  $\gamma$ :

$$(DSS) \wedge \frac{\alpha^{i-1} \|d^{i-1}\|}{(f(x^i) - f_*) + \alpha^{i-1}} \|d^{i-1}\| \leq \gamma^i$$

“as  $f(x^i) \rightarrow f_*$ , deflection  $\searrow$ ”

This algorithm gets the optimal  $O(1/\varepsilon^2)$  on average, sadly not worst case.

### 20.3 Smoothed gradient methods

Let us assume that the target function is a **Lagrangian function**.

**Definition 20.3** (Lagrangian function). *Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function of the following shape:*

$$f(x) = \max\{x^T A z : z \in Z\}$$

where  $Z$  is convex and bounded.

Let us assume that  $Z$  is also compact.

A graphical example in the case of  $f(x) = |x| = \max\{x, -x\} = \max\{zx : z \in [-1, 1]\}$  is shown in Figure 20.2.

In the case of the absolute value, the nasty trick is “to make it have only one optimal solution” in the point 0.




FIGURE 20.2: Let us take the absolute value function  $f(x) = |x|$ . This function would be differentiable, if it wasn't for some nasty points (in this case 0, were the optimization problem has many optimal solutions). In 0 there are many subgradients, so it has many optima.

This result is obtained adding a small quadratic term:  $f(x) = \max\{x^T A z - \mu \|z\|^2 : z \text{ in } G\}$  that is shown in Figure 20.3. At this point the new function  $f_\mu$  is not the original function anymore, but it is very close to it whenever the  $\mu$  is small.

Notice that this new function is smooth (differentiable).




FIGURE 20.3: Geometric intuition of the usage of variable  $\mu$ .

It might be not ery easy to compute once chosen the value for  $\mu$ .

We are solving a problem which is different from the original one and would be exactly the same if  $\mu = 0$ .

On the other hand for  $\mu = 0$  we have the problem which is not differentiable once again, so we need to keep close to 0 but not too close.

At this point we are in the situation of  $f_\mu(x) \leq f(x) \leq f_\mu(x) + \mu R$ , such that as  $\mu \searrow 0$ , “ $\operatorname{argmin} \{f_\mu(x)\} \rightarrow x_*$ ”.

The new function is not only convex, but it is also Lipschitz continuous:  $\nabla f_\mu$  Lipschitz with  $L = \|A\|^2/\mu$ , but it is “less and less Lipschitz” as  $\mu \searrow 0$ .

**Fact 20.3.** *If  $f_* > -\infty$  and picking a very special value of  $\mu$  ( $\mu = \varepsilon/(2R)$ ), then an appropriate ACCG obtains  $f(x^i) - f_* \leq \varepsilon$  for  $i \geq 4 \cdot \|A\| \cdot \|x_*\| \cdot \sqrt{R}/\varepsilon$ .*

We observe that the convergence is much better, because it depends on  $O(1/\varepsilon)$  instead of  $O(1/\varepsilon^2)$ .



FIGURE 20.4: At the beginning we will make a lot of bad steps (the upper gray line). We can improve (pick the black line) changing the  $\varepsilon$  value and we obtain something that looks more stable. The more precision we want, the smaller the step we make. At the ending it pays, but at the beginning it is not so.

## 20.4 Cutting-plane algorithm

We cheated to get first order information, we want to do more. We want to cheat and have also the second order information.

We want to use the same idea of limited memory quasi Newton methods. Using some limited memory of the Hessian, in order to understand the curvature.

The point is that the directional derivatives are defined and (if computed massively) give me a hint of the curvature of the function (cfr. Figure 20.5). Notice that it is not possible to build a matrix, because it is not defined.

Let us say we have performed  $i$  iterates, then we have collected  $i$  subgradients ( $\mathcal{B} = \{(x^i, f^i = f(x^i), g^i \in \partial f(x^i))\}$   $\equiv$  bundle of first-order information) and function values.

We can now define a piece wise linear function defined as the maximum of the first order model:

$$f_{\mathcal{B}}(x) = \max\{f^i + g^i(x - x^i) : (x^i, f^i, g^i) \in \mathcal{B}\}$$

At this point we can apply Newton method: first we minimize the model and the use the minimum as next point. We collect information in that specific point and then we repeat.

Notice that the model is always below the objective function ( $f_{\mathcal{B}}(x) \leq f(x) \forall x$ ), hence  $\min\{f_{\mathcal{B}}(x)\} \leq f_*$ , so  $x_{\mathcal{B}}^* \in \operatorname{argmin}\{f_{\mathcal{B}}(x)\} \approx x_*$ .




FIGURE 20.5: The idea is that we do not compute a subgradient and discard it. We store it, thus resulting in a model

This function has many kinky points, so how to minimize it? Dirty trick from Ricerca Operativa:

$$\min\{f_{\mathcal{B}}(x)\} = \min\{v : v \geq f^i + g^i(x - x^i), \text{ where } (x^i, f^i, g^i) \in \mathcal{B}\}$$

And on this problem, we can use the simplex method.




FIGURE 20.6: A geometric representation of how the model (blue line) changes after one iteration.

At this point we have obtained both an upperbound and a lower bound, which go one towards the other.

We may decide to stop iterating when they are “close enough”.

The problem is that at each step we need to solve a minimization problem and the convergence is very slow.

It's also possible that the blue function (the model) does not have a minimum (unbounded below).

How to overcome this problem? Use so-called bundle methods.

## 20.5 Bundle methods

The intuition behind this is that we want to keep quite close to the point where the model corresponds to the function, because the further we get the more the difference from the function.

A way to express this is to add a quadratic quantity to the function that grows when we move outside the current point.

**Definition 20.4** (Stabilized master problem). *We term **stabilized master problem** the following*

$$\min\{f_{\mathcal{B}}(x) + \mu\|x - \bar{x}\|^2/2\}$$

This improved model cannot be undounded below (quadratic function), but we need to choose  $\mu$  and  $\bar{x}$  wisely.

A possible way is to move the **stability center** whenever the current value is better than the best encountered so far.

---

ALGORITHM 20.2 Pseudocode for boundle method.

---

```

1: procedure PBM( $f, g, \bar{x}, m_1, \varepsilon$ )
2:   choose  $\mu$ ;
3:    $\mathcal{B} \leftarrow \{(\bar{x}, f(\bar{x}), g(\bar{x}))\}$ ;
4:   while ( true ) do
5:      $x^* \leftarrow \operatorname{argmin} \{f_{\mathcal{B}}(x) + \mu\|x - \bar{x}\|^2/2\}$ ;
6:     if ( $\mu\|x^* - \bar{x}\|_2 \leq \varepsilon$ ) then
7:       break;
8:     end if
9:     if ( $f(x^*) \leq f(\bar{x}) + m_1(f_{\mathcal{B}}(x^*) - f(\bar{x}))$ ) then
10:       $\bar{x} \leftarrow x^*$ ;
11:      possibly decrease  $\mu$ ;
12:    else
13:      possibly increase  $\mu$ ;
14:    end if
15:     $\mathcal{B} \leftarrow \mathcal{B} \cup (x^*, f(x^*), g(x^*))$ ;
16:  end while
17: end procedure

```

---

This algorithm may never move (without cycling, luckily), but at least we gained some information.

The bundle method converges in few steps, although each step is quite costly.

We reached a point where to solve an unconstrained problem we need to solve a constrained one, so from next lecture we will start dealing with constrained optimization problems.

# 21 30th of November 2018 — A. Frangioni

## 21.1 Constrained optimization

In this lecture we address the problem of finding the **optimum** of a function in a subset of its domain, called  $X$ . The term optimum differs from the minimum, because the optimum in that subset may not be a minimum of the whole function.

$$f_* = \min\{f(x) : x \in X\}$$

**Definition 21.1** (Local optimum). *Given a function  $f$  and a constraint set  $X$ , we denote **local optimum** the point where the function assumes the minimum value inside the set  $X$ . Formally,  $\min\{f(x) : x \in \mathcal{B}(x_*, \varepsilon) \cap X\}$  for some  $\varepsilon > 0$ .*

Notice that the only points in which the constraint adds some informations are the ones on the boundary, as shown in Figure 21.1




FIGURE 21.1: The red line is level set of the function corresponding to the smallest value that touches the set  $X$ . The point in the intersection is not a saddle point of the function  $f$ , although it is the minimum.

There are two kinds of constraints:

**FAKE ONES:** this first kind is such that the minimum of the function lies inside the set  $X$ , hence there is no need to use the constraints at all;

**REAL ONES:** when the optimal is on the boundary. This is the case of linear functions, because the gradient is constant  $\nabla f(x) = c$ .

At this point we want to decide if a point on the boundary is an optimum. In this context it is important how the boundary is defined.

### 21.1.1 Linear equality constraints

A constraint of this kind is very simple: it is a subspace, as shown in Figure 21.2.

$\min\{f(x) : Ax = b\}$ , where the rank of  $A$  counts the number of linearly independent rows.




FIGURE 21.2: Linear constraint and a point on the boundary.




FIGURE 21.3: The gradient is orthogonal to the level set in that point, when the function is smooth. The same holds for matrix  $A$ .




FIGURE 21.4: If we take any other point in the space it has to be orthogonal to  $A$ .

Let us assume that there are not linearly independent rows in  $A$ , then or this behaviour is reflected in  $B$  or the system does not have any solution.

In the case of presence of linearly dependent columns such columns may be eliminated to ease the computation without loss of information.

The intuition behind what follows is that each linear constraint kills one degree of freedom, formally  $\det(A_B) \neq 0 \Rightarrow Ax = b \equiv x_B = A_B^{-1}(b - A_N x_N) \Rightarrow$

We want to extract a submatrix  $A_B$  from  $A \in M(m, n, \mathbb{R})$ , such that  $A_B \in M(m, \mathbb{R})$  and then the system induces a partitioning in the variables as well.

$A = [A_B, A_N]$ ,  $x = [x_B, x_N]$ , so the system becomes  $A_B x_B + A_N x_N = b$  and (since  $A_B$  is non singular)  $x_B + A_B^{-1} x_N = A_B^{-1} b$  in other words, given the independent variables we can compute the values of the dependent ones and this is a linear operation.

The original optimization problem becomes an optimization problem on a reduced space, formally  $\min\{r(w) = f(Dw + d) : w \in \mathbb{R}^{n-m}\}$ , where  $D = \begin{bmatrix} -A_B^{-1} A_N \\ I \end{bmatrix}$  and  $d = \begin{bmatrix} A_B^{-1} b \\ 0 \end{bmatrix}$ .

For each point in the smaller space we can compute the function in the larger space.

How can we compute the gradient of  $r(w)$ ? The gradient is  $\nabla r(w) = D^T \nabla f(Dw + d)$ . The fact that  $w^*$  is an optimum implies that  $\nabla r(w^*) = 0$ .

$$D = \begin{bmatrix} -A_B^{-1} A_N \\ I \end{bmatrix} \text{ then } AD = [A_B, A_N] \cdot \begin{bmatrix} -A_B^{-1} A_N \\ I \end{bmatrix} = -A_B \cdot A_B^{-1} \cdot A_N + A_N = 0.$$

Now the point is taking a multiple of matrix  $A$ , finding a feasible  $x$  and corresponding  $w$  (because there is a bijection), then finding the value  $\mu$  that allows the equality.

**Theorem 21.1.** Let  $Ax = b$  be a linear system and  $w$  such that  $x = [A_B^{-1}(b - A_N w), w]$ . If  $\exists \mu \in \mathbb{R}^m$  s.t.  $\mu A = \nabla f(x)$  then  $r(w) = D^T \nabla f(x) = 0$ , see Figure 21.5. In other words, it is equivalent to find a stationary point  $x$  for the original problem (P) or finding the stationary point  $w$  for (R).

**Definition 21.2** (Poorman's Karush Kuntaker conditions). A point is a good candidate for being a minimum of the constrained problem if and only if it satisfy **Poorman's KKT conditions**, namely the problem is feasible and that  $\exists \mu \in \mathbb{R}^m$  s.t.  $\mu A = \nabla f(x)$ .

**Theorem 21.2.** Let  $f$  be a convex function, then KKT conditions are enough for optimality.

A very naive explanation of the theorem is that if the function is convex also the restriction is convex and a tationaly point of a convex function is a minimum.

Our idea is to characterize the directions we can move along in order to find new points that satisy the constraint. Formally,  $Ax = b$  is our constraint and we want to move towards  $x + d$  and stay in the feasible region. How?  $A(w + d) = b \Leftrightarrow Ax + Ad = b \Leftrightarrow Ad = 0$ , since  $Ax = b$ . The only way to move along the constraint is choosing a direction which scalar product with  $A$  is 0, hence 0 scalar product with the gradient.

From now on we would like to study the behaviour on contrained problems where the constraints are equalities, but inequalities.

In order to do this we need some mathematical background.

### 21.1.2 Background for linear inequality constraints

**Definition 21.3** (Tangent cone). We call **tangent cone** of  $X$  at  $x$   $\mathbf{T}_X(x) = t$ .

$$\{d \in \mathbb{R}^n : \exists \{z_i \in X\} \rightarrow x \wedge \{t_i \geq 0\} \rightarrow 0 \text{ s.t. } d = \lim_{i \rightarrow \infty} \frac{z_i - x}{t_i}\}$$




FIGURE 21.5: When the boundary has this shape we can move along directions that point inside the constraints. Tangent directions are not allowed.




FIGURE 21.6: Geometric representation of tangent cone, which is the region of the space where we can pick the directions. The intuition is to zoom in  $x$  and the result of this zooming is a cone.

**Theorem 21.3.** Let  $\mathcal{C}$  be a cone,  $\forall x \in \mathcal{C} \ \alpha x \in \mathcal{C}, \ \forall \alpha > 0$ .

**Theorem 21.4.** Given a function  $f$ , where  $x$  is a **local** optimum  $\langle \nabla f(x), d \rangle \geq 0 \ \forall d \in T_X(x)$ .

*Proof. Proof by contraddiction:* Assume  $\exists d \in T_X(x)$  such that  $\langle \nabla f(x), d \rangle < 0$ , but  $x$  is a local optimum.

By definition  $\exists X \supset \{z_i\} \rightarrow x$  and  $\{t_i\} \rightarrow 0$  such that  $d = \lim_{i \rightarrow \infty} \frac{z_i - x}{t_i}$ .

First order Taylor  $f(z_i) - f(x) = \langle \nabla f(x), (z_i - x) \rangle + R(z_i - x)$ .

$$\begin{aligned}
 \lim_{i \rightarrow \infty} \frac{f(z_i - x)}{t_i} &= \lim_{i \rightarrow \infty} \left\langle \nabla f(x), \frac{z_i - x}{t_i} \right\rangle + \frac{R(z_i - x)}{t_i} \\
 &\stackrel{*}{=} \langle \nabla f(x), d \rangle + \lim_{i \rightarrow \infty} \frac{R(z_i - x)}{t_i} \\
 &\stackrel{(1)}{=} \langle \nabla f(x), d \rangle \\
 &< 0
 \end{aligned} \tag{21.1}$$

Where,  $\stackrel{(1)}{=}$  follows from  $\lim_{i \rightarrow \infty} \frac{R(z_i - x)}{t_i} = 0$  by Taylor.  $\square$

**Observation 21.1.** The optimum of Theorem 21.4 is global when the function is convex, because in that case  $X \subseteq x + T_X(x)$ . For a geometric idea see Figure 21.7.




FIGURE 21.7: Convex function.

**Observation 21.2.** Notice that the rule  $\langle \nabla f(x), d \rangle \geq 0 \forall d \in T_X(x) \Rightarrow x$  local optimum does not hold. Let us see a counter example:  $\min\{x_2 : x_2 \geq x_1^3\}$ , displayed in Figure 21.9.




FIGURE 21.8: Let us suppose we pick the direction towards the left part of the function in the saddle point. That direction is promising and actually the value of the function decreases. In this case the problem is that the constraint is not convex.

Now we need a more manageable object for  $T_X(x)$ .

**Definition 21.4** (Cone of feasible directions).

Intuitively, we are in  $x$  and we want to find all directions **feasible cone** such that there exist small but not 0 steps such that all the points on this direction are feasible. Formally, a **feasible cone** is  $F_X(x) = \{d \in \mathbb{R}^n : \exists \bar{\varepsilon} > 0 \text{ such that } x + \varepsilon d \in X, \forall \varepsilon \in [0, \bar{\varepsilon}]\}$ .

**Fact 21.5.** The properties of such cone are:

1.  $T_X$  closed,  $F_X$  in general not (hence the cone of feasible directions is the tangent cone minus the tangent directions);
2.  $cl(F_X) \subseteq T_X$ , where  $cl(F_X)$  is the closure of the cone of feasible directions;

3. If  $X$  convex then the cones coincide:  $T_X$  and  $F_X$  convex and  $\text{cl } F_X = T_X$ .

We are now interested in finding a better characterization of the cone of feasible directions, hence we introduce a new characterization of the set of constraints  $X$ .

FIRST REPRESENTATION:

$$X = \{x \in \mathbb{R}^n : g_i(x) \leq 0 \ i \in \mathcal{I}, \ h_j(x) = 0 \ j \in \mathcal{J}\}$$

where  $\mathcal{I}$  is the set of inequality constraints and  $\mathcal{J}$  is the set of equality constraints;

SECOND REPRESENTATION:

$$X = \{x \in \mathbb{R}^n : G(x) \leq 0, H(x) = 0\}$$

where  $G = [g_i(x)]_{i \in \mathcal{I}} : \mathbb{R}^n \rightarrow \mathbb{R}^{|\mathcal{I}|}$  and  $H = [h_i(x)]_{i \in \mathcal{J}} : \mathbb{R}^n \rightarrow \mathbb{R}^{|\mathcal{J}|}$ ;

THIRD REPRESENTATION: (hiding equalities)

$$X = \{x \in \mathbb{R}^n : g_i(x) \leq 0 \ i \in \mathcal{I}, \ h_j(x) \leq 0 \ \wedge \ h_j(x) \geq 0 \ j \in \mathcal{J}\}$$

FOURTH REPRESENTATION: (hiding inequalities into a single function)

$$X = \{x \in \mathbb{R}^n : g(x) = \max\{g_i(x) : i \in \mathcal{I}\} \leq 0 \ i \in \mathcal{I}, \ h_j(x) = 0 \ j \in \mathcal{J}\}$$

**Definition 21.5** (Active constraints). We term **active constraints** at  $x \in X$  the following set

$$\mathcal{A}(x) = \{i \in \mathcal{I} : g_i(x) = 0\} \subseteq \mathcal{I}$$

Let us introduce some useful notation on the subject: let  $\mathcal{B} \subseteq \mathcal{I}$  a subset of indices.

We denote  $G_{\mathcal{B}} = [g_i(x)]_{i \in \mathcal{B}} : \mathbb{R}^n \rightarrow \mathbb{R}^{|\mathcal{B}|}$  the corresponding set of inequalities.

**Definition 21.6** (First-order feasible direction cone). We term **First-order feasible direction cone at  $x \in X$** :

$$D_X(x) = \{d \in \mathbb{R}^n : \langle \nabla g_i(x), d \rangle \leq 0 \ i \in \mathcal{A}(x), \ \langle \nabla h_j(x), d \rangle = 0 \ j \in \mathcal{J}\} = \{d \in \mathbb{R}^n : (JG_{\mathcal{A}(x)}(x))d \leq 0,$$

Intuitively, the fact that we are looking at the active set means that we are zooming very close to 0.

In this cone we require that all the directions inside it have a negative scalar product with the gradient of the constraints.

A visual example of a first order feasible direction cone is displayed in Figure 21.9.

**Fact 21.6.** The tangent cone is a subset of the first-order feasible direction cone. Formally,  $T_X(x) \subseteq D_X(x)$ .




FIGURE 21.9: The first-order feasible direction cone is made of those directions that are orthogonal to the gradient of the constraints  $g_1(x)$  and  $g_2(x)$  in  $x$ .




FIGURE 21.10: The circle represents the quadratic constraint, while the semi-plane represents the first degree constraint.

We would like the first-order feasible direction cone to be exactly equal to the tangent cone and this is almost always true, except for some pathological cases.

**Example 21.1.** Let our minimum problem be  $\min\{\dots : x_1^2 + (x_2 - 1)^2 - 1 \leq 0, x_2 \leq 0\}$ .  
The plot of such functions is shown in Figure 21.10.

Our claim is that the only feasible point is  $X = \{x = [0, 0]\}$ .

The only feasible direction is 0, hence cone of feasible direction and the tangent cone are the singleton  $\{[0, 0]\}$ .

On the other hand, the set of directions that have non-negative scalar product with both  $g_1$  and  $g_2$  are all the  $x$  axis.

We would like to ensure we are not in one of these pathological cases and to do this we introduce some conditions.

**Fact 21.7.** The following holds:

AFFINE CONSTRAINTS (AFFC): Let  $g_i$  and  $h_j$  be affine constraints. Then,  $\forall i \in \mathcal{I}$  and  $j \in \mathcal{J}$   $T_X(x) = D_X(x) \forall x \in X$ .

**SLATER'S CONDITION (SLAC):** Let  $g_i$  convex  $\forall i \in \mathcal{I}$  and let  $h_j$  affine  $\forall j \in \mathcal{J}$   $\exists \bar{x} \in X$  s.t.  $g_i(\bar{x}) < 0 \forall i \in \mathcal{I}$ . Then  $T_X(x) = D_X(x) \forall x \in X$ ;

**LINEAR INDEPENDENCE (LINI):**  $\bar{x} \in X \wedge$  the vectors  $\{\nabla g_i(\bar{x}) : i \in \mathcal{A}(\bar{x})\} \cup \{\nabla h_j(\bar{x}) : j \in \mathcal{J}\}$  linearly independent  $\implies T_X(\bar{x}) = D_X(\bar{x})$ . Among all these conditions this is the only local one.

It goes without saying that we cannot check all the directions in order to exclude the nasty pathological cases.

**Definition 21.7** (Dual cone). Let  $D_X$  be a **polyhedral cone**  $\mathcal{C} = \{d \in \mathbb{R}^n : Ad \leq 0\}$ , for some  $A \in \mathbb{R}^{k \times n}$ .

We term **dual cone**  $\mathcal{C}^* = \{c = \sum_{i=1}^k \lambda_i A_i : \lambda \geq 0\}$ .

**Lemma 21.8** (Farka's lemma). Intuitively, this lemma says that pick a vector: either it belongs to the dual cone or there exists a vector in the polyhedral cone which has a negative scalar product with it.

Equivalently, either  $c \in \mathcal{C}^*$  or  $c \notin \mathcal{C}^*$ .

More formally, either  $\exists \lambda \geq 0$  s.t.  $c = \sum_{i=1}^k \lambda_i A_i$  or  $\exists d$  s.t.  $Ad \leq 0 \wedge \langle c, d \rangle > 0$ .

**Theorem 21.9** (Karush-Kuhn-Tucker conditions). Let us assume that we found an optimal solution  $x_*$  and the constraints qualification holds.

Then  $\exists \lambda \in \mathbb{R}_+^{|\mathcal{I}|}$  and  $\mu \in \mathbb{R}^{|\mathcal{J}|}$  such that

$$\nabla f(x) + \sum_{i \in \mathcal{A}(x)} \lambda_i \nabla g_i(x) + \sum_{j \in \mathcal{J}} \mu_j \nabla h_j(x) = 0$$

It is interesting to notice that we did not impose  $\mu \geq 0$ . Let us take an equality constraint  $h_j(x) = 0$ . This is equivalent to write  $h_j(x) \leq 0 \wedge h_j(x) \geq 0$ , thus leading to two different multipliers, say  $\lambda_j^+$  and  $\lambda_j^-$ .

The term of the sum concerning  $h_j$  looks like this  $\lambda_j^+ \nabla h_j(x_*) - \lambda_j^- \nabla h_j(x_*) = (\lambda_j^+ - \lambda_j^-) \cdot \nabla h_j(x_*)$ , where both  $\lambda_j^+$  and  $\lambda_j^-$  are  $\geq 0$ , hence their difference (denoted by  $\mu_j$ ) may be either positive or negative.

**Fact 21.10.** The Karush-Kuhn-Tucker conditions are also written as:

FEASIBILITY:  $x \in X \equiv g_i(x) \leq 0 \quad i \in \mathcal{I}, \quad h_j(x) = 0 \quad j \in \mathcal{J}$

KKT-G:  $\nabla f(x) + \sum_{i \in \mathcal{I}} \lambda_i \nabla g_i(x) + \sum_{j \in \mathcal{J}} \mu_j \nabla h_j(x) = 0$

COMPLEMENTARITY SLACKNESS:  $\sum_{i \in \mathcal{I}} \lambda_i g_i(x) = 0$

where the second and the third equations formalize the definition of KKT, where the terms of the first sum should be summed only if their constraints are active.

**Fact 21.11.** Let  $(P)$  be a convex problem. In this case, if the Karush-Kuhn-Tucker conditions hold then  $x$  is a global optimum.

## 22 6th of December 2018 — A. Frangioni

### 22.1 Duality



Do you recall?

The KKT-G condition:

$$\nabla f(x) + \sum_{i \in \mathcal{I}} \lambda_i \nabla g_i(x) + \sum_{j \in \mathcal{J}} \mu_j \nabla h_j(x) = 0$$

We would like to understand what is the function of which this is the gradient.

**Definition 22.1** (Lagrangian function). *Let  $(P)$  be our minimum problem over  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . We term **Lagrangian function** the following:*

$$L(x, \lambda, \mu) = f(x) + \sum_{i \in \mathcal{I}} \lambda_i g_i(x) + \sum_{j \in \mathcal{J}} \mu_j h_j(x)$$

**Fact 22.1.** *A necessary condition for optimality is that the gradient of the Lagrangian function is 0. Formally,  $\nabla L(x; \lambda, \mu) = 0$ .*

A first approach to use the Lagrangian function would be taking  $\lambda$  and  $\mu$  such that the Lagrangian is positive semidefinite.

But this is a too strict requirement, in fact we impose:

**Definition 22.2** (Critical cone). *Let us assume  $(x, \lambda, \mu)$  satisfies (KKT). We define the **critical cone** as*

$$C(x, \lambda, \mu) = \left\{ d \in \mathbb{R}^n : \begin{array}{ll} < \nabla g_i(x), d > = 0 & i \in \mathcal{A}(x) \text{ s.t. } \lambda_i^* > 0 \\ < \nabla g_i(x), d > \leq 0 & i \in \mathcal{A}(x) \text{ s.t. } \lambda_i^* = 0 \\ < \nabla h_j(x), d > = 0 & i \in \mathcal{J} \end{array} \right\}$$

**Theorem 22.2.** *Let us assume we have a point  $(x, \lambda, \mu)$  that satisfies the Karush-Khun-Tucker conditions and satisfies the linear independence of the constraints. If  $x$  is local optimum then  $d^T \nabla_{xx}^2 L(x, \lambda, \mu) d \geq 0 \forall d \in C(x, \lambda, \mu)$ .*

*Informally, if the hypothesis holds, then the Hessian of the Lagrangian function is  $\succeq 0$  on the critical cone.*

**Observation 22.1.**  $(x, \lambda, \mu)$  satisfies (KKT)  $\wedge \nabla_{xx}^2 L(x, \lambda, \mu) \succ 0$  on  $C(x, \lambda, \mu)$  then  $x$  local optimum.

We would like to say something more about  $\lambda$  and  $\mu$ . Until now we considered the Lagrangian as a function of  $x$ , but what if we consider the Lagrangian in terms of  $\lambda$  and  $\mu$ ?

## 22.2 Lagrangian duality

**Definition 22.3** (Lagrangian relaxation). Let us consider the Lagrangian function. We term **Lagrangian relaxation** the function where we fixed  $\lambda$  and  $\mu$  and minimize on  $x$ :

$$\psi(\lambda, \mu) = \min\{L(x, \lambda, \mu) : x \in \mathbb{R}^n\}$$

The relaxation leads to an unconstrained problem and we learnt how to solve one of those. This Lagrangian relaxation leads to the definition of **lagrangian dual**  $\psi$ .

**Property 22.3.** The dual function has the following properties:

- $\psi$  is concave, but  $\psi(\lambda, \mu) = -\infty$ ;
- $\psi$  is non differentiable, although  $f$ ,  $g_i$  and  $h_j$  are;
- Let  $\bar{x}$  be optimal in  $(R_{\lambda, \mu})$ , then  $[\sum_{i \in \mathcal{I}} \nabla g_i(\bar{x}) + \sum_{j \in \mathcal{J}} \nabla h_j(\bar{x})] \in \partial\psi(\lambda, \mu)$
- $\forall$  fixed  $\lambda \geq 0, \mu, \bar{x} \in X$   $\psi(\lambda, \mu) = \min_x L(x, \lambda, \mu) \leq L(\bar{x}, \lambda, \mu) \leq f(\bar{x})$

$$\psi(\lambda, \mu) = \begin{pmatrix} \lambda_1 g_1(\bar{x}) \\ \vdots \\ \lambda_k g_k(\bar{x}) \\ \lambda_1 \mu_1(\bar{x}) \\ \vdots \\ \lambda_p \mu_p(\bar{x}) \end{pmatrix} \text{ is such that } \begin{pmatrix} g_1(\bar{x}) \\ \vdots \\ g_k(\bar{x}) \\ \mu_1(\bar{x}) \\ \vdots \\ \mu_p(\bar{x}) \end{pmatrix} \text{ belongs to the supergradient.}$$

**Theorem 22.4** (Weak duality).  $\forall$  fixed  $\lambda \in \mathbb{R}^+ \cup \{0\}, \mu \in \mathbb{R}$  such that  $\psi(\lambda, \mu) \leq v(P)$  and let us take any feasible  $\bar{x} \in X$  and  $g(\bar{x}) \leq 0, h(\bar{x}) = 0$ .

It can be proved that  $\psi(\lambda, \mu) = \min_x L(x, \lambda, \mu) \leq L(\bar{x}, \lambda, \mu) \leq f(\bar{x})$ .

*Proof.*  $L(x, \lambda, \mu) = f(x) + \sum_i \lambda_i g_i(x) + \sum_j \mu_j h_j(x)$ . Let us assume we have some  $\bar{\lambda}, \bar{\mu}, \bar{x}$ , where  $\bar{x}$  is feasible (aka  $g_i(\bar{x}) \leq 0$  and  $h_i(\bar{x}) = 0$  and  $\bar{\lambda} \geq 0$ ).

The value of the dual function is

$$\psi(\bar{\lambda}, \bar{\mu}) = \min_x \{f(x) + \sum_i \lambda_i g_i(x) + \sum_j \mu_j h_j(x)\} \leq f(\bar{x}) + \sum_i \bar{\lambda}_i g_i(\bar{x}) + \sum_j \bar{\mu}_j h_j(\bar{x}),$$

where the last term is cancelled, since  $h_i(\bar{x}) = 0$ .

On the other hand,  $\sum_i \bar{\lambda}_i g_i(x) \leq 0$ , which implies that  $f(x) \leq 0$ .  $\square$

**Observation 22.2.** This theorem gives us the way to prove that a point is an optimum. Let us assume that we found  $\lambda_* \leq 0, x_*$  and  $\mu_*$  such that an equality holds:  $\psi(\lambda_*, \mu_*) = f(x_*)$ . The value of  $\psi$  gives us the information about how far we are from the optimum. Let us assume we have  $\bar{\lambda}, \bar{\mu}$  and  $\bar{x}$  then  $\psi(\bar{\lambda}, \bar{\mu}) \leq f(\bar{x}) \leq \psi(\bar{\lambda}, \bar{\mu}) + \varepsilon$  and this tells that we are far from the optimum of a factor  $\varepsilon$ .

If everything in the original function was convex then the Lagrangian is convex and the computations are easy.

At this point we want to find the biggest lower bound and it may be computed since the function is concave.

$$\max\{\psi(\lambda, \mu) : \lambda \in \mathbb{R}_+^{|\mathcal{I}|}, \mu \in \mathbb{R}^{|\mathcal{J}|}\}$$

Notice that the constraints on  $\lambda$  are very easy, so the problem may be considered somehow unconstrained.

We can use local methods to compute the maximum of this function.

If we are able to compute the gradient of  $\psi$  then we have the supergradients of the function  $f$ .

Is (P) equal to (D)? Yes, provided that everything is convex. In general the Lagrangian gives a lowerbound.

$$v(D) \leq v(P)$$

**Example 22.1.** Let us take a concave objective function  $\min\{-x^2 : 0 \leq x \leq 1\}$ , such that its Lagrangian function is  $L(x, \lambda) = -x^2 + \lambda_1(x - 1) - \lambda_2x$ . It goes without saying that the Lagrangian function is unbounded below (upside-down parabola).

Formally,  $\psi(\lambda) = \min_{x \in \mathbb{R}} L(x, \lambda) = -\infty$ ,  $\forall \lambda \in \mathbb{R}^2$ , which means that the Lagrangian dual is  $v(D) = -\infty < v(P) = -1$ .

**Theorem 22.5.** Let us assume that  $f$ ,  $g$  and  $h$  are convex, and constraints qualification holds. If the problem has an optimum then the value of the primal and the value of the dual are the same.

$$\text{Formally, } T_X(x_*) D_X(x_*) \Rightarrow v(D) = v(P).$$

*Proof.* We are assuming  $x_*$  is an optimum, so the necessary conditions hold, hence we have the Karush Khun Tucker conditions, then we can find  $(\lambda^*, \mu^*)$  that satisfy the KKT with  $x$ .

Then our claim is that  $(\lambda^*, \mu^*)$  is an optimal solution of the dual (D) and the value of the dual is equal to the value of the primal.

$x_*$  is a stationary point for the Lagrangian function, since it satisfies the KKT conditions then  $x_*$  is exactly the minimum, since the Lagrangian function is convex.

Hence, the value of the dual  $v(D) \geq \psi(\lambda^*, \mu^*) = L(x_*, \lambda^*, \mu^*) = f(x_*) = v(P) \geq v(D)$ .  $\square$

How many solution may  $\psi(\lambda, \mu)$  have? In principle many, but what if  $f$  is strongly convex, than everything in the Lagrangian is strongly convex, then the solution of  $\psi$  is unique and it is also differentiable, but typically not two times differentiable. At this point the Lagrangian dual has a single solution and the optimum of the Lagrangian dual corresponds to an optimum for  $f$ .

We only translated a minimum problem on convex functions to another minimum problem on other convex functions. We will see soon that in some cases this approach is advantageous in others it is not.

## 22.3 Specialized dual

### 22.3.1 Linead programs

$$(P) \min\{cx : Ax \geq b\}$$

Lagrangian function:  $L(x, \lambda) = cx + \lambda(b - Ax) = \lambda b + (c - \lambda A)x$

Dual function:

$$\psi(\lambda) = \min_{x \in \mathbb{R}^n} L(x, \lambda) = \begin{cases} -\infty & \text{if } c - \lambda A \neq 0 \\ \lambda b & \text{if } c - \lambda A = 0 \end{cases}$$

Since we can find a minimum only on a function of the second case we have that:

$$(D) \max \{\psi(\lambda) : \lambda \geq 0\} \equiv \max \{\lambda b : \lambda A = c, \lambda \geq 0\}$$

Since the lagrangian is far from having an unique optimal solution the dual does not have a unique maximum so the do not match usually.

### 22.3.2 Quadratic programs

Notice that thwe quadratic case is simpler than the linear case.

$$(P) \min \left\{ \frac{1}{2} \|x\|_2^2 : Ax = b \right\} \text{ (linear least-norm solution)}$$

$L(x, \mu) = \frac{1}{2} \|x\|_2^2 + \mu(Ax - b)$ ,  $\nabla_x L = x + \mu A = 0 \iff x = -\mu A$ , which is stricltly convex since  $x^2$  is strictly convex.

$$\psi(\mu) = \min_{x \in \mathbb{R}^n} L(x, \mu) = L(-\mu A, \mu) = -\frac{1}{2} \mu^T (AA^T) \mu - \mu b$$

$$\text{Then the Lagrangian dual (D) } \max \left\{ -\frac{1}{2} \mu^T (AA^T) \mu - \mu b : \mu \in \mathbb{R}^m \right\}$$

We can generalize with quadratic functions for general problems:

Strictly convex QP: (P)

$$\min \left\{ \frac{1}{2} x^T Q x + q x : Ax \geq b \right\}, Q \succ 0 \implies (D) \max \left\{ \lambda b - \frac{1}{2} v^T Q^{-1} v : \lambda A - v = q, \lambda \geq 0 \right\}$$

strong duality  $\equiv v(P) = v(D)$  (almost) always holds

### 22.3.3 Conic program

We can do duals of things that are not quadratic programs. Sometimes we want to have non linear things to be able to draw different shapes.

**Definition 22.4** (Conic program). We term **conic program**  $\min\{cx : Ax \geq_K b\}$ , where  $x \geq_K y \equiv x - y \in K$ , where  $K$  is a convex cone.

**Example 22.2.** It is easy to see that the  $\geq$  constraints we saw before are a special case of conic program, where the conic is  $\mathbb{R}_+^n$ .

$$-x_1 - 2x_2 \leq 2$$

$$x_1 + x_2 \geq 1$$

$$2x_1 - x_2 \geq 0$$

Let us write  $-x_1 - 2x_2 - 2 = s_0$ ,  $x_1 + x_2 - 1 = s_1$  and  $2x_1 - x_2 = s_2$ , such that  $s_0, s_1, s_2 \geq 0$ .

We have a mapping from  $\mathbb{R}^n$  (where the variables live) to  $\mathbb{R}^m$  (where the constraints live) and  $Ax - b \in K$ .

Rather than writing  $s_0, s_1, s_2 \geq 0$  we could write  $s_0 \geq \sqrt{s_1^2 + s_2^2}$  and this would still be a conic program.




FIGURE 22.1:  $x^3 \geq \sqrt{x_1^2 + x_2^2}$  is a convex cone.

The idea is to hide the non linear part in the cone, in particular in  $\geq_K$ . Let us see what happens to cones depending on the function.

There are three interesting cones:

- $K = \mathbb{R}_+^n \equiv$  sign constraints  $\equiv$  Linear Program;
- $K = \mathbb{L} = \{x \in \mathbb{R}^n : x_n \geq \sqrt{\sum_{i=1}^{n-1} x_i^2}\} \equiv$  Second-Order Cone (or Lorentz cone) Program, that generalizes linear programs;
- $K = \mathbb{S}_+ = \{A \succeq 0\} \equiv \succeq$  constraints  $\equiv$  SemiDefinite Program.

Given the problem that looks like linear except for the cone.

**Definition 22.5** (Conic dual). *Conic dual: (D)  $\max\{yb : yA = c, y \geq_{K^D} 0\}$ , where  $K^D = \{z : \langle z, x \rangle \geq 0 \ \forall x \in K\}$  is a dual cone.*

Sometimes constraints qualification does not hold, since the conic program is not linear sometimes.

Explicit form of second order cone program (SOCP) (“explicit data”  $D_i, d_i, p_i, q_i$ )  $\min\{cx : \|D_i x - d_i\|_2 \leq p_i x - q_i \ i = 1, \dots, m\}$ .

It collapses to a linear program for  $D_i = 0$  and  $d_i = 0$ .

It turns out that the dual has just this explicit form:

$$\max\left\{\sum_{i=1}^m \lambda_i d_i + \nu_i q_i : \sum_{i=1}^m \lambda_i D_i + \nu_i p_i = c, \|\lambda_i\|_2 \leq \nu_i, i = 1, \dots, m\right\}$$

where the  $\nu_i$  are the dual variables of the rightmost part, while the  $\lambda_i$  are the dual variables of the leftmost part.

We can write the explicit form of semidefinite problems as  $\min\{cx : \sum_{i=1}^n x_i A^i \succeq B\}$ , where  $A^i, B \in M(k, \mathbb{R})$ .

There is an even more general form of duality, that we are going to introduce next.

## 22.4 Fenchel's duality

**Definition 22.6** (Fenchel's conjugate). We denote **Fenchel's conjugate** of  $f$   $f^*(z) = \sup_x \{zx - f(x)\}$ .

We can observe that  $f^*$  is always convex, even if  $f$  is not and it is closed if  $f$  is.

The following functions are such that  $f^*$  can be computed easily:

1.  $f(x) = \frac{1}{2} \|x\|_2^2 \implies f^*(z) = \frac{1}{2} \|z\|_2^2$  (only function s.t.  $f^* = f$ );
2.  $(\|\cdot\|_1)^*(z) = \beta_{\mathcal{B}_\infty(0,1)}(z)$ ,  $(\|\cdot\|_\infty)^*(z) = \beta_{\mathcal{B}_1(0,1)}(z)$ ;
3.  $f(x) = \max\{g_i x - \alpha_i \mid i \in I\} \implies f^*(z) = \min \{ \sum_{i \in I} \alpha_i \theta_i \mid \sum_{i \in I} g_i \theta_i = z, \sum_{i \in I} \theta_i = 1, \theta_i \geq 0 \mid i \in I \}$ .

**Fact 22.6.** If we are minimizing the sum of two convex functions  $(P) \min\{f(x) + g(x)\}$ , this problem is the same of maximizing the sum of “almost” the two conjugates:  $(D) - \min\{f^*(z) + g^*(-z)\}$ .

## 23 12th of December 2018 — A. Frangioni

### 23.1 Quadratic problem with linear equality constraints

Let  $A$  be a matrix in  $M(m, n, \mathbb{R})$ , where  $m < n$  (otherwise the system is either fully determined or impossible) and  $\text{rk}(A) = m$ . The constrained quadratic problem may be written as

$$\min \left\{ \frac{1}{2}x^T Qx + qx : Ax = b \right\}$$

where  $Q \succeq 0$ .

A way to solve this problem is through Karush Kuhn Tucker system:

$$\begin{array}{ll} \text{(a)} & \left[ \begin{array}{cc} Q & A^T \\ A & 0 \end{array} \right] \left[ \begin{array}{c} x \\ \mu \end{array} \right] = \left[ \begin{array}{c} -q \\ b \end{array} \right] \\ \text{(b)} & \end{array} \quad (23.1)$$

where the first row (a) says that the gradient is a linear combination of the normals of the gradient of the constraints and the second one is just feasibility.

Everything is linear here. This system is symmetric, although indefinite, because it contains many 0s.

We are left with solving the KKT system:

**REDUCED KKT:** in this method we first add the hypothesis of non-singularity to the matrix  $Q$  so we get the following

$$\begin{cases} Qx + A^T\mu = -q & \text{(a)} \\ Ax = b & \text{(b)} \end{cases}$$

Multiply (a) by  $AQ^{-1}$

$$\begin{cases} AQ^{-1}Qx + AQ^{-1}A^T\mu = -AQ^{-1}q & \text{(a)} \\ Ax = b & \text{(b)} \end{cases}$$

Multiply (b) by  $-1$  and add to it (a)

$$\begin{cases} Ax + AQ^{-1}A^T\mu = -AQ^{-1}q & \text{(a)} \\ Ax + AQ^{-1}A^T\mu - Ax = -AQ^{-1}q - b & \text{(b)} \end{cases}$$

Multiply (a) by  $A^{-1}$

$$\begin{cases} x + Q^{-1}A^T\mu = -Q^{-1}q & \text{(a)} \\ AQ^{-1}A^T\mu = -AQ^{-1}q - b & \text{(b)} \end{cases}$$

Isolate  $x$

$$\begin{cases} x = -Q^{-1}(A^T \mu + q) & \text{(a)} \\ AQ^{-1}A^T \mu = -AQ^{-1}q - b & \text{(b)} \end{cases}$$

Notice that  $0 \preceq AQ^{-1}A^T = M \in M(m, \mathbb{R})$  and may be much smaller than the original one, since its size depends on the number of constraints. The issue here is that the matrix  $M$  is less sparse than both  $A$  and  $Q$ .

**NULL SPACE METHOD:** In this method we need no assumption on  $Q$ .

First of all we rearrange the matrix  $A$  in order to have a small square matrix  $A_B$  and then the rest of the columns ( $A_N$ ):  $A = [A_B, A_N]$ ,  $x = [x_B, x_N]$ ,  $\det(A_B) \neq 0$ .

Replacing  $A$  in (b) we get  $A_B x_B + A_N x_N = b \iff x_B = A_B^{-1} \cdot (b - A_N x_N)$ , hence resulting in  $x = Dx_N + d$ , where

$$d = \begin{bmatrix} b \\ 0 \end{bmatrix}, D = \begin{bmatrix} -A_B^{-1} A_N \\ I \end{bmatrix} \in M(m, n-m)$$

Notice that  $D$  is a basis of the **null space** (or kernel) of  $A$  and it is not mandatory to build it like we did above, it is only important to obtain a basis of the kernel.

Let us multiply (a) by  $D^T$  and obtain

$$\begin{aligned} D^T(Qx + A^T \mu) &= -D^T q \\ D^T Qx + D^T A^T \mu &= -D^T q \\ D^T Qx + (AD)^T \mu &= -D^T q \\ D^T Q(Dx_N + d) &= -D^T q \end{aligned} \tag{23.2}$$

Where in the last step we applied the definition  $x = Dx_N + d$  and hence,  $(D^T Q D)x_N = -D^T(Qd + q)$ .

We term  $H = D^T Q D \in M(n-m, \mathbb{R})$  the reduced Hessian of the problem and notice that whenever the number of constraints is close to the number of variables this matrix is very small.

It is important to note that in order to solve an equality constrained problem we can choose either the reduced KKT method or the null space method, depending on the structure of our problem.

## 23.2 Inequality constrained problems

The problem, in this case, is written as follows

$$(P) \min\{f(x) : Ax \leq b\}$$

### 23.2.1 Projected gradient method

The intuition behind this kind of algorithm is that we are interested in finding a direction for the line search not the opposite of the gradient (because then the step size should be put to 0 if we lie on the boundaries, see Figure 23.1), but it is enough to pick a direction which has a negative scalar product with the gradient.

The rationale is to pick the direction that minimizes the norm of the difference with the gradient. Formally




FIGURE 23.1: Geometric idea of how to choose the direction in case of lying on the boundary.

$$d = \operatorname{argmin}\{\|\nabla f(x) - d\|^2 : d \in \mathcal{D}_X(x)\}$$

where  $\mathcal{D}_X(x)$  is the feasible cone. Notice that if the difference between the gradient and the direction is 0 it means that we can stop, since the descent direction would bring us outside the feasible set.

A more formal definition of the feasible cone  $\mathcal{D}_X(x) = \{d \in \mathbb{R}^n : A_{\mathcal{A}(x)}d \leq 0\}$ , where  $\mathcal{A}(x)$  is the set of all the active constraints.

For sake of clarity we denote  $\bar{A} = A_{\mathcal{A}(x)}$ .

At this point we are ready to project the problem in such a way that inequality constraints become equality constraints:

$$\min \left\{ \frac{1}{2} \|\nabla f(x) + d\|^2 = \frac{1}{2} d^T Id + \nabla f(x)d : \bar{A}d = 0 \right\}$$

Note that in this case the Hessian is the identity matrix  $I \succ 0$ , hence the KKT conditions become simpler:

$$\begin{cases} Qx + A^T \mu = -q \iff d + A^T \mu = -\nabla f(x) \\ Ad = 0 \end{cases}$$

since  $x = d, A = I, b = \nabla f(x)d$ , so we get

$$\begin{aligned} d &= -\nabla f(x) - A^T \mu \\ A\bar{d} &= -A\nabla f(x) - AA^T \mu \\ AA^T \mu &= -A\nabla f(x) \end{aligned} \tag{23.3}$$

resulting in

$$\begin{cases} Qx + A^T \mu = -q \iff d + A^T \mu = -\nabla f(x) \\ Ad = 0 \end{cases}$$

Therefore, we need to solve a system in  $\mu$  and then we have the direction. If the number of active constraints is small, solving the system is very fast.

**Fact 23.1.** *We can restrict to solve*

$$\begin{cases} \mu = -(\bar{A}\bar{A}^T)^{-1}\bar{A}\nabla f(x) \\ d = (I - \bar{A}^T(\bar{A}\bar{A}^T)^{-1}\bar{A})(-\nabla f(x)) \end{cases}$$

*Proof.*  $\bar{A}$  has full row rank, then is non singular and may be inverted.

$$\begin{cases} \mu = -(\bar{A}\bar{A}^T)^{-1}\bar{A}\nabla f(x) \\ d = -\nabla f(x) + \bar{A}^T(\bar{A}\bar{A}^T)^{-1}\bar{A}\nabla f(x) \end{cases} \iff \begin{cases} \mu = -(\bar{A}\bar{A}^T)^{-1}\bar{A}\nabla f(x) \\ d = (I - \bar{A}^T(\bar{A}\bar{A}^T)^{-1}\bar{A})(-\nabla f(x)) \end{cases}$$

□

If the set of active constraints contains some linear dependent ones, we take the maximal subset of our constraints such that the matrix  $\bar{A}$  has full rank.

This procedure is formalized in Algorithm 23.1, where at each stage we remove one of the constraints that lead to a negative  $\mu$ , keeping a set of linearly independent constraints.

In the 14-th line of the algorithm we find that our step size is takes as the minimum step size among the ones that satisfy some subsets of the constraints.

---

ALGORITHM 23.1 Pseudocode for projected gradient method for quadratic functions.

---

```

1: procedure PGM( $f, A, b, x, \varepsilon$ )
2:   for ( $; ;$ ) do
3:      $B \leftarrow$  maximal  $\subseteq \mathcal{A}(x)$  s.t.  $\text{rank}(A_B) = |B|$ ;
4:     for ( $; ;$ ) do
5:        $d \leftarrow (I - A_B^T (A_B A_B^T)^{-1} A_B)(-\nabla f(x))$ ;
6:       if  $\langle \nabla f(x), d \rangle \leq \varepsilon$  then
7:          $\mu_B \leftarrow -(A_B A_B^T)^{-1} A_B \nabla f(x)$ ;
8:          $\mu_i \leftarrow 0 \forall i \notin B$ ;
9:         if  $\mu_B \geq 0$  then return
10:        end if
11:         $h \leftarrow \min\{i \in B : \mu_i < 0\}$ ;
12:      end if
13:       $\bar{\alpha} \leftarrow \min\{\alpha_i = (b_i - A_i x)/A_i d : A_i d > 0, i \notin B\}$ ;
14:       $x \leftarrow x + \alpha d$ ;
15:      if  $\bar{\alpha} > 0$  then
16:        break;
17:      end if
18:       $k \leftarrow \min\{i \notin B : A_i d > 0 : \alpha_i = 0\}$ ;
19:       $B \leftarrow B \cup \{k\}$ ;
20:    end for
21:     $\alpha \leftarrow \text{Line\_Search}(f, x, d, \bar{\alpha})$ ;
22:     $x \leftarrow x + \alpha d$ ;
23:  end for
24: end procedure

```

---

**Fact 23.2.** *The following holds:*

1.  $d = 0 \wedge \mu \geq 0 \implies x$  optimal (from KKT);
2.  $d = 0 \wedge \exists h \in B$  s.t.  $\mu_h < 0 \implies \exists x' \in \{x \in \mathbb{R}^n : A_{B \setminus \{h\}}x = b_{B \setminus \{h\}}, A_h x \leq b_h\}$  s.t.  $f(x') < f(x)$ ;
3.  $d \neq 0$  descent direction:  $H = I - A_B^T [A_B A_B^T]^{-1} A_B$  symmetric and idempotent  $HH = H^T H = H \implies \langle d, \nabla f(x) \rangle < 0$ .

*Proof.*

trivial;

- The intuition is that if we remove  $h$  from  $B$  next time we will have a descent direction.  $\exists d$  s.t.  $A_{B \setminus \{h\}}d = 0 \wedge A_h d < 0 \implies \langle \nabla f(x), d \rangle = \langle -\mu A_B, d \rangle = -\mu_h A_h d < 0$ ;

- $\langle d, \nabla f(x) \rangle = \langle -H\nabla f(x), \nabla f(x) \rangle = -\nabla f(x)^T H \nabla f(x) = -(H \nabla f(x))^T H \nabla f(x)$ .

□

Once we found the optimal face we move inside that face and we are dealing with a steepest descent, which is slow.

At a certain point of the execution the set  $B$  of the active constraints will stabilize and become the one of the optimal solution.

At this point the smart thing to do is to use the KKT conditions, since we know the set of active constraints. This idea is pursued in Section 23.2.3.

### 23.2.2 Projected gradient method with box constraints

Given the non linear minimum problem of the following shape

$$(P) \min \left\{ f(x) : l \leq x \leq u \right\}$$

a slightly different version of the projected gradient method forces the algorithm to put to 0 the component of the direction which would bring us to go outside the feasible region. This intuition is formalized in Algorithm 23.2.

---

ALGORITHM 23.2 Pseudocode for projected gradient method for quadratic functions in the case of box constraints.

---

```

1: procedure PGMBC( $f, l, u, x, \varepsilon$ )
2:    $d = -\nabla f(x)$ ;
3:    $\bar{\alpha} = \infty$ ;
4:   for ( $i = 1 \dots n$  s.t.  $d_i \neq 0$ ) do
5:     if ( $d_i < 0$ ) then
6:       if ( $x_i = l_i$ ) then
7:          $d_i = 0$ ;
8:       else
9:          $\bar{\alpha} \leftarrow \min\{\bar{\alpha}(x_i - l_i)/d_i\}$ ;
10:      end if
11:    else
12:      if ( $x_i = u_i$ ) then
13:         $d_i = 0$ ;
14:      else
15:         $\bar{\alpha} \leftarrow \min\{\bar{\alpha}(u_i - x_i)/d_i\}$ ;
16:      end if
17:    end if
18:    if ( $\langle \nabla f(x), d \rangle \leq \varepsilon$ ) then return
19:    end if
20:     $\alpha \leftarrow \text{Line\_Search}(f, x, d, \bar{\alpha})$ ;
21:     $x \leftarrow x + \alpha d$ ;
22:  end for
23: end procedure

```

---

Notice that we can assume that  $l_i < u_i \forall i$ , because otherwise that component would be fixed.

### 23.2.3 Active-set method for quadratic programs

Let us be given the following minimum problem

$$\min \left\{ \frac{1}{2} x^T Q x + q x : Ax \leq b \right\}$$

where an important hypothesis is that the problem is quadratic, once we know  $\mathcal{A}(x_*)$ .

We start from a certain point, such that the constraints are satisfied as equalities. According to KKT conditions, a solution  $\bar{x}$  is optimal if  $\bar{x}$  is feasible and  $\mu \geq 0$ . Otherwise, if  $\mu$  is not positive we eliminate something from the active set and start again. In the case of  $x$  unfeasible, we know the descent direction, we only need to revise the step size.

All this machinery is formalized in Algorithm 23.3.

---

ALGORITHM 23.3 Pseudocode for active set method for quadratic functions.

---

```
1: procedure ASMQP( $Q, q, A, b, x, \varepsilon$ )
2:   for ( $B \leftarrow \mathcal{A}(x); ;$ ) do
3:     solve ( $P_B$ )  $\min\{\frac{1}{2}x^T Qx + qx : Ax = b\}$  for ( $\bar{x}, \bar{\mu}_B$ );
4:     if ( $A_i \bar{x} \leq b_i \forall i \notin B$ ) then
5:       if ( $\bar{\mu}_B \geq 0$ ) then return
6:       end if
7:        $h \leftarrow \min\{i \in B : \mu_i < 0\};$ 
8:        $B \leftarrow B \setminus \{h\};$ 
9:       continue;
10:      end if
11:       $d \leftarrow \bar{x} - x;$ 
12:       $\bar{\alpha} \leftarrow \min\{\alpha_i = (b_i - A_i x) / A_i d : A_i d > 0, i \notin B\};$ 
13:       $x \leftarrow x + \bar{\alpha}d;$ 
14:       $B \leftarrow \mathcal{A}(x);$ 
15:    end for
16: end procedure
```

---

Notice that this algorithm allows easy solving of the same problem, under box constraints, because we end up having two sets:  $L$ , indexes of variables fixed to the lower bound, and  $U$ , indexes of variables fixed to the upper bound.

$\{1, \dots, n\} \setminus (L \cup U)$  is the set of the indexes of the free variables.

Thanks to this consideration the problem to solve becomes

$$\min \left\{ \frac{1}{2}x_F^T Q_{FF}x_F + (q_F + u_U^T Q_{UF})x_F \right\} [ + \frac{1}{2}x_U^T Q_{UU}x_U + q_U u_U ]$$

#### 23.2.4 Frank-Wolfe method

Let us take a non-linear function and the following problem

Supposing  $f$  is convex we can make a first order model and minimize it over our constraints. In this case we use the right constraints and an approximation of the function.

---

ALGORITHM 23.4 Pseudocode for Frank-Wolf method for non linear functions.

---

```
1: procedure FWM( $f, A, b, x, \varepsilon$ )
2:   while ( $\|\nabla f(x)\| > \varepsilon$ ) do
3:      $\bar{x} \leftarrow \text{argmin } \{\langle \nabla f(x), y \rangle : Ay \leq b\};$ 
4:      $d \leftarrow \bar{x} - x;$ 
5:      $\alpha \leftarrow \text{Line\_Search}(f, x, d, 1);$ 
6:      $x \leftarrow x + \alpha d;$ 
7:   end while
8: end procedure
```

---

Provided that the computation at line 3 of Algorithm 23.4 is performed efficiently, then either  $d = 0$  and we are in the optimum or  $d > 0$  and we are moving toward the optimum.

The linear model is below the function in the quadratic case, hence we get a lower bound.

## 24 14th of December 2018 — A. Frangioni

In the previous lecture we addressed the problem of linear constrained optimization. Our first approach was to deal very little with constraints (projected gradient method), after a few improvements we took all of them and modify the function (Frank-Wolfe method).

### 24.1 Dual methods for linear constrained optimization

In this class of methods constraints are first class citizens.

Let us be given the following optimization problem:

$$\min \left\{ \frac{1}{2}x^T Qx + qx : Ax \leq b \right\}$$

We would like to work with dual feasibility.

$\forall$  fixed  $\lambda \geq 0$ , this is the lagrangian problem  $\psi(\lambda) = \min_x \left\{ \frac{1}{2}x^T Qx + qx + \lambda(b - Ax) \right\} \leq v$  and the Lagrangian dual is the following

$$(D) \max \{\psi(\lambda) : \lambda \geq 0\}$$

which is equivalent to the primal problem.

The solution of the Lagrangian problem is **unique** and this is due to the fact that  $\psi$  is concave and  $Q \succ 0$ . The optimal solution is then  $x(\lambda) = Q^{-1}(\lambda A - q)$ .

The function is differentiable in the solution, because we have only one sub (or super) gradient which is  $\nabla \psi(\lambda) = b - Ax(\lambda)$ .

At this point we only need to solve the dual problem, which has the shape of a box constrained optimization, where the box has only one boundary and this can be solved via quasi-Newton methods.

Notice that  $\psi \notin C^2$  so the Hessian is not defined.

This dual approach is advantageous in the case of a small number of constraints, because the size of the problem decreases. For example, in the case of one constraint, the Lagrangian dual becomes a problem in one variable, hence solvable through a line search.

In this method the degenerate case (more than one constraint active at a time) is not an issue.

If the quadratic function is convex we may use quasi Newton method. Otherwise the global optimality is not guaranteed and may be used if we accept not to be able to solve the original problem, but only to find a lower bound.

#### 24.1.1 Separable problems and partial dual

Let us assume that our constraints are separable, which means that it is not mandatory to work with all of them, but they can be splitted into constraints of groups of variables.

$$\min \{f(x) : Ax \leq b, Ex \leq d\}$$

We can decide to use a **partial dual**, writing the Lagrangian problem picking only some constraints that we chose:

$$\psi(\lambda) = \min_x \{f(x) + \lambda(b - Ax) : Ex \leq d\}$$

The Lagrangian dual method may be better than projected gradient or worse and it depends on the instance.

In the dual approaches we can't move inside the feasible solution. We find an optimum for the dual, which surely breaks feasibility. Then, if the variable is above the upperbound it gets decreased to the upper bound, otherwise if it is below the lower bound it takes the value of the lower bound.

This way we get an upperbound for the function to be minimized.

## 24.2 Primal/dual methods or barrier methods

This kind of methods are designed to overcome the cons of dual approaches, namely the fact that  $\psi$  does not have the Hessian (and this creates problems to quasi Newton method) and the fact that  $x$  is not feasible until the end.

At the same time this methods keep the unconstrained property of the Lagrangian dual.

### 24.2.1 Barrier function and central path

The rationale behind this algorithm is to minimize a function which penalizes the value of the original function when the solution is getting closer and closer to the boundaries of the feasible set.

$$(P_\mu) \min\{f_\mu(x) = f(x) - \mu \sum_{i=1}^m \log(b_i - A_i x)\}$$

The parameter  $\mu$  is there to weight the proximity to the boundary.

**Property 24.1.** • If  $f$  is convex,  $f_\mu$  is strictly convex;

- If  $f \in \mathcal{C}^2$  then  $f_\mu \in \mathcal{C}^2$ , since  $\log \in \mathcal{C}^\infty$ ;
- $\forall \mu \exists! x_\mu$  optimal of  $(P_\mu)$ , since  $\mu \sum_{i=1}^m \log(b_i - A_i x)$  is strictly convex;
- As  $\mu \rightarrow 0$   $x_\mu$  converges to the analytic center of the optimal face. An example of this behaviour may be seen in Figure 24.1.

Another interesting property is that, since the barrier function is **self concordant**  $x^i$  gets “close” to  $x(\mu^i)$  in very few Newton's steps.

In one step  $x^{i+1}$  is “much closer” to  $x(\mu^i)$  than  $x^i$  was and  $x^{i+1}$  is “close” to  $x(\mu^{i+1})$ , with  $\mu^{i+1} \ll \mu^i$  (more formally,  $\mu^{i+1} = \tau \mu^i$ ,  $\tau < 1$ ).

This behaviour may be observed in Figure 24.2




FIGURE 24.1: The trajectory converges to the optimal solution of the problem, when  $\mu \rightarrow 0$ .




FIGURE 24.2: The dotted line represents a region where the Newton method is very efficient. We are starting from a point  $x_1$ , which belongs to that region and we want to move towards  $x_{\mu^1}$ . At next iterate  $x_2$  is closer to  $x_{\mu^2}$  than the current iterate.

The convergence is exponential if  $\tau$  is very small, but these iterations are very costly, because the Hessian changes at each step, hence it needs to be recomputed.

Let us focus on computing the **Newton's step**.

First, we write the Karusich-Kuhn-Tucker conditions:

PRIMAL FEASIBILITY:  $Ax + s = b, s \geq 0$ ;

DUAL FEASIBILITY:  $Qx + \lambda A = -q, \lambda \geq 0$ ;

COMPLEMENTARY SLACKNESS:  $\lambda_i s_i = 0, i = 1, \dots, m$ .

We can write a slackened version of KKT, imposing  $\lambda_i s_i = \mu, i = 1, \dots, m$ , where  $\mu \in \mathbb{R}^n$  and should be decreased over iterations until it gets closer enough to 0.

Let us construct  $\Lambda, S \in \text{Diag}(m, \mathbb{R})$  such that the diagonal is made of  $\lambda_i$  and  $s_i$  respectively.

At this point, we rewrite the problem in terms of the displacement from the fixed current point we are in:

- $x \rightarrow x + \Delta x$
- $s \rightarrow s + \Delta s$
- $\lambda \rightarrow \lambda + \Delta \lambda$

The KKT system becomes:

$$\text{PRIMAL FEASIBILITY: } Ax + A\Delta x + s + \Delta s = b, \quad s \geq 0;$$

$$\text{DUAL FEASIBILITY: } Qx + Q\Delta x + \lambda A + \Delta \lambda A = -q, \quad \lambda \geq 0;$$

$$\text{COMPLEMENTARY SLACKNESS: } \lambda_i s_i + \lambda_i \Delta s_i + s_i \Delta \lambda_i + \Delta \lambda_i \Delta s_i = \mu, \quad i = 1, \dots, m.$$

In this new system of coordinates the first two KKT remain linear, while the third one is no longer linear ( $\Delta \lambda_i \Delta s_i$ ).

$$\begin{bmatrix} Q & A^T & 0 \\ A & 0 & I \\ 0 & S & \Lambda \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta \lambda \\ \Delta s \end{bmatrix} \stackrel{(1)}{=} \begin{bmatrix} -(Qx + q) - \lambda A \\ b - Ax - s \\ \mu u - \Lambda S u - \Delta \Lambda \Delta S u \end{bmatrix} \approx \begin{bmatrix} 0 \\ 0 \\ \mu u - \Lambda S u \end{bmatrix} \quad (24.1)$$

Where (1) holds since  $\Delta \lambda A = A^T \Delta \lambda^T$ , although  $\Delta \lambda$  is written without the “transpose” syntax to ease notation.

Notice that  $u = \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix} \in \mathbb{R}^m$  and has the purpose of adjusting dimension:

$$S \Delta \lambda = \begin{bmatrix} s_1 \Delta \lambda_1 \\ \vdots \\ s_m \Delta \lambda_m \end{bmatrix} \in \mathbb{R}^m; \quad \Lambda \Delta s = \begin{bmatrix} \lambda_1 \Delta s_1 \\ \vdots \\ \lambda_m \Delta s_m \end{bmatrix} \in \mathbb{R}^m; \quad \mu u = \begin{bmatrix} \mu \\ \vdots \\ \mu \end{bmatrix} \in \mathbb{R}^m;$$

$$\Lambda S u = \begin{bmatrix} -s_1 \lambda_1 \\ \vdots \\ -s_m \lambda_m \end{bmatrix} \in \mathbb{R}^m; \quad \Delta \Lambda \Delta S u = \begin{bmatrix} -\Delta \lambda_1 \Delta s_1 \\ \vdots \\ -\Delta \lambda_m \Delta s_m \end{bmatrix} \in \mathbb{R}^m;$$

The Newton method can be applied if we discard the non-linear part (written in red), pretending it does not exist. Notice that vector  $u$  is the vector of all 1s.

### 24.3 Primal-dual interior point method

This method is based on the observation that we can solve the dual problem

$$(D) \quad \max \left\{ -\lambda b - \frac{1}{2} x^T Q x : Qx + \lambda A = -q, \lambda \geq 0 \right\}$$

thus obtaining both a lower and upper bound for the solution  $x$ .

We term **complementarity gap**  $(\frac{1}{2} x^T Q x + q x) - (-\lambda b - \frac{1}{2} x^T Q x) = \lambda s = \mu$ .

Once we found a solution for Equation (24.1), we perform a step and compute a new couple of primal and dual solutions and reduce the gap  $\mu$ .

We are left with solving Equation (24.1). The trick is to express one between  $\Delta s$  and  $\Delta \lambda$  as a linear combination of the other.

For example, let us take the third line of Equation (24.1):

$$\begin{aligned}
0\Delta x + S\Delta\lambda + \Lambda\Delta s &= \mu u - \Lambda S u \\
\Lambda\Delta s &= \mu u - \Lambda S u - S\Delta\lambda \\
\Delta s &= \Lambda^{-1}\mu u - A^T \cancel{\Lambda} S u - \Lambda^{-1} S \Delta\lambda \\
\Delta s &= \Lambda^{-1}\mu u - S u - \Lambda^{-1} S \Delta\lambda \\
\Delta s &= \Lambda^{-1} \cdot (\mu u - S\Delta\lambda) - S u \\
\Delta s &= \Lambda^{-1} \cdot (\mu u - S\Delta\lambda) - s
\end{aligned} \tag{24.2}$$

We obtain the modified normal equations (or KKT system)

$$\begin{bmatrix} Q & A^T \\ A & -\Lambda^{-1}S \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta\lambda \end{bmatrix} = \begin{bmatrix} 0 \\ s - \mu\Lambda^{-1}u \end{bmatrix} \tag{24.3}$$

where the last row is derived working on the second row of Equation (24.1) ( $A\Delta x + 0\Delta\lambda + I\Delta s = 0 \Leftrightarrow A\Delta x = -\Delta s$ ), substituting Equation (24.2):

$$\begin{aligned}
A\Delta x - [\Lambda^{-1}S]\Delta\lambda &= -\Delta s - [\Lambda^{-1}S]\Delta\lambda \\
&= -\Lambda^{-1}(\mu u - S\Delta\lambda) + s - \Lambda^{-1}S\Delta\lambda \\
&= -\Lambda^{-1}\mu u + \cancel{\Lambda^{-1}S\Delta\lambda} + s - \cancel{\Lambda^{-1}S\Delta\lambda} \\
&= s - \mu\Lambda^{-1}u
\end{aligned} \tag{24.4}$$

With respect to normal equations of ??, we have in position (2, 2) a quantity  $(-\Lambda^{-1}S)$ , which is not 0, but it is the opposite of a strictly positive definite matrix.

A possible approach for solving this system is making some calculations and obtain something of the shape of reduced KKT (see Section 23.1).

$$\begin{cases} Q\Delta x + A^T\Delta\lambda = 0 \\ (Q + A^T\Lambda S^{-1}A)\Delta x = A^T(\lambda - \mu S^{-1}u) \end{cases} \tag{24.5}$$

where the first set of equations follows from the expansion of the first row of the KKT system (Equation (24.3)) and the second one is obtained taking the same row of the same system and substituting the value of  $\Delta\lambda$  as follows:

$$\begin{aligned}
A\Delta x - \Lambda^{-1}S\Delta\lambda &= s - \mu\Lambda^{-1}u \\
\Lambda^{-1}S\Delta\lambda &= A\Delta x - s + \mu\Lambda^{-1}u \\
\Delta\lambda &= (\Lambda^{-1}S)^{-1}A\Delta x - (\Lambda^{-1}S)^{-1}s + (\Lambda^{-1}S)^{-1}\mu\Lambda^{-1}u \\
\Delta\lambda &= S^{-1}\Lambda A\Delta x - S^{-1}\Lambda s + \mu S^{-1} \cancel{\Lambda} \cancel{A} u \\
\Delta\lambda &= S^{-1}\Lambda A\Delta x - S^{-1}\Lambda s + \mu S^{-1}u \\
\Delta\lambda &= \mu S^{-1}u + \Lambda S^{-1}A\Delta x - \Lambda S^{-1}s \\
\Delta\lambda &= \mu S^{-1}u + \Lambda S^{-1}A\Delta x - \Lambda u \\
\Delta\lambda &= \mu S^{-1}u + \Lambda S^{-1}A\Delta x - \lambda
\end{aligned} \tag{24.6}$$

Hence,

$$\begin{aligned}
Q\Delta x + A^T \Delta \lambda &= 0 \\
Q\Delta x + A^T \cdot (\mu S^{-1}u + \Lambda S^{-1}A\Delta x - \lambda) &= 0 \\
Q\Delta x + A^T \mu S^{-1}u + A^T \Lambda S^{-1}A\Delta x - A^T \lambda &= 0 \\
Q\Delta x + A^T \Lambda S^{-1}A\Delta x &= -A^T \mu S^{-1}u + A^T \lambda \\
(Q + A^T \Lambda S^{-1}A)\Delta x &= A^T(\lambda - \mu S^{-1}u)
\end{aligned} \tag{24.7}$$

We term  $M = Q + A^T \Lambda S^{-1}A$  and the following holds.

**Fact 24.2.** *If  $A$  has full column rank (aka it is invertible),  $M \succ 0$ .*

At this point we need to factorize the matrix  $M$ , that changes at each iteration (since  $\Lambda S^{-1}$  does) and this is the bottleneck.

Cholesky factorization may be used, although its complexity is cube. Another downside of this approach is that the matrix  $M$  is much denser than  $A$ ,  $\Lambda$  and  $S^{-1}$ .

An orthogonal approach to the reduced KKT is called **predictor-corrector** and it works computing a solution without taking into account the non linear term  $\Delta \Lambda \Delta S u$ , then computing it according to the approximated solution and repeat until convergence.

The bottleneck again is solving the system in Equation (24.1).

For what concerns implementation, we should start from a triplet  $(x, \lambda, s)$ , that could be not feasible and then compute the residuals and iterate until feasibility is reached.

$$r^D = -(Qx + q) - \lambda A, r^P = b - Ax - s.$$

When dealing with the step size we need to highlight the fact that  $\lambda + \Delta \lambda \geq 0$ ,  $s + \Delta s \geq 0$  should hold.

In order to achieve this we find the maximum  $\alpha$  that satisfies the equality and then multiply it by a constant  $\bar{\alpha} = 0.995$  (or 0.9995), in order to get closer.

Let us assume that we also have a bunch of box constraints, hence our problem becomes

$$(P) \quad \min \left\{ \frac{1}{2} x^T Q x + q x : Ax = b, 0 \leq x \leq u \right\}$$

In this special case, things simplify a lot.