COMP0147_18-19

Sheet 4

Due Monday 20 February. Hand-in-solutions to questions 2, 4, 5, 6 91.23 u93.43=91, 21 3143. Eq. class of $\{1,2,3,4,5\}$ and $Y=\{3,4\}$. Define a relation \mathcal{R} on the power set P(X)PRoflexivity: ARA > since AUY: AUY. =[51,23, 51,2,3], 51,2,4] of X by $A\mathcal{R}B \text{ iff } A \cup Y = B \cup Y$ Symmetry ivalence relation. (۱، ۲، ۲، ۲) (a) Prove that R is an equivalence relation. (b) What is the equivalence class of $\{1, 2\}$? 2. Let X be $\mathbb{Z} \times \mathbb{Z}$, i.e. X is the set of all ordered pairs of the form (x, y) with $x, y \in \mathbb{Z}$. Define the relation R on X as follows: rewritten as Is it an equivalence relation? $(x_1, x_2)R(y_1, y_2) \quad \text{iff} \quad x_1^2 + x_2^2 = y_1^2 + y_2^2.$ $(x_1, x_2)R(y_1, y_2) \quad (x_1, x_2)R(x_1, x_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(y_1, y_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ $(x_1, x_2)R(x_1, x_2) \Rightarrow x_1^2 + x_2^2 = x_1^2 + x_2^2$ E(A, B)= | OA| = 10B) F(A ! W) => (OA) = 104/3. 3. Let $X = \mathbb{R} \times \mathbb{R}$. Define the relation R on X as follows: $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_1, y_1)R(x_1, y_1)R(x_1, y_1) \quad y_1 - y_1 = 2(x_1 - x_1) \Rightarrow 0 = 0.$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad (x_1, y_1)R(x_1, y_1) \quad \text{if} \quad (x_1, y_1)R(x_1,$ $(x_1, y_1)R(x_2, y_2) \quad \text{iff} \quad y_1 - y_2 = 2(x_1 - x_2).$ $\text{Reflexivity} \quad (x_1, y_1) R(x_1, y_1) \quad y_1 - y_1 = 2(x_1 - x_2).$ (a) Is it an equivalence relation? Symmetry, $(x_1, y_1) R(x_2, y_2) \quad y_1 - y_2 = 2(x_1 - x_2)$ by permutations and hence calculate expressing this first as a permutation and then as a symmetry of P.

5. Let G be a group and let H_1 and H_2 be subgroups of G. Show that $H_1 \cap H_2$ is a subgroup of G.

HIEG hihzeHI (hI) GHZ

HINHZEG. hihzeHI (hI) GHZ

HINHZEG. hihzeHZ (hI) GHINHZ

HINHZEG. hihzeHZ. hihzeHINHZ. ClosureN

hihzeH, Associativity

Linheriet. Associativity

hi, hzeHZ. JE is also included since EeHI

EEH=

6. Let $G = \{1_G, g, h, k\}$ be a group with 4 elements and suppose G is not cyclic. Using Lagrange's Theorem show that g, h and k all have order 2 and write down a table for the group operation. (b) Show that the number of elements in G is either 1 or a prime number. 19 is 8 = neutral element Using Lagrange's Theorem, we know that, 141 divides However, only subgroup H = \$16} [1-11=1. if g*h=1 h=g=2g. X if g*h=9 h=1 X if g*h=h g=1 X if g*h=k only possible.