

## 1. The Wave Function

### 1.1 The Schrödinger Equation

$$i\hbar \frac{\partial \Psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \Psi}{\partial x^2} + V\Psi$$

$$\text{or } i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \vec{\nabla}^2 \Psi(\vec{r}, t) + V(\vec{r}, t) \Psi(\vec{r}, t)$$

$$\text{where } \vec{\nabla}^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$$

Solve for the particle's wave function  $\Psi(x, t)$

$$\hbar = \frac{h}{2\pi} = 1.054572 \times 10^{-34} \text{ Js}$$

### 1.2 The Statistical Interpretation

$$\int_a^b |\Psi(x, t)|^2 dx = \{P \text{ of finding the particle btwn } a \text{ and } b, \text{ at } t\}$$

### 1.3 Probability

$$\text{Standard deviation: } \sigma = \sqrt{\langle j^2 \rangle - \langle j \rangle^2}$$

$$\text{Expectation value of } x \text{ given } \Psi: \langle x \rangle = \int x |\Psi|^2 dx$$

$$\text{Probability current: } J(x, t) = \frac{i\hbar}{2m} (\Psi \frac{\partial \Psi^*}{\partial x} - \Psi^* \frac{\partial \Psi}{\partial x})$$

### 1.4 Normalization

$$\int_{-\infty}^{+\infty} |\Psi(x, t)|^2 dx = 1$$

The Schrödinger equation produces unitary time evolution:

$$\vec{J} = -\frac{i\hbar}{2m} (\psi(\vec{r}, t)^* \vec{\nabla} \psi(\vec{r}, t) - \psi(\vec{r}, t) \vec{\nabla} \psi^*(\vec{r}, t)^*)$$

The probability density satisfies the continuity equation,

$$\frac{\partial}{\partial t} \rho + \vec{\nabla} \cdot \vec{J} = 0$$

Because the probability for finding the particle at infinity is 0

(otherwise non-normalizable),  $\vec{J} = 0$  at infinity.

Therefore,  $\frac{d}{dt} \int_{-\infty}^{\infty} \rho d^3\vec{r} = \frac{d}{dt} P = 0$ , where  $P$  is the total probability  
→ the total probability is constant in time.

### 1.5 Momentum

For a particle in state  $\Phi$ , the expectation value of  $x$  and  $p$  is

$$\langle x \rangle = \int_{-\infty}^{+\infty} x |\Psi(x, t)|^2 dx \quad \langle p \rangle = m \frac{d\langle x \rangle}{dt} = -i\hbar \int (\Psi^* \frac{\partial \Psi}{\partial x}) dx$$

To calculate the expectation value of any quantity,  $Q(x, p)$ :

$$\langle Q(x, p) \rangle = \int \Psi^* Q(x, \frac{\hbar}{i} \frac{\partial}{\partial x}) \Psi dx$$

Position and momentum operators:  $\hat{r} = \vec{r}$ ,  $\hat{p} = -i\hbar \vec{\nabla}$

### 1.6: The Uncertainty Principle

The wavelength of  $\Psi$  is related to the momentum of the particle by the de Broglie formula:

$$p = \frac{h}{\lambda} = \frac{2\pi\hbar}{\lambda}$$

The more precisely determined a particle's position is, the less precisely is its momentum. The Heisenberg's uncertainty principle:

$$\sigma_x \sigma_p \geq \frac{\hbar}{2}$$

Commutation relation btwn position and momentum:

$$\hat{p}_x (\hat{x} \psi(x, t)) = -i\hbar \frac{\partial}{\partial x} [x \psi(x, t)] = -i\hbar \psi(x, t) - i\hbar x \frac{\partial}{\partial x} \psi(x, t)$$

$$\hat{x} (\hat{p}_x \psi(x, t)) = x (-i\hbar \frac{\partial}{\partial x} \psi(x, t))$$

$$\hat{x} \hat{p}_x - \hat{p}_x \hat{x} = [\hat{x}, \hat{p}_x] = i\hbar$$

$$[\hat{x}_i, \hat{p}_j] = i\hbar \delta_{ij}, [\hat{x}_i, \hat{x}_j] = [\hat{p}_i, \hat{p}_j] = 0,$$

where  $\delta_{ij} = 1$  for  $i = j$  and  $\delta_{ij} = 0$  for  $i \neq j$

Given three operators  $\hat{A}, \hat{B}, \hat{C}$ , we have  $[\hat{A}, \hat{B}\hat{C}] = [\hat{A}, \hat{B}]\hat{C} + \hat{B}[\hat{A}, \hat{C}]$ .

### Other: Blackbody Spectrum

$$E = h\nu = \hbar\omega$$

The wave number  $k$  is  $k = 2\pi/\lambda = \omega/c$

Only two spin states occur (quantum number  $m$  is +1 or -1).

$$\rho(\omega) = \frac{\hbar\omega^3}{\pi^2 c^3 (e^{\hbar\omega/k_b T} - 1)}$$

$$\text{Wien displacement law: } \lambda_{\max} = \frac{2.90 \times 10^{-3} \text{ mK}}{T}$$

## 2. Time-Independent Schrödinger Equation

### 2.1 Stationary States

Suppose PE is independent of time,  $V(\vec{r}, t) = V(\vec{r})$ .

Separation of variables:  $\Psi(\vec{r}, t) = \psi(\vec{r})\varphi(t)$

Eq of motion for  $\varphi(t)$ :  $\varphi(t) = e^{-iEt/\hbar}$

Eq of motion for  $\psi(\vec{r})$  is the time-independent Schrödinger equation:

$$-\frac{\hbar^2}{2m} \frac{d^2 \psi(\vec{r})}{dx^2} + V(\vec{r})\psi(\vec{r}) = E\psi(\vec{r})$$

TD of the wavefunction that corresponds to the constant  $E$  is easily written once we solve the TISE:  $\Psi_E(\vec{r}, t) = \psi_E(\vec{r})e^{-iEt/\hbar}$

Properties of solutions for TI potentials:

- **The constant  $E$  must be real.**
- **Stationary wavefunction.**  
 $\mathcal{P}(\vec{r}, t) = |\psi_E(\vec{r}, t)|^2 = |\psi_E(\vec{r})|^2$  (TD cancels out).
- **Stationary wavefunction is a state of definite energy.**  
The total energy (kinetic plus potential) is the Hamiltonian:  
 $H(x, p) = \frac{p^2}{2m} + V(x)$ .  
Hamiltonian operator:  $\hat{H} = -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x)$   
Thus the TISE can be written as  $\hat{H}\psi = E\psi$   
 $\langle \hat{H} \rangle = E$ ,  $\langle \hat{H}^2 \rangle = E^2$ ,  $\Delta E = \sqrt{\langle \hat{H}^2 \rangle - \langle \hat{H} \rangle^2} = 0$
- Spatial part of stationary wavefunction can be chosen to be real.  
 $\psi^*(\vec{r})$  is a soln w/ same  $E$   
Solns can be chosen to be real:  $\psi(\vec{r}) + \psi^*(\vec{r})$  and  $\frac{\psi(\vec{r}) - \psi^*(\vec{r})}{i}$ .
- **Parity symmetry: even and odd wavefunctions.** Suppose  $V(-\vec{r}) = V(\vec{r})$ . Then,  $\psi_E(-\vec{r})$  is a soln w the same energy.  
 $\psi_E(\vec{r}) + \psi_E(-\vec{r})$  is even under reflection,  $\psi_E(\vec{r}) - \psi_E(-\vec{r})$  is odd.  
When the potential is symmetric under reflection, we can choose the stationary states to be either even or odd under reflection.
- **Orthogonality/orthonormality.**  
 $\int \psi_m(\vec{r})^* \psi_n(\vec{r}) d^3\vec{r} = \delta_{mn}$  where  $\delta_{mn}$  is 0 if  $m \neq n$  and 1 if  $m = n$ .
- **Linearity.**

The SE is linear. Given stationary states, a linear combo of these

$$\psi(\vec{r}, t) = \sum_n c_n \psi_n(\vec{r}, t) = \sum_n c_n \psi_n(\vec{r}) e^{-\frac{i}{\hbar} E_n t}$$

where  $c_n$  are complex constants, is a solution to the TDSE

$$i\hbar \frac{\partial \psi(\vec{r}, t)}{\partial t} = \hat{H} \psi(\vec{r}, t)$$

- **Time evolution.** Given

$$\psi(\vec{r}, 0) = \sum_n c_n \psi_n(\vec{r}, 0) = \sum_n c_n \psi_n(\vec{r})$$

at time  $t$ , the time evolution is

$$\psi(\vec{r}, t) = \sum_n c_n \psi_n(\vec{r}, t) = \sum_n c_n \psi_n(\vec{r}) e^{-\frac{i}{\hbar} E_n t}$$

Once we've expanded a given initial wavefunction in terms of a linear combo of the stationary wavefunctions  $\psi_n(\vec{r})$ , the time evolution follows simply by putting a factor of  $e^{-i/h E_n t}$  to each term containing  $\psi_n(\vec{r})$ .

- **Normalization.**  
The constant coefficients are constrained by  $\sum_n |c_n|^2 = 1$
- **Completeness.**

The stationary states form a complete set if

$$\sum_n \psi_n(\vec{r}', t)^* \psi_n(\vec{r}, t) = \delta^3(\vec{r}' - \vec{r})$$

where  $\delta^3(\vec{r}' - \vec{r})$  is the Dirac-delta function in 3D defined by

$$\int d^3\vec{r}' \psi(\vec{r}', t) \delta^3(\vec{r}' - \vec{r}) = \psi(\vec{r}, t)$$

**Euler's formula:**  $e^{i\theta} = \cos \theta + i \sin \theta$

**sin and cos:**  $\cos(\theta) = \frac{1}{2}(e^{i\theta} + e^{-i\theta})$ ,  $\sin(\theta) = \frac{1}{2i}(e^{i\theta} - e^{-i\theta})$

**Delta function:** Given  $f(x)$ ,  $\delta(x - x')$  is defined as

$$f(x') = \int f(x) \delta(x - x') dx$$

$\int \delta(x - x') dx = 1$ , note this is not the area

$$\delta_\alpha(x) = \frac{1}{\alpha\sqrt{\pi}} e^{-\frac{x^2}{\alpha^2}}, \delta_\alpha(x) = \frac{1}{\pi x} \sin\left(\frac{x}{\alpha}\right), \delta_\alpha(x) = \frac{\alpha}{\pi x^2} \sin^2\left(\frac{x}{\alpha}\right)$$

### One-dimensional systems

Wavefunction for a system containing a single particle of mass  $m$  in 1D with TI potentials.

$$-\frac{\hbar^2}{2m} \frac{d^2 \psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

Once we find the wavefunction  $\psi_E(x)$  of energy  $E$ , its time dependence follows easily:

$$\psi_E(x, t) = \psi_E(x) e^{-\frac{i}{\hbar} E t}$$

### Boundary conditions

1. When the potential  $V(x)$  has a finite jump at  $x = a$ , both  $\psi(x)$  and  $\psi'(x)$  are continuous across  $x = a$ .

2. When the potential  $V(x)$  has an infinite jump at  $x = a$ ,  $\psi(x)$  is continuous but  $\psi'(x)$  is discontinuous across  $x = a$ .

Futhermore, the wavefunction must vanish at  $x = \pm\infty$  for a normalizable wavefunction.

### 2.2 The Infinite Square Well

$$V(x) = \begin{cases} 0, & \text{if } 0 \leq x \leq a \\ \infty, & \text{otherwise} \end{cases}$$

$$\psi(x) = 0 \text{ for } x < 0 \text{ and } x > a$$

For  $0 \leq x \leq a$ ,  $V(x) = 0$  and the Schrödinger equation reduces to

$$\psi''(x) + k^2 \psi(x) = 0, \text{ where } k = \sqrt{\frac{2mE}{\hbar^2}} \text{ and } E > 0$$

Classic **simple harmonic oscillator**:  $\psi(x) = A \sin(kx) + B \cos(kx)$

### Boundary conditions:

Continuity of  $\psi(x)$  at  $x = 0$  sets  $\psi(0) = B = 0 \rightarrow \psi(x) = A \sin(kx)$   
at  $x = a$  sets  $\psi(a) = A \sin(ka) = 0$

$$k_n = \frac{n\pi}{a}, n = 1, 2, \dots$$

$$\psi_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi}{a} x\right) \quad E_n = \frac{\hbar^2 k_n^2}{2m} = \frac{n^2 \pi^2 \hbar^2}{2ma^2}$$

$\psi_1$  is the ground state, others are excited states.

Properties of  $\psi_n(x)$ :

1. Alternatively even and odd.
2. As you go up in energy, each successive state has one more node.
3. They are mutually orthogonal, in the sense that  $\int \psi_m(x)^* \psi_n(x) dx = 0$  whenever  $m \neq n$ .  
 $\int \psi_m(x)^* \psi_n(x) dx = \delta_{mn}$  where  $\delta_{mn}$  (Kronecker delta) is 0 if  $m \neq n$  and 1 if  $m = n$ . We say that the  $\phi$ 's are orthonormal.
4. They are complete, in the sense that any other function,  $f(x)$ , can be expressed as a linear combination of them (Fourier series), Dirichlet's theorem:

$$f(x) = \sum_{n=1}^{\infty} c_n \psi_n(x) = \sqrt{\frac{2}{a}} \sum_{n=1}^{\infty} c_n \sin\left(\frac{n\pi}{a} x\right)$$

Fourier's trick:  $c_n = \int \psi_n(x)^* f(x) dx$

$$c_m = \frac{2}{a} \int_0^a f(x) \sin\left(\frac{m\pi x}{a}\right) dx$$

$|c_n|^2$  tells you the probability that a measurement of the energy would yield the value  $E_n$ .

Sum of these probabilities should be 1:

$$\sum_{n=1}^{\infty} |c_n|^2 = 1$$

The expectation value of the energy is

$$\langle H \rangle = \sum_{n=1}^{\infty} |c_n|^2 E_n$$

Conservation of energy in QM

### 2.3 The Harmonic Oscillator

Hooke's law (mass  $m$  w/ spring constant  $k$ ):  $F = -kx = m \frac{d^2x}{dt^2}$

Solution is  $x(t) = A \sin(\omega t) + B \cos(\omega t)$ , where  $\omega = \sqrt{\frac{k}{m}}$

Potential energy:  $V(x) = \frac{1}{2} kx^2 = \frac{1}{2} m\omega^2 x^2$

Expanding  $V(x)$  in a **Taylor series** about the min:

$$V(x) = V(x_0) + V'(x_0)(x - x_0) + \frac{1}{2} V''(x_0)(x - x_0)^2 + \dots$$

Simple harmonic oscillation,  $V(x) \cong \frac{1}{2} V''(x_0)(x - x_0)^2$ ,  $k = V''(x_0)$

The Schrödinger Equation for the harmonic oscillator:

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} + \frac{1}{2} m\omega^2 x^2 \psi = E\psi$$

Boundary conditions:  $\psi(-\infty) = 0$ ,  $\psi(+\infty) = 0$

#### 1. Simplify notation with change of variables

Introduce  $\xi \equiv \sqrt{\frac{m\omega}{\hbar}} x$ .

SE becomes  $\frac{d^2\psi}{d\xi^2} = (\xi^2 - K)\psi$ , where  $K \equiv \frac{2E}{\hbar\omega}$ .

#### 2. Asymptotic behavior

Working in the large  $\xi^2 \gg K$  region,

**Hermite eqn:**  $H''(\xi) - 2\xi H'(\xi) + (K - 1)H(\xi) = 0$

**Hermite polynomials:**  $H_0 = 1$ ,  $H_1 = 2\xi$ ,  $H_2 = 4\xi^2 - 2$ ,

$H_3 = 8\xi^3 - 12\xi$ ,  $H_4 = 16\xi^4 - 48\xi^2 + 12$ ,  $H_5 = 32\xi^5 - 160\xi^3 + 120\xi$

#### 3. Method of power series

The recursion formula:  $a_{j+2} = \frac{(2j+1-K)}{(j+1)(j+2)} a_j$

Recursion formula for allowed  $K$ :

$$a_{j+2} = \frac{2(n-j)}{(j+1)(j+2)} a_j$$

$$h(\xi) = a_0 + a_1\xi + a_2\xi^2 + \dots = \sum_{j=0}^{\infty} a_j \xi^j$$

$$h(\xi) = h_{\text{even}}(\xi) + h_{\text{odd}}(\xi)$$

#### 4. Infinite series produces a diverging function

For large  $n$ , we have  $a_{n+2} \approx \frac{2}{n} a_n$

#### 5. Truncate series

$K = 2n + 1$ , so  $E_n = (n + \frac{1}{2})\hbar\omega$

The normalized stationary states:

$$\psi_n(x) = \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} \frac{1}{\sqrt{2^n n!}} H_n(\xi) e^{-\xi^2/2}$$

Rodrigues formula:  $H_n(\xi) = (-1)^n e^{\xi^2} \left(\frac{d}{d\xi}\right)^n e^{-\xi^2}$

### 2.4 The Free Particle

$$\frac{\partial^2 \xi}{\partial x^2} = -k^2 \xi, k = \frac{\sqrt{2mE}}{\hbar}$$

General solution to the TISE: wave packet,

$$\Psi(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \psi(k) e^{i(kx - \frac{\hbar k^2}{2m} t)} dk$$

$$\phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{+\infty} \Psi(x, 0) e^{-ikx} dx$$

Plancherel's theorem:

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} F(k) e^{ikx} dk \leftrightarrow F(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-kx} dx$$

$F(k)$  is the Fourier transform of  $f(x)$ ;  $f(x)$  is the inverse Fourier transform of  $F(k)$

Phase velocity: speed of individual ripples; group velocity: speed of the envelope

Dispersion relation: the formula for  $\omega$  as a function of  $k$

### 2.5 The Delta-Function Potential

**Dirac delta function** is an infinitely high, infinitesimally narrow spike at the origin, whose area is 1:

$$\delta(x) = \begin{cases} 0, & \text{if } x \neq 0 \\ \infty, & \text{if } x = 0 \end{cases}$$

$f(x)\delta(x-a) = f(a)\delta(x-a)$  bc the product is 0 anyway except at  $a$ .

In particular,  $\int_{-\infty}^{\infty} f(x)\delta(x-a)dx = f(a)$ .  $a - \epsilon$  to  $a + \epsilon$  would do, for any  $\epsilon > 0$ .

Consider  $V(x) = -\alpha\delta(x)$ , where  $\alpha$  is some positive constant.


$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} - \alpha\delta(x)\psi = E\psi$$

**Bound states** ( $E < 0$ ):

**Scattering states** ( $E > 0$ ):

### 2.6 The Finite Square Well

$$V(x) = \begin{cases} -V_0, & \text{for } -a < x < a \\ 0, & \text{for } |x| > a \end{cases}$$



where  $V_0$  is a positive constant.

Both bound states ( $E < 0$ ) and scattering states ( $E > 0$ ).

**Bound states:**

Potential is piecewise and discontinuous, can split into regions.

**REGION I**

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} = E\psi, \text{ or } \psi''_I(x) - \kappa^2 \psi_I(x) = 0, \quad \kappa \equiv \sqrt{-\frac{2mE}{\hbar^2}}$$

where  $E < 0$  for a bound state.

General sol:  $\psi_I(x) = Ae^{-\kappa x} + Be^{\kappa x}$ .

$x = -\infty \rightarrow \psi(x) = 0$ , so  $A = 0$ , and we have  $\psi_I(x) = Be^{\kappa x}$

**REGION II**

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} - V_0\psi = E\psi, \text{ or } \psi'' = -l^2\psi, \quad l \equiv \sqrt{\frac{2m(E + V_0)}{\hbar^2}}$$

General sol:  $\psi(x) = C \sin(lx) + D \cos(lx)$ , for  $-a < x < a$

**REGION III**

SE and general sol same as region I, but  $x = \infty \rightarrow \psi(x) = 0$ , so

$G = 0$  and  $\psi_{III}(x) = Fe^{-\kappa x}$

**Even bound states**

$\psi(-x) = \psi(x)$ ,  $\psi_{II}(x) = D \cos(lx)$

Bc the potential has only a finite discontinuity at  $x = \pm a$ , both  $\psi$  and  $\psi'$  must be continuous at  $x = \pm a$ .

$x = a$ ,  $\psi_{II}(a) = \psi_{III}(a)$  imposes  $D \cos(la) = F e^{-\kappa a}$

$x = a$ ,  $\psi'_{II}(a) = \psi'_{III}(a)$  imposes  $-lD \sin(la) = -\kappa F e^{-\kappa a}$

Continuity of  $\psi(x)$  and  $\psi'(x)$  at  $x = -a$  does not add anything new.

Dividing the above two, we get  $\kappa = l \tan(la)$

This is a formula for the allowed energies, since  $\kappa$  and  $l$  are both functions of  $E$ . Let  $z \equiv la$ , and  $z_0 \equiv \frac{a}{\hbar} \sqrt{2mV_0}$ .  $\kappa^2 + l^2 = 2mV_0/\hbar^2$ ,

so  $\kappa a = \sqrt{z_0^2 - z^2}$ .

Transcendental eq for  $z$  (and hence  $E$ ) as a function of  $z_0$  (which is a measure of size of well):  $\tan z = \sqrt{(\frac{z_0}{z})^2 - 1}$

**Odd bound states**

$\psi_{II}(x) = C \sin(lx)$

$x = -a$ ,  $\psi_{II}(-a) = \psi_I(-a)$  imposes  $C \sin(la) = B e^{-\kappa a}$


$x = -a$ ,  $\psi'_{II}(-a) = \psi'_I(-a)$  imposes  $lC \cos(la) = -\kappa B e^{-\kappa a}$

Dividing the above two,  $l \cot(la) = -\kappa$ .

Rewriting this in terms of  $z$  and  $z_0$ ,

$$\cot(z) = -\sqrt{(\frac{z_0}{z})^2 - 1}$$

$V_1$  does not support an odd bound state, since there is no intersection pt,  $V_2$  produces only one bound state, and  $V_3$  produces two bound states. Finite well potential supports at least one even state, the ground state, and it may not support any of the excited states.



**Wide and deep well**

$z_n \approx \frac{n\pi}{2}$ ,  $n = 1, 2, 3, \dots$

Using the def of  $z$  and solving for  $E$ ,

$$E_n = -V_0 + \frac{\hbar^2 \pi^2 n^2}{2m(2a)^2}, n = 1, 2, \dots$$

Thus, the energy levels of the infinite square well of width  $2a$  are reproduced for  $E_n - (-V_0) = E_n + V_0$ , which is the energy above the bottom of the well. As  $V_0 \rightarrow \infty$ , finite sq well goes to infinite sq well.

**Shallow and narrow well**

Any shallow or narrow well supports at least one bound state. But we

need at least  $z_0 = \sqrt{\frac{2mV_0 a^2}{\hbar^2}} \geq \frac{\pi}{2}$  to support any odd state.

### Gaussian integrals

$$\int_{-\infty}^{\infty} e^{-ax^2} dx = \sqrt{\frac{\pi}{a}} \text{ and } \int_0^{\infty} e^{-ax^2} dx = \frac{1}{2} \sqrt{\frac{\pi}{a}}$$

$$\int_{-\infty}^{\infty} x^{2n+1} e^{-ax^2} dx = 0 \text{ for } n = 0, 1, 2, \dots$$

$$\int_{-\infty}^{\infty} x^{2n} e^{-ax^2} dx = (-1)^n \frac{d^n}{da^n} \sqrt{\frac{\pi}{a}} \text{ for } n = 0, 1, 2, \dots$$

$$\int_{-\infty}^{\infty} e^{-ax^2+bx} dx = \sqrt{\frac{\pi}{a}} e^{\frac{b^2}{4a}} \quad \int_{-\infty}^{\infty} x^n e^{-ax^2+bx} dx = \frac{d^n}{db^n} \sqrt{\frac{\pi}{a}} e^{\frac{b^2}{4a}}$$