

Flexible learning through reuse and reconfiguration of stable functional clusters

Jiayi Zhang¹, Jae-hyun Kim², Nuo Li², Tatiana A. Engel³

¹Lewis-Sigler Institute of Integrative Genomics, Princeton University, Princeton, NJ, USA

²Department of Neurobiology, Duke University, Durham, NC, USA

³Princeton Neuroscience Institute, Princeton University, Princeton, NJ, USA




Digital poster
COSYNE2026
3-158

Motivation

- Animals **acquire** and **reuse** motor skills across diverse contexts throughout their lifetime.
- When learning new task contexts, the motor cortex generates new population activity patterns while preserving some of the existing ones [1].
- How neural circuits reorganize** to support flexible skill learning remains unknown.


Longitudinal imaging across contingency reversals



Figs. adapted from Kim and Li, 2024 [1]

- Mice first learned context 1, and upon achieving proficiency, learned context 2 with reversed rule.
- Same populations** of ALM neurons (4 animals, average 1366 neurons per animal) tracked **across contexts**.

Latent circuit inference



- From neural recording y , the latent circuit inference [2] jointly infers a **subspace** Q (containing task-relevant dynamics) and a **latent circuit** (low-dimensional RNN) that generates dynamics x within Q such that $y \approx Qx$.
- Q is orthonormal; maps between latent nodes and neurons.

Model training and selection


- For each animal and task context, we fit 200 latent circuit models of each latent dimensionality (varying $n=3-12$), subject to

$$\mathcal{L}_{\text{total}} = \lambda_x \mathcal{L}_x + \mathcal{L}_z + \lambda_w \mathcal{L}_w$$


Trajectory reconstruction Task performance L1-penalty on recurrent weights

- We selected model dimensionality that balances explained variance and model uniqueness ($n=6, 8, 9, 11$ for four mice).
- Average total variance explained: 60%, average behavioral variance explained: 91%.


Model selection criteria for one mouse



Distinct functional clusters in ALM



- Neurons typically exhibited a **dominant loading on a single latent node**, forming distinct **functional clusters**.
- This is reflected in sparse rows in Q -matrices, quantified by the Hoyer sparsity index.



Cluster reorganization across contexts

Is cluster membership stable?


- We computed correspondence between clustered populations across contexts for each animal. Given Q_1, Q_2 learned for context 1, 2:

- Maximally align Q_1 with Q_2 through permutation P^* :

$$P^* = \underset{P \in \mathcal{P}}{\operatorname{argmax}} \operatorname{Tr}((PQ_1^T)Q_2)$$

- Compute $G^* = (P^*Q_1^T)Q_2$. The diagonal entries g^* quantify the cosine-similarity between matched clusters.

- 18 of 34 clusters showed $g^* > 0.5$, indicating **substantial reuse of clusters**.




Do clusters preserve or alter their response profiles?

- We computed cross-context correlations of latent trajectories on trials with 1) the same choice 2) the same stimulus.
- Most clusters **preserved** both their **temporal dynamics and selectivity** for either stimulus (example 1) or choice (example 3).
- A subset changed response profiles (example 2).

Circuit motifs

- In both contexts, the inferred recurrent weights formed chains with **feedforward excitation** and **feedback inhibition**, selective for the left and right choices. This motif was hypothesized in theoretical models [3].



Conclusions

- We identified **distinct functional clusters** in the mouse motor cortex that remain **largely stable across task contingency reversal**.
- After relearning, **most clusters preserved their selectivity and temporal profiles**, while a subset changed their responses.
- The inferred circuits exhibit a **feedforward chain** motif across contexts, consistent with theoretical predictions.

1. Kim, J. H., Daie, K., & Li, N. (2025). *Nature*, 637(8046), 663-672.

2. Langdon, C., & Engel, T. A. (2025). *Nature Neuroscience*, 28(3), 665-675.

3. Daie, K., Fontolan, L., Druckmann, S., & Svoboda, K. (2023). *bioRxiv*, 2023-08.