

Math Notebook

Markkus Hansen

Created: July 7, 2026
Last updated: July 7, 2026

Contents

1	Analysis	2
1.1	Conventions and notation	2
1.1.1	Abbreviations for references	2
1.1.2	Standard number sets	2
1.1.3	Placeholder sets	2
1.1.4	Mathematical statements $P(x)$ or $P(n)$	2
1.1.5	Explicit dependence e.g. $N = N(\varepsilon)$, $\delta = \delta(\varepsilon)$	2
1.2	Given definitions and prerequisites	3
1.2.1	Sequences	3
1.2.2	Series	3
1.2.3	Limits over real functions	4
1.2.4	Continuity	4
1.2.5	Derivatives	4
1.3	Derivatives of standard functions	4
1.3.1	Polynomials	4
1.4	L'Hôpital's rule	5
1.4.1	Motivation	5
1.4.2	Statement	5
1.4.3	Required lemmas	5
1.4.4	Proof	5
1.4.5	Examples of L'Hôpital's rule	5

1 Analysis

1.1 Conventions and notation

1.1.1 Abbreviations for references

- TL: Kalkulus - Tom Lindstrøm - 4. udgave
- An1: Analyse 1 - Matthias Christandl, Søren Eilers og Henrik Schlichtkrull - Syvende udgave

1.1.2 Standard number sets

Notation of standard sets of numbers in the analysis section of this notebook.

- $\mathbb{N} = \{1, 2, 3, \dots\}$
- $\mathbb{N}_0 = \{0, 1, 2, \dots\}$
- $\mathbb{Z}^+ = \mathbb{N}$
- $\mathbb{R}^+ = \{x \in \mathbb{R} \mid x > 0\}$

1.1.3 Placeholder sets

Unless otherwise stated, uppercase Latin letters $A, B, C, \dots$ denote subsets of $\mathbb{R}$. I.e. whenever something like $f : A \rightarrow B$ is written without further context of the domain and codomain A and B , assume that $A, B \subseteq \mathbb{R}$.

This assumption also holds for functions defined like $f : D_f \rightarrow V_f$.

1.1.4 Mathematical statements $P(x)$ or $P(n)$

Not to be confused with the power set $\mathcal{P}(S)$, the notation $P(x)$ or $P(n)$ refers to a mapping $P : D_P \rightarrow \{\perp, \top\}$ that assigns a truth value based on whether the statement holds for a given input value.¹ In this notebook, unless otherwise stated, assume that $D_P \subseteq \mathbb{R}$ when $P(x)$ is written, and assume that $D_P \subseteq \mathbb{N}$ when $P(n)$ is written.

1.1.5 Explicit dependence e.g. $N = N(\varepsilon)$, $\delta = \delta(\varepsilon)$

When you see something like " $\varepsilon \in \mathbb{R}^+$, $\delta = \delta(\varepsilon)$ " in this notebook, you may read it as "Let ε be a positive real number, and let δ be a number where its value depends in some way on the value of ε ". This notation is simply for ease of understanding the structure of proofs and theorems, and may show itself in epsilon-N proofs (for sequences) like

$$\forall \varepsilon \in \mathbb{R}^+ \exists N = N(\varepsilon) \in \mathbb{N} \text{ s.t. } n > N \implies |a_n - a| < \varepsilon \quad (1.1.5.1)$$

¹The $\{\perp, \top\}$ is equivalent to $\{\text{False}, \text{True}\}$, $\{0, 1\}$, etc.

or epsilon-delta proofs like

$$\forall \varepsilon \in \mathbb{R}^+ \exists \delta = \delta(\varepsilon) \in \mathbb{R}^+ \text{ s.t. } 0 < |x - a| < \delta \implies |f(x) - b| < \varepsilon \quad (1.1.5.2)$$

In a similar manner, if a variable y is defined as an expression containing a variable x , one may also write $y = y(x)$ since y depends on the value of x .

1.2 Given definitions and prerequisites

The following concepts are assumed known:

- Basic set theory notation
- Real number arithmetic
- Functions and mappings
- Mathematical induction
- Proof techniques

1.2.1 Sequences

A sequence is a function whose domain is an ordered subset of the integers (usually $\mathbb{N}$). If we let $n \in \mathbb{N}$ and $a_n \in \mathbb{R}$, then $(a_n)_{n \in \mathbb{N}} = (a_1, a_2, a_3, \dots)$ is said to be a real sequence. Each number a_n is called a term of the sequence.

One can view a sequence as simply being a map $a : D_a \rightarrow V_a$. The codomain may consist of any set that contains the sequence's possible outputs, for example, $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$, $\mathbb{R}$, $\mathbb{C}$, or even vector spaces like $\mathbb{R}^2$ or $\mathbb{C}^2$. A sequence $(a_n)_{n \in A}$ may also be finite over a set $A = \{n_1, n_2, \dots, n_k\}$ with $k \in \mathbb{N}$. Unless otherwise stated though, assume that any given sequence is a map $a : \mathbb{N} \rightarrow \mathbb{R}$.

A finite sequence of length two may be denoted as (a, b) . To avoid confusing it with the notation of an open interval, assume that (a, b) refers to an open interval if it is not stated to be a finite sequence.

1.2.2 Series

A series is an expression representing the sum of the terms in a finite or infinite sequence. For $(a_n)_{n \in \mathbb{N}}$ we define its corresponding series as the expression

$$\sum_{n=1}^{\infty} a_n \quad (1.2.2.1)$$

A partial sum, S_n , of the series $\sum_{k=1}^{\infty} a_k$ is the sum of the terms within a finite subsequence $(a_k)_{k \in A}$ with $A = \{k \in \mathbb{N} \mid 1 \leq k \leq n\}$.² The n -th partial sum S_n may then be defined as

$$S_n = \sum_{k=1}^n a_k \quad (1.2.2.2)$$

²Pretentious way of writing $A = \{1, 2, 3, \dots, n\}$

1.2.2.1 Limit of partial sum

The limit of the partial sum S_n converges to $S \in \mathbb{R}$ as n approaches infinity, if and only if

$$\forall \varepsilon \in \mathbb{R}^+ \exists N = N(\varepsilon) \in \mathbb{N} \text{ s.t. } n > N \implies |S_n - S| < \varepsilon \quad (1.2.2.3)$$

i.e.

$$\lim_{n \rightarrow \infty} S_n = S \quad (1.2.2.4)$$

which means that the series converges to the value S :

$$\sum_{n=1}^{\infty} a_n = S \quad (1.2.2.5)$$

1.2.3 Limits over real functions

Symbolic definition of limit for $f : \mathbb{R} \rightarrow \mathbb{R}$

$$\forall \varepsilon \in \mathbb{R}^+ \exists \delta = \delta(\varepsilon) \in \mathbb{R}^+ \text{ s.t. } \forall x \in D_f : 0 < |x - a| < \delta \implies |f(x) - b| < \varepsilon \quad (1.2.3.1)$$

This defines the standard limit notation of

$$\lim_{x \rightarrow a} f(x) = b \quad (1.2.3.2)$$

Definition is taken from TL definition 5.4.1.

1.2.4 Continuity

A function $f : D_f \rightarrow V_f$ is said to be continuous at a point $a \in D_f$ if the following holds:

$$\forall \varepsilon \in \mathbb{R}^+ \exists \delta = \delta(\varepsilon) \in \mathbb{R}^+ \text{ s.t. } \forall x \in D_f : |x - a| < \delta \implies |f(x) - f(a)| < \varepsilon \quad (1.2.4.1)$$

Definition is taken from TL definition 5.1.1.

1.2.5 Derivatives

Given a function $f : D_f \rightarrow V_f$, the derivative of f at a point $a \in D_f$ is defined as

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} \quad (1.2.5.1)$$

1.3 Derivatives of standard functions

1.3.1 Polynomials

1.3.1.1 Degree-1 polynomials

Let a function $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = x$. For the derivative for a fixed point $a \in \mathbb{R}$, we have from equation (1.2.5.1) that

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{x \rightarrow a} \frac{x - a}{x - a} = \lim_{x \rightarrow a} \frac{1}{1} = 1 \quad (1.3.1.1)$$

Redefine f to the general form $f(x) = mx + b$. We then have

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a} = \lim_{x \rightarrow a} \frac{mx + b - ma - b}{x - a} = \lim_{x \rightarrow a} \frac{m(x - a)}{x - a} = \lim_{x \rightarrow a} \frac{m}{1} = m \quad (1.3.1.2)$$

1.4 L'Hôpital's rule

1.4.1 Motivation

1.4.2 Statement

1.4.3 Required lemmas

1.4.4 Proof

1.4.5 Examples of L'Hôpital's rule