

# Understanding the spectrum of a blackbody

---

Sudheendra B.R.

Suhas P.K.

Shahsank K.K.

March 2020

Guided by V.H. Belvadi

Dept. of Physics

Yuvaraja's College, Mysuru

# Overview

---

## Part 1

The blackbody problem

Rayleigh's intensity function

The ultraviolet catastrophe

Planck's empirical formula

## Part 2

Intensity of a blackbody spectrum and the problem

Einstein's work on blackbodies

## Part 3

Amplitude-based description of identical particles

n-Boson systems

Probability-based emission and absorption of blackbodies


## Part 1

---


An idealised object that can **absorb**, and subsequently **emit**, all the radiation incident on it is called a **blackbody**.

## Rayleigh's law

---




## Rayleigh's law



electron

A diagram showing a small grey circle representing an electron. A horizontal arrow points from the circle to the right, ending in a triangular arrowhead. The word "electron" is written below the circle.



gaseous medium

A light grey rectangular area representing a gaseous medium. The words "gaseous medium" are written at the bottom center of the rectangle.

- Consider a gaseous medium into which a charged particle, say an electron, enters.


## Rayleigh's law



- Consider a gaseous medium into which a charged particle, say an electron, enters.
- Due to the atoms in the gas, the electron collides with the atoms several times



## Rayleigh's law



- Consider a gaseous medium into which a charged particle, say an electron, enters.
- Due to the atoms in the gas, the electron collides with the atoms several times
- As a result the gaseous atoms impart some of their kinetic energy to the electron.

Kinetic energy of the electron is,

$$\frac{1}{2} kT$$


And the total kinetic energy will be,

$$kT$$

- Since the electron is radiating energy  
**the system cannot immediately  
attain equilibrium**


## Rayleigh's law

- Since the electron is radiating energy  
**the system cannot immediately  
attain equilibrium**



## Rayleigh's law

- Since the electron is radiating energy **the system cannot immediately attain equilibrium**
- The inner walls of the box is totally reflective.



The electron behaves like an oscillator whose energy radiated away per second is be given by,

$$\frac{1}{Q} = \frac{\left(\frac{dW}{dt}\right)}{\omega_0 W}$$

$Q$  is the radiation reaction

$\omega_0$  is the natural frequency of vibration

$W$  is the total energy content of the oscillator

Also, we have

$$\frac{1}{Q} = \frac{\gamma}{\omega_0}$$

Where  $\gamma$  is the damping constant.

Average energy loss due to radiation is given by,

$$\left\langle \frac{dW}{dt} \right\rangle = \gamma kT$$

Since the degree of freedom is 3

$$\left\langle \frac{dW}{dt} \right\rangle = 3\gamma kT$$



- $\mathbf{I}(\omega)d\omega}$  is the incident light energy within a frequency range  $d\omega$ .
- **Asumption:** The radiation which falls on a *cross-section*  $\sigma_S$  is absorbed completely.
- The total energy scattered is the product of  $\mathbf{I}(\omega)d\omega$  and  $\sigma_S$ .

The cross-section is given by

$$\sigma_s = \frac{8\pi r_0^2}{3} \left[ \frac{\omega^4}{(\omega^2 - \omega_0^2)^2 + \gamma^2 \omega^2} \right]$$

At equilibrium,  $\omega = \omega_0$

$$\sigma_s = \frac{2\pi r_0^2 3\omega_0^2}{3(\omega - \omega_0)^2 + \frac{\gamma^2}{4}}$$

$$\frac{dW_s}{dt} = \int_0^\infty \mathbf{I}(\omega) \sigma_s d\omega = \int_0^\infty \frac{2\pi r_0^2 \omega_0^2 \mathbf{I}(\omega) d\omega}{3 \left[ (\omega - \omega_0)^2 + \frac{\gamma^2}{4} \right]}$$

$\because \omega = \omega_0,$

$$\frac{dW_s}{dt} = \frac{2}{3} \pi r_0^2 \omega_0^2 \mathbf{I}(\omega_0) \int_{-\infty}^{\infty} \frac{d\omega}{(\omega - \omega_0)^2 + \left(\frac{\gamma}{2}\right)^2}$$

The integral reduces to an arctan function and we get  $2\pi/\gamma$ .

$$\frac{dW_s}{dt} = \frac{dW}{dt} = 3\gamma kT \quad (\text{as seen earlier})$$


$$\therefore \mathbf{I}(\omega) = \frac{9\gamma^2 kT}{4\pi^2 r_0^2 \omega_0^2}$$

$$\Rightarrow \gamma = \frac{2}{3} \frac{r_0 \omega_0^2}{c}$$

The intensity distribution function  $\mathbf{I}(\omega)$  is,


$$\mathbf{I}(\omega) = \frac{\omega^2 kT}{\pi^2 c^2}$$

## The experimental observation



**Figure 1:** Spectrum of a blackbody.

A theoretical explanation for the obtained experimental results were needed.



- The harmonic oscillator can take up energies only as multiples of  $\hbar\omega$ .



- The harmonic oscillator can take up energies only as multiples of  $\hbar\omega$ .
- The probability of occupying an energy level  $\mathbf{E}$  is  $\mathbf{P}(\mathbf{E}) = \alpha e^{-\hbar\omega/kT}$ .

- The number of oscillators in ground state is  $N_0$  .

- The number of oscillators in ground state is  $N_0$  .
- The number of oscillator in first state is  $N_1 = N_0 e^{-\hbar\omega/kT}$  .

- The number of oscillators in ground state is  $N_0$  .
- The number of oscillator in first state is  $N_1 = N_0 e^{-\hbar\omega/kT}$  .
- Then  $N_n = N_0 x^n$  , where  $x = e^{-\hbar\omega/kT}$

- The energy in the ground state is zero.

- The energy in the ground state is zero.
- The energy in the first state is  $N_1 \hbar \omega$  or  $N_0 \hbar \omega$

- The energy in the ground state is zero.
- The energy in the first state is  $N_1 \hbar \omega$  or  $N_0 \hbar \omega x$
- Total energy is given by

$$E_{(total)} = N_0 \hbar \omega (0 + x + 2x^2 + \dots)$$

- The total number of oscillators is

$$N_{(total)} = N_0(1 + x + x^2 + \cdots)$$



- The total number of oscillators is

$$N_{(total)} = N_0(1 + x + x^2 + \dots)$$

- The average energy is

$$\langle E \rangle = \frac{E_{(total)}}{N_{(total)}} = \frac{N_0 \hbar \omega (0 + x + 2x^2 + \dots)}{N_0(1 + x + x^2 + \dots)}$$

$$Y = 1 + 2x + 3x^2 + \cdots$$

$$Yx = x + 2x^2 + 3x^3 + \cdots$$

$$Y(1 - x) = 1 + x + x^2 + \cdots$$

$$Y = \frac{1}{(1-x)^2}$$

Average energy

$$\langle E \rangle = \frac{N_0 \hbar \omega x / (1-x)^2}{N_0 / (1-x)}$$


$$\langle E \rangle = \frac{\hbar\omega e^{-\hbar\omega/kT}}{1 - e^{-\hbar\omega/kT}} \quad \because x = e^{-\hbar\omega/kT}$$

$$\boxed{\langle E \rangle = \frac{\hbar\omega}{e^{\hbar\omega/kT} - 1}}$$

(1)

Hence the intensity of radiation for frequency  $\omega$  is given by

$$I(\omega)d\omega = \frac{\hbar\omega^3 d\omega}{\pi^2 c^2 (e^{\hbar\omega/kT} - 1)} \quad (2)$$



**Figure 2:** Spectrum of a blackbody.

## Part 2

---

- The intensity of radiation for frequency  $\omega$  is given by

$$I(\omega)d\omega = \frac{\hbar\omega^3 d\omega}{\pi^2 c^2 (e^{\hbar\omega/kT} - 1)}$$



- The intensity of radiation for frequency  $\omega$  is given by

$$I(\omega)d\omega = \frac{\hbar\omega^3 d\omega}{\pi^2 c^2 (e^{\hbar\omega/kT} - 1)}$$

- Planck considered that the matter was quantised and not light.

- The intensity of radiation for frequency  $\omega$  is given by

$$I(\omega)d\omega = \frac{\hbar\omega^3 d\omega}{\pi^2 c^2 (e^{\hbar\omega/kT} - 1)}$$

- Planck considered that the matter was quantised and not light.
- Einstein came up with an alternate idea to solve Black body problem.

Einstein considered that even the light was quantised and considered it to be made up of photons with energy  $\hbar\omega$



- Consider two energy levels, say the  $n^{th}$  and  $m^{th}$  levels.

- Consider two energy levels, say the  $n^{th}$  and  $m^{th}$  levels.
- The probability of transitioning is proportional to intensity of the incident light.

- Consider two energy levels, say the  $n^{th}$  and  $m^{th}$  levels.
- The probability of transitioning is proportional to intensity of the incident light.

$$R_{n \rightarrow m} = N_n B_{nm} I(\omega) \quad (3)$$

- Einstein considers that there are two types of emissions that take place in an atom.

- Einstein considers that there are two types of emissions that take place in an atom.
  1. Spontaneous emission.
  2. Stimulated emission.



- The combined mathematical expression is

$$R_{m \rightarrow n} = N_m[A_{mn} + B_{mn}I(\omega)] \quad (4)$$

- At thermal equilibrium, the number of atoms going to higher energy level must be equal to the number of atoms coming to lower energy state.

- At thermal equilibrium, the number of atoms going to higher energy level must be equal to the number of atoms coming to lower energy state.
- The ratio of  $N_m$  to  $N_n$  is given by  $e^{-\hbar\omega/kT}$

At thermal equilibrium both  $R_{n \rightarrow m}$  and  $R_{m \rightarrow n}$  must be equal.

At thermal equilibrium both  $R_{n \rightarrow m}$  and  $R_{m \rightarrow n}$  must be equal.

$$N_n B_{nm} I(\omega) = N_m [A_{mn} + B_{mn} I(\omega)]$$

At thermal equilibrium both  $R_{n \rightarrow m}$  and  $R_{m \rightarrow n}$  must be equal.

$$N_n B_{nm} I(\omega) = N_m [A_{mn} + B_{mn} I(\omega)]$$

On dividing the previous equation by  $N_m$  we get,

$$B_{nm}I(\omega)e^{\hbar\omega/kT} = A_{mn} + B_{mn}I(\omega)$$

$$B_{nm}I(\omega)e^{\hbar\omega/kT} = A_{mn} + B_{mn}I(\omega)$$

The equation that Einstein got for the intensity of radiation for frequency is

$$I(\omega) = \frac{A_{mn}}{B_{nm}e^{\hbar\omega/kT} - B_{mn}} \quad (5)$$



Since Einstein considered that the formula given by Planck was correct, we should get

Since Einstein considered that the formula given by Planck was correct, we should get

$$B_{nm} = B_{mn}$$

$$\frac{A_{mn}}{B_{nm}} = \frac{\hbar\omega^3 d\omega}{\pi^2 c^2}$$

This is how Einstein derived Planck's formula but this time from a quantum mechanical viewpoint.

## Part 3

---

**P** = probability

$\phi$  = Probability amplitude

$$\mathbf{P} = |\phi|^2$$


$$\phi = \phi_1 + \phi_2$$

$$\mathbf{P} = |\phi_1 + \phi_2|^2$$

- Detector 1 is set to detect only  $\alpha$  particles and Detector 2 is set to detect only oxygen atoms.

- Detector 1 is set to detect only  $\alpha$  particles and Detector 2 is set to detect only oxygen atoms.
- Probability amplitude of the scattering is given by  $f(\theta)$  at angle  $\theta$ .

- Detector 1 is set to detect only  $\alpha$  particles and Detector 2 is set to detect only oxygen atoms.
- Probability amplitude of the scattering is given by  $f(\theta)$  at angle  $\theta$ .
- The probability of this event =  $|f(\theta)|^2$







- Set up the detectors such that the detectors would detect either  $\alpha$  particle or oxygen atom.


- Set up the detectors such that the detectors would detect either  $\alpha$  particle or oxygen atom.
- We will not distinguish which particle is which entering the detector.

- Set up the detectors such that the detectors would detect either  $\alpha$  particle or oxygen atom.
- We will not distinguish which particle is which entering the detector.
- This means that if oxygen atom in position  $\theta$ , then  $\alpha$  particle on the opposite side is at an angle  $\pi - \theta$ .







- Probability amplitude of oxygen atom =  $f(\pi - \theta)$




- Probability amplitude of oxygen atom =  $f(\pi - \theta)$
- Probability amplitude of  $\alpha$  particle =  $f(\theta)$



- Probability amplitude of oxygen atom =  $f(\pi - \theta)$
- Probability amplitude of  $\alpha$  particle =  $f(\theta)$
- The probability of a particle being detected at detector 1 =  $|f(\theta)|^2 + |f(\pi - \theta)|^2$




- Consider if both are  $\alpha$  particles,




- Consider if both are  $\alpha$  particles,
- Then we would not know which particle entered the detector, so the total probability changes to,







- Consider if both are  $\alpha$  particles,
- Then we would not know which particle entered the detector, so the total probability changes to,
- The probability of a  $\alpha$  particle being detected at detector 1 =  $|f(\theta) + f(\pi - \theta)|^2$



- If  $\theta = \frac{\pi}{2}$ , then applying this to the expression  $|f(\theta) + f(\pi - \theta)|^2$  we get,




- If  $\theta = \frac{\pi}{2}$ , then applying this to the expression  $|f(\theta) + f(\pi - \theta)|^2$  we get,
- Probability =  $4|f\left(\frac{\pi}{2}\right)|^2$ , if the particles are indistinguishable.



- Suppose the particles were distinguishable, then the probability for  $\theta = \frac{\pi}{2}$  when applied for

$$|f(\theta)|^2 + |f(\pi - \theta)|^2$$

is given by




- Suppose the particles were distinguishable, then the probability for  $\theta = \frac{\pi}{2}$  when applied for

$$|f(\theta)|^2 + |f(\pi - \theta)|^2$$

is given by

- Probability =  $2 \left| f\left(\frac{\pi}{2}\right) \right|^2$



- Suppose the particles were distinguishable, then the probability for  $\theta = \frac{\pi}{2}$  when applied for

$$|f(\theta)|^2 + |f(\pi - \theta)|^2$$

is given by

- Probability =  $2 \left| f\left(\frac{\pi}{2}\right) \right|^2$
- This shows that the probability gets doubled for indistinguishable particles.

- *Can we apply the same logic to the electron-electron scattering ?*

- *Can we apply the same logic to the electron-electron scattering ?*
- **OBSERVATION:** When we have situation in which the identity of the electron which is arriving at a point is exchanged with another one, the new amplitude interfere with old one with an opposite phase.



- *Can we apply the same logic to the electron-electron scattering ?*
- *OBSERVATION:* When we have situation in which the identity of the electron which is arriving at a point is exchanged with another one, the new amplitude interfere with old one with an opposite phase.
- In electrons case , the interfering amplitude for exchange interfere with a negative sign.

- *Can we apply the same logic to the electron-electron scattering ?*
- *OBSERVATION:* When we have situation in which the identity of the electron which is arriving at a point is exchanged with another one, the new amplitude interfere with old one with an opposite phase.
- In electrons case , the interfering amplitude for exchange interfere with a negative sign.  
Probability for electron =  $|f(\theta) - f(\pi - \theta)|^2$

Fraction of cases	Particle 1	Particle 2	Spin at D1	Spin at D2	Probability
$\frac{1}{4}$	up	up	up	up	$ f(\theta) - f(\pi - \theta) ^2$
$\frac{1}{4}$	down	down	down	down	$ f(\theta) - f(\pi - \theta) ^2$
$\frac{1}{4}$	up	down	up	down	$ f(\theta) ^2$
			down	up	$ f(\pi - \theta) ^2$
$\frac{1}{4}$	down	up	up	down	$ f(\pi - \theta) ^2$
			down	up	$ f(\theta) ^2$

**Identical particles**, also called **indistinguishable particle**, are particles that cannot be distinguished from one another.

## Identical indistinguishable particles

---

- Consider particle 'a' and particle 'b'.

## Identical indistinguishable particles

---

- Consider particle 'a' and particle 'b'.
- Let the two particles collide and get scattered in two different directions say '1' and '2' over a surface element  $ds_1$  and  $ds_2$  of the detector respectively.

## Identical indistinguishable particles

---

- Consider particle 'a' and particle 'b'.
- Let the two particle collide and get scattered in two different directions say '1' and '2' over a surface element  $ds_1$  and  $ds_2$  of the detector respectively.
- If the particles are indistinguishable then the amplitudes of these process will add up.

## Identical indistinguishable particles

---

- Consider particle 'a' and particle 'b'.
- Let the two particles collide and get scattered in two different directions say '1' and '2' over a surface element  $ds_1$  and  $ds_2$  of the detector respectively.
- If the particles are indistinguishable then the amplitudes of these processes will add up.
- Probability that the two particles arrive at  $ds_1$  and  $ds_2$  is



## Identical indistinguishable particles

---

- Consider particle 'a' and particle 'b'.
- Let the two particles collide and get scattered in two different directions say '1' and '2' over a surface element  $ds_1$  and  $ds_2$  of the detector respectively.
- If the particles are indistinguishable then the amplitudes of these processes will add up.
- Probability that the two particles arrive at  $ds_1$  and  $ds_2$  is

$$|\langle 1|a\rangle \langle 2|b\rangle + \langle 2|a\rangle \langle 1|b\rangle|^2 ds_1 ds_2$$

- Integrating over the area of the detector , if we let  $ds_1$  and  $ds_2$  range over the whole area  $(\Delta S)$  , we could count each part of the area twice since the expression  $|\langle 1|a\rangle \langle 2|b\rangle + \langle 2|a\rangle \langle 1|b\rangle|^2 ds_1 ds_2$  contains everything that can happen with any pair of surface elements  $ds_1$  and  $ds_2$ .

- Integrating over the area of the detector , if we let  $ds_1$  and  $ds_2$  range over the whole area  $(\Delta S)$  , we could count each part of the area twice since the expression  $|\langle 1|a\rangle \langle 2|b\rangle + \langle 2|a\rangle \langle 1|b\rangle|^2 ds_1 ds_2$  contains everything that can happen with any pair of surface elements  $ds_1$  and  $ds_2$ .

- $Probability_{BOSE} = \frac{(4 |a|^2 |b|^2)}{2} (\Delta S)$

- Integrating over the area of the detector , if we let  $ds_1$  and  $ds_2$  range over the whole area  $(\Delta S)$  , we could count each part of the area twice since the expression  $|\langle 1|a\rangle \langle 2|b\rangle + \langle 2|a\rangle \langle 1|b\rangle|^2 ds_1 ds_2$  contains everything that can happen with any pair of surface elements  $ds_1$  and  $ds_2$ .

- $Probability_{BOSE} = \frac{(4 |a|^2 |b|^2)}{2} (\Delta S)$

- This is just twice what we got the probability for distinguished particles.

- Consider  $n$  particles say  $a, b, c \dots$  scattered in  $n$  direction say  $1, 2, 3 \dots$

- Consider n particles say a, b, c...scattered in n direction say 1, 2, 3 ...
- Probability that each particle acting alone would go into an element of the surface  $ds$  of the detector is  $|\langle \cdot \cdot \cdot \rangle|^2 ds$ .

- **Assumption:** All particles are distinguishable.

- **Assumption:** All particles are distinguishable.
- Probability that  $n$  particles will be counted together in  $n$  different surface elements =  
 $|a_1 b_2 c_3 \dots|^2 ds_1 ds_2 \dots$



- **Assumption:** All particles are distinguishable.
- Probability that n particles will be counted together in n different surface elements =  $|a_1 b_2 c_3 \dots|^2 ds_1 ds_2 \dots$
- If the amplitude does not depend on where ds is located in the detector , then the

$$Probability = (|a|^2 |b|^2 \dots) (ds_1 ds_2 \dots)$$

- Integrating each  $dS$  over the surface  $\Delta S$  of the dectector

- Integrating each  $dS$  over the surface  $\Delta S$  of the dectector

$$(P_n)_{different} = (|a|^2 |b|^2 \dots) (\Delta S)^n$$

- Integrating each  $dS$  over the surface  $\Delta S$  of the dectector

$$(P_n)_{different} = (|a|^2 |b|^2 \dots) (\Delta S)^n$$

- Now suppose that all the particle are bosons.
- For  $n$  particles, there are  $n!$  different , but indistinguishable possibilities for which we must add the amplitudes.

- Integrating each  $dS$  over the surface  $\Delta S$  of the dectector

$$(P_n)_{different} = (|a|^2 |b|^2 \dots) (\Delta S)^n$$

- Now suppose that all the particle are bosons.
- For  $n$  particles, there are  $n!$  different , but indistinguishable possibilities for which we must add the amplitudes.

- Probability that  $n$  particles will be counted on the  $n$  surface elements is given by

- Probability that  $n$  particles will be counted on the  $n$  surface elements is given by  

$$\text{Probability} = (|a_1 b_2 c_3 \dots + a_1 b_3 c_2 \dots|^2) (ds_1 ds_2 \dots)$$
- $\text{Probability} = (|n! abc \dots|^2) (ds_1 ds_2 \dots)$
- Integrate each  $ds$  over the area  $\Delta S$  of the detector

- Probability that  $n$  particles will be counted on the  $n$  surface elements is given by  

$$\text{Probability} = (|a_1 b_2 c_3 \dots + a_1 b_3 c_2 \dots|^2) (ds_1 ds_2 \dots)$$
- $\text{Probability} = (|n! abc \dots|^2) (ds_1 ds_2 \dots)$
- Integrate each  $ds$  over the area  $\Delta S$  of the detector  

$$(P_n)_{\text{BOSE}} = n! (|abc \dots|^2) (\Delta S)^n$$



- Comparing the probability when the particles are distinguishable and indistinguishable

- Comparing the probability when the particles are distinguishable and indistinguishable  
 $(P_n)_{BOSE} = n! (|abc\dots|^2) (\Delta S)^n$

- Comparing the probability when the particles are distinguishable and indistinguishable

$$(P_n)_{BOSE} = n! (|abc \dots|^2) (\Delta S)^n$$

$$(P_n)_{different} = (|a|^2 |b|^2 \dots) (\Delta S)^n$$

- $(P_n)_{BOSE} = n! (P_n)_{different}$

- What is the probability that a boson will go into particular state when there are already  $n$  particles present ?

- When the light is emitted, a photon is "created".

## Emission and Absorption of photons

---

- When the light is emitted, a photon is "created".
- Consider that there are some atom emitting  $n$  photons.

- When the light is emitted, a photon is "created".
- Consider that there are some atom emitting  $n$  photons.
- *OBSERVATION*: The probability that an atom will emit a photon into a particular final state is increased by the factor  $(n+1)$  if there are already  $n$  photons in that state.

- In quantum mechanics we can show that



- In quantum mechanics we can show that

$$\langle \chi | \phi \rangle = \langle \phi | \chi \rangle^*$$

- In quantum mechanics we can show that

$$\langle \chi | \phi \rangle = \langle \phi | \chi \rangle^*$$

- The amplitude to get from any condition  $\phi$  to any other condition  $\chi$ .

- In quantum mechanics we can show that
$$\langle \chi | \phi \rangle = \langle \phi | \chi \rangle^*$$
- The amplitude to get from any condition  $\phi$  to any other condition  $\chi$ .
- We have that amplitude that a photon will be added to some state, say  $j$ , when there are already  $n$  photons present we can express this condition as

- In quantum mechanics we can show that
$$\langle \chi | \phi \rangle = \langle \phi | \chi \rangle^*$$
- The amplitude to get from any condition  $\phi$  to any other condition  $\chi$ .
- We have that amplitude that a photon will be added to some state, say  $j$ , when there are already  $n$  photons present we can express this condition as
$$\langle n + 1 | n \rangle = (\sqrt{n + 1})a$$

- In quantum mechanics we can show that

$$\langle \chi | \phi \rangle = \langle \phi | \chi \rangle^*$$

- The amplitude to get from any condition  $\phi$  to any other condition  $\chi$ .
- We have that amplitude that a photon will be added to some state, say  $j$ , when there are already  $n$  photons present we can express this condition as

$$\begin{aligned}\langle n+1 | n \rangle &= (\sqrt{n+1}) a \\ \langle n | n+1 \rangle &= (\sqrt{n+1}) a^*\end{aligned}$$

- In quantum mechanics we can show that

$$\langle \chi | \phi \rangle = \langle \phi | \chi \rangle^*$$

- The amplitude to get from any condition  $\phi$  to any other condition  $\chi$ .
- We have that amplitude that a photon will be added to some state, say  $j$ , when there are already  $n$  photons present we can express this condition as

$$\langle n + 1 | n \rangle = (\sqrt{n + 1}) a$$

$$\langle n | n + 1 \rangle = (\sqrt{n + 1}) a^*$$

where  $a = \langle j | a \rangle$  is the amplitude when there are no other photons are present.

- The amplitude to absorb a photon when there are  $n$  photons present is given by

- The amplitude to absorb a photon when there are  $n$  photons present is given by  
$$\langle n - 1 | n \rangle = (\sqrt{n}) a^*$$



- The amplitude to absorb a photon when there are  $n$  photons present is given by

$$\langle n-1|n\rangle = (\sqrt{n})a^*$$

- $\langle n+1|n\rangle = (\sqrt{n+1})a$

- The amplitude to absorb a photon when there are  $n$  photons present is given by

$$\langle n-1|n\rangle = (\sqrt{n})a^*$$

- $\langle n+1|n\rangle = (\sqrt{n+1})a$   
 $\langle n|n+1\rangle = (\sqrt{n+1})a^*$

- The amplitude to absorb a photon when there are  $n$  photons present is given by

$$\langle n-1|n\rangle = (\sqrt{n})a^*$$

- $\langle n+1|n\rangle = (\sqrt{n+1})a$   
 $\langle n|n+1\rangle = (\sqrt{n+1})a^*$

- The above two equations show that they are symmetric in nature.

- *Thought Experiment:* Lets us consider that there are  $n$  photons that are created in the same state, that of same frequency but they cannot be distinguished .

- *Thought Experiment:* Lets us consider that there are  $n$  photons that are created in the same state, that of same frequency but they cannot be distinguished .
- The probability that an atom can emit another photon into same state is

- *Thought Experiment:* Lets us consider that there are  $n$  photons that are created in the same state, that of same frequency but they cannot be distinguished .
- The probability that an atom can emit another photon into same state is  
 $(Probability)_{emit} = (n + 1) |a|^2$

- *Thought Experiment:* Lets us consider that there are  $n$  photons that are created in the same state, that of same frequency but they cannot be distinguished .
- The probability that an atom can emit another photon into same state is  
 $(Probability)_{emit} = (n + 1)|a|^2$
- The probability that an atom absorbs a photon into the same state is

- *Thought Experiment:* Lets us consider that there are  $n$  photons that are created in the same state, that of same frequency but they cannot be distinguished .
- The probability that an atom can emit another photon into same state is  
 $(Probability)_{emit} = (n + 1)|a|^2$
- The probability that an atom absorbs a photon into the same state is  
 $(Probability)_{absorb} = (n)|a|^2$



- Rate at which an atom will make a transition to downwards has two parts.

- Rate at which an atom will make a transition to downwards has two parts.
- Probability that it will make a spontaneous transition  $|a|^2$  is proportional to the number of photons.

- Rate at which an atom will make a transition to downwards has two parts.
- Probability that it will make a spontaneous transition  $|a|^2$  is proportional to the number of photons.
- The co-efficient of absorption, of induced emission and spontaneous emission are all equal and are related to the probability of spontaneous emission.

## The Blackbody Spectrum

---

- For each light frequency  $\omega$ , there are certain N number of atoms which have two energy states separated, given by the equation  $E = \omega\hbar$ .

## The Blackbody Spectrum

---

- For each light frequency  $\omega$ , there are certain  $N$  number of atoms which have two energy states separated, given by the equation  $E = \omega\hbar$ .
- Let  $N_e$  and  $N_g$  be the average numbers of atoms that are in excited state and ground state.

## The Blackbody Spectrum

---

- For each light frequency  $\omega$ , there are certain  $N$  number of atoms which have two energy states separated, given by the equation  $E = \omega\hbar$ .
- Let  $N_e$  and  $N_g$  be the average numbers of atoms that are in excited state and ground state.
- In thermal equilibrium at temperature  $T$ , from statistical mechanics

## The Blackbody Spectrum

---

- For each light frequency  $\omega$ , there are certain  $N$  number of atoms which have two energy states separated, given by the equation  $E = \omega\hbar$ .
- Let  $N_e$  and  $N_g$  be the average numbers of atoms that are in excited state and ground state.
- In thermal equilibrium at temperature  $T$ , from statistical mechanics

$$\frac{N_e}{N_g} = e^{\left(\frac{-\Delta E}{\omega\hbar}\right)}$$

## The Blackbody Spectrum

- For each light frequency  $\omega$ , there are certain  $N$  number of atoms which have two energy states separated, given by the equation  $E = \omega\hbar$ .
- Let  $N_e$  and  $N_g$  be the average numbers of atoms that are in excited state and ground state.
- In thermal equilibrium at temperature  $T$ , from statistical mechanics
$$\frac{N_e}{N_g} = e^{\left(\frac{-\Delta E}{\omega\hbar}\right)}$$
- NOTE: Each atom in the ground state can absorb a photon and go into the excited state and each atom in the excited state can emit a photon and go to the ground state.



- At equilibrium, the rate of these two process must be equal.

- At equilibrium, the rate of these two process must be equal.
- Rate is proportional to the probability of the event and the number of atoms present.

- At equilibrium, the rate of these two process must be equal.
- Rate is proportional to the probability of the event and the number of atoms present.
- $\bar{n}$  is the average number of photons present in a given state with the frequency  $\omega$ .

- The absorption rate from the state is  $N_g \bar{n} |a|^2$ , and the emission rate into that state is  $N_e (\bar{n} + 1) |a|^2$ .
- At equilibrium  $N_g \bar{n} |a|^2 = N_e (\bar{n} + 1) |a|^2$

- Solving for the average number of photons present in a given state with the frequency  $\omega$

- Solving for the average number of photons present in a given state with the frequency  $\omega$

$$\bar{n} = \frac{1}{e^{\hbar\omega/k_B T} - 1}$$

- Solving for the average number of photons present in a given state with the frequency  $\omega$

$$\bar{n} = \frac{1}{e^{\hbar\omega/k_B T} - 1}$$

- The energy of each photon is given by  $\frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$

- Solving for the average number of photons present in a given state with the frequency  $\omega$

$$\bar{n} = \frac{1}{e^{\hbar\omega/k_B T} - 1}$$

- The energy of each photon is given by  $\frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$
- For any harmonic oscillator, the quantum mechanical energy levels are equally spaced with a separation  $\hbar\omega$ .



- The energy levels are equally spaced and the  $n^{th}$  energy level is the the mean enegry of the oscillator.

- The energy levels are equally spaced and the  $n^{th}$  energy level is the the mean enegry of the oscillator.

$$(E)_{mean} = \frac{\hbar\omega}{e^{\hbar\omega/k_B t} - 1}$$

- The energy levels are equally spaced and the  $n^{th}$  energy level is the the mean enegy of the oscillator.

$$(E)_{mean} = \frac{\hbar\omega}{e^{\hbar\omega/k_B t} - 1}$$

Considering the boson which do not interact with each other, and in that state the whole system of particles behaves (for all quantum mechanical purpose) exactly like an harmonic oscillator.

- Analysing the Electro-magnetic field in a box, it show the properties of an harmonic oscillation.
- Thus, the number of photons in a particular state in a box, can be equated to the number of energy levels associated with the particular modes of oscillation of the electromagnetic fields.

- Mean energy in any particular modes in a box at a temperature  $T$  is given by

- Mean energy in any particular modes in a box at a temperature T is given by

$$(E)_{mean} = \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$$

- Mean energy in any particular modes in a box at a temperature T is given by


$$(E)_{mean} = \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$$

### *ASSUMPTION*

For every mode there are some atoms in the box, which have energy levels that can radiate into that mode so that each mode can get into thermal equilibrium.



- There will be billions of modes in the box and there will be many small frequency intervals  $\Delta\omega$ .



- The wave number  $k$  is given by  $k = \frac{j\pi}{\lambda}$ .
- The  $\delta k$  between successive modes is given by

- The wave number  $k$  is given by  $k = \frac{j\pi}{\lambda}$ .
- The  $\delta k$  between successive modes is given by

$$\delta k = k_{j+1} - k_j = \frac{\pi}{L}$$

- The wave number  $k$  is given by  $k = \frac{j\pi}{L}$ .
- The  $\delta k$  between successive modes is given by

$$\delta k = k_{j+1} - k_j = \frac{\pi}{L}$$

- An assumption is made that  $kL$  is large that in small interval  $\Delta k$ , there are many modes.

- $\Delta W$  is the number of modes in the interval  $\Delta k$ .

- $\Delta W$  is the number of modes in the interval  $\Delta k$ .
- This is given by  $\Delta W = \frac{\Delta k}{\delta k}$  and  $\delta k = \frac{j\pi}{L}$ .

- $\Delta W$  is the number of modes in the interval  $\Delta k$ .
- This is given by  $\Delta W = \frac{\Delta k}{\delta k}$  and  $\delta k = \frac{j\pi}{L}$ .
- Thus,  $\Delta W = \frac{L(\Delta k)}{\pi}$


- $\Delta W$  is the number of modes in the interval  $\Delta k$ .

- This is given by  $\Delta W = \frac{\Delta k}{\delta k}$  and  $\delta k = \frac{j\pi}{L}$ .

- Thus,  $\Delta W = \frac{L(\Delta k)}{\pi}$

$$\Delta W = \frac{L(\Delta k)}{2\pi}$$





- A standing wave in a rectangular box must have an integral number of half waves along each axis.
- Thus,  $\Delta W$  the number of modes for a vector wave number  $\mathbf{k}$  between the axes components  $k$  and  $k + \Delta k$  is

- A standing wave in a rectangular box must have an integral number of half waves along each axis.
- Thus,  $\Delta W$  the number of modes for a vector wave number  $\mathbf{k}$  between the axes components  $k$  and  $k + \Delta k$  is

$$\Delta W = \frac{L_x L_y L_z}{(2\pi)^3} (\Delta k_x \Delta k_y \Delta k_z) \quad (6)$$

$$dW(K) = V \frac{d^3 k}{(2\pi)^3} \quad (7)$$

- Applying the above result to find number of photon modes for photons with frequencies in the range  $\Delta k$ .
- In vacuum the magnitude of  $\mathbf{k}$  is related to the frequency by

- Applying the above result to find number of photon modes for photons with frequencies in the range  $\Delta k$ .
- In vacuum the magnitude of  $\mathbf{k}$  is related to the frequency by

$$|k| = \frac{\omega}{c}.$$

- Applying the above result to find number of photon modes for photons with frequencies in the range  $\Delta\omega$ .
- In vacuum the magnitude of  $\mathbf{k}$  is related to the frequency by

$$|\mathbf{k}| = \frac{\omega}{c}.$$

- In the frequency interval  $\Delta\omega$ , these are all the modes which correspond to  $\mathbf{k}$ 's with magnitude between  $k$  and  $k + \Delta k$ , independent of the direction.
- The "volume in the  $\mathbf{k}$ -space" between  $k$  and  $k + \Delta k$  is a spherical shell of volume

- Applying the above result to find number of photon modes for photons with frequencies in the range  $\Delta k$ .
- In vacuum the magnitude of  $\mathbf{k}$  is related to the frequency by

$$|\mathbf{k}| = \frac{\omega}{c}.$$

- In the frequency interval  $\Delta\omega$ , these are all the modes which correspond to  $\mathbf{k}$ 's with magnitude between  $k$  and  $k + \Delta k$ , independent of the direction.
- The "volume in the  $\mathbf{k}$ -space" between  $k$  and  $k + \Delta k$  is a spherical shell of volume  $4\pi(k^2)\Delta k$ .

- The number of modes is then ,



- The number of modes is then ,

$$\Delta W = \frac{V 4\pi k^2 \Delta k}{(2\pi)^3}.$$

- The number of modes is then ,

$$\Delta W = \frac{V 4\pi k^2 \Delta k}{(2\pi)^3}.$$

$$\text{substitute } k = \frac{\omega}{c}$$

- The number of modes is then ,

$$\Delta W = \frac{V 4\pi k^2 \Delta k}{(2\pi)^3}.$$

$$\text{substitute } k = \frac{\omega}{c}$$

$$\Delta W(\omega) = \frac{V 4\pi \omega^2 \Delta \omega}{(2\pi c)^3}$$

- These modes are independent, we must for double the number of modes.
- This is given by,

- These modes are independent, we must for double the number of modes.
- This is given by,

$$\Delta W(\omega) = \frac{V(\omega^3)\Delta\omega}{\pi^2 c^3} \text{ (for light).}$$

- We have shown that each mode has an average the energy

- We have shown that each mode has an average the energy

$$\bar{n}\hbar\omega = \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$$

- We have shown that each mode has an average the energy

$$\bar{n}\hbar\omega = \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1}$$

- Multiplying This by the number of modes, we get the energy  $\Delta E$  in the modes that lie in the interval  $\Delta\omega$  :

$$\Delta E = \left( \frac{\hbar\omega}{e^{\hbar\omega/k_B T} - 1} \right) \left( \frac{V\omega^3 \Delta\omega}{\pi^2 c^3} \right)$$



The photons are the bosons, which have tendency to try to get to all into the same state.

Thank you