

Metabolic routes and enzyme effects

S = substrate (cleared by it) · I = inhibitor (raises other substrates) · S/I = both.
Every pharmacokinetic interaction in the dataset is an I in one row meeting an S in the same column.

	1A2	2B6	2C8	2C9	2C19	2D6	2E1	3A4	Glucuronidation	Hydrolysis	Renal, unchanged	Unabsorbed
Acetaminophen							S		S	S		
Amphetamine IR						S					S	
Amphetamine XR						S					S	
Aspirin									S	S		
Caffeine	S											
Cimetidine	I				I	I		I			S	
Colestipol												S
Daridorexant								S				
Dextromethorphan						S		S				
Diclofenac			S	S				S	S			
Diphenhydramine	S			S		S/I						
Esomeprazole					S/I			S				
Famotidine											S	
Ibuprofen			S	S					S			
Levocetirizine											S	
Loperamide		S	S			S		S				
Melatonin	S				S							
Naproxen	S			S					S			
Propranolol	S				S	S			S			
Pseudoephedrine											S	
Sertraline		S		S/I	S/I	S/I		S/I				
Simethicone												S
Temazepam								S	S			
Triazolam								S				

Read the two contrast pairs off this grid. Cimetidine carries four I marks; famotidine carries none — same drug class, same indication. Triazolam is a single S under 3A4 with no alternative route; temazepam sits under glucuronidation, which no inhibitor in this set touches. That is why cimetidine + triazolam is major and cimetidine + temazepam is nothing at all.

cytochrome P450

phase II and non-metabolic