DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

V e r s i o n 2 . 0

KISS Study

Trial Statistical Analysis Plan

15 March 2023

This document contains confidential information. Nothing herein is to be disclosed without written consent from the sponsor.

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

TABLE OF CONTENTS

Page

SIGNATURE PAGE **3**
Introduction **5**
1. Statistical design **5**
2. Selection of Study Population **5**
2.1 Inclusion Criteria 5
2.2 Exclusion Criteria 5
3. Definition of the Analysis Population **7**
4. Sample size determination **7**
5. Handling of Missing Data **7**
6. Endpoints **8**
6.1 Primary endpoint 8
6.2 Secondary endpoints 8
7. Statistical Methods **9**
7.1 General principles 9
7.2 Patient, procedural and imaging characteristics 9
7.3 Analysis of primary endpoint 10
7.4 Analysis of secondary endpoints 11
7.5 Subgroup Analyses 11
7.6 Other data 11
8. Refences **12**
9. Appendix **12**

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

SIGNATURE PAGE

TITLE: KISS clinical trial

PROTOCOL NUMBER: MED-04-Version 2.0 (2020-07-01)

Responsible Personnel

DocuSigned by:
C4981411F64E4D6...

22-mars-23

Stéphane Cook, M.D., PhD, General Manager, CERIC, Villars-sur-Glâne, Switzerland — Date

DocuSigned by:
524718D9E9AE4E5...

23-mars-23

Bernard Chevalier, M.D., Coordinating Investigator, Hôpital Privé Jacques Cartier, Massy, France — Date

DocuSigned by:
Kris Bogaerts
B352E3687BCE477...

22-Mar-23

Kris Bogaerts, Ph.D., Trial Statistician, I-BioStat, KU Leuven, Belgium — Date

DocuSigned by:
Peter Smits
0945CA73022D46A...

23-Mar-23

Peter Smits, M.D., PhD, Medical Director, CERC, Massy, France — Date

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

History of changes:

Date	Changes
15/03/2023	Added 4 Corelab tables

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

INTRODUCTION

This statistical analysis plan for KISS contains the following:

- 1. Statistical design/model of the study
- 2. Definition of the analysis populations
- 2. Handling of missing data
- 3. Statistical methods

1. Statistical design

The KISS trial is a randomized (1:1), parallel, two-arm, multicenter, prospective, post-market, non-inferiority study.

Five hundred ninety-six (596) patients with coronary heart disease presenting with de novo coronary bifurcation lesion with >70% lesion in the main vessel requiring revascularization. If all eligibility criteria and none of the exclusion criteria (See section 2) are fulfilled, patients will be allocated by permuted block randomization in a 1:1 ratio to

a) No side branch (SB) intervention (test arm)

b) Side branch ballooning (control arm)

Follow-up for all randomized subjects will be 12 months. The randomization is stratified per site.

The primary objective is to evaluate the non-inferiority of no side branch (SB) intervention: (test arm), versus side branch ballooning (control arm), in the setting of single stenting with systematic proximal optimization technique (POT), in the treatment of bifurcation lesion. Secondary objectives are to observationally describe the differences between the two arms regarding procedural and clinical endpoints at 1 month and 12 months.

2. Selection of Study Population

All patients will be expected to meet all inclusion and exclusion criteria before entering the study:

2.1 Inclusion Criteria

1. Coronary artery disease requiring revascularization that is amenable to percutaneous coronary intervention (PCI)
2. Any de novo Medina type bifurcation lesion with >70% lesion in the main vessel by visual assessment except Medina 0,0,1 if the planned strategy does not include a main branch stenting
3. Side branch diameter compatible with a PCI with stent implantation (≥ 2.25mm)
4. Side branch requiring a protection wire according to operator evaluation

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

5. Anatomy compatible with a proximal optimization technique
6. Post main branch treatment (stenting then POT): Absence of side branch occlusion Or low flow associated with ongoing ischemia (either chest pain or ST changes)
7. Male or female patient >18 years old.

2.2 Exclusion Criteria

1. Acute myocardial infarction with ongoing ST-elevation
2. Cardiogenic shock
3. Requirement for ongoing hemodialysis
4. Life-expectancy limited to <12 months due to co-morbid condition
5. Lesion involves the left main coronary artery
6. Lesion requiring a double stenting
7. Plan to treat >1 other coronary vessel at the time of the index procedure
8. Two lesions to be treated in the same session: unsuccessful PCI on the single nonbifurcation lesion in a non-target vessel before the treatment of the target lesion
9. Chronic total occlusion of any target vessel
10. Left ventricular ejection fraction <20%
11. Side branch TIMI Flow <3
12. Pre-dilatation of the Side Branch during the procedure prior to randomization
13. Known allergy to Aspirin
14. Known allergy to Clopidorel and Prasugrel and Ticagrelor
15. Known allergy to stent drug elutant
16. Known allergy to any other component of Onyx Resolute stent
17. Ongoing participation in another investigational device or drug study
18. Inability to provide informed consent
19. Patients under judicial protection, tutorship or curatorship
20. Pregnant and breast-feeding women or intention to become pregnant prior to completion of all follow-up procedures.

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

3. Definition of the Analysis Population

Three analysis populations are defined as follows:

Intention-to-treat set (ITT): based on randomized treatment.
The ITT set includes all randomized patients. Patients will be analyzed based on their allocated treatment arm, irrespective of any crossing-over or treatment protocol violations.

Per Protocol set (PP): based on randomized treatment.
Patients from the ITT set with major protocol deviations will be excluded from the PP set. Patients in the PP set will be analysed based on the procedure arm to which they were originally allocated to.The PP set will be reviewed and finalized at a Blinded Data Review Meeting (BDRM) which will take place after all data have been collected and cleaned and prior to the database lock. All major protocol deviations that lead to exclusion from the PPS will be fully documented in the Analysis Sets Specification Document that will be dated and signed at the end of the BDRM.

As-treated set (AT): based on received treatment.
AT set includes all patients randomized and who underwent the procedure. Patients in the AT set are analysed in the treatment group of which procedure actually was received.

The primary analyses will be performed on the ITT and PP set. The AT set will serve for sensitivity analyses.

4. Sample size determination

Assuming a rate of 15% myocardial infarction (MI) as defined by ARC-2 criteria in the control group, taking a 6% drop out rate into account and using an absolute non-inferiority limit of 7.5%, 596 subjects randomized equally between the 2 arms will have 80% power to show non-inferiority at a one-sided significance level of 0.05.

5. Handling of Missing Data

Best efforts will be made to collect complete data for the clinical endpoints of interest, regardless of whether or not the subject received study device.

Multiple imputation is used to account for missing data for the primary endpoint. A total of 100 imputations will be done, and the seed number will be set to 95872. As for each patient the primary endpoint is binary (MI versus no MI at discharge or 48h post-procedure), the imputation is based on a logistic regression model.

A logistic regression model will be estimated with the following covariates:

- Treatment group
- Age at time of randomization

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

- Sex.

Based on this model, the missing data will be imputed.

If necessary, for the imputation of the clinical variables, the fully conditional specification method will be used with a normal or logistic regression.

For a sensitivity analysis, a similar imputation method will be applied with the exception that for all deceased patients within the first 48 hours are assumed to show MI at discharge or 48h post-procedure.

For the secondary endpoints nor the subgroup variables, no imputation is foreseen.

6. Endpoints

6.1 Primary endpoint

The primary endpoint is the rate of peri-procedural MI at discharge or 48h post-procedure whichever comes first as judged by clinical evaluation in association with day-1 troponin evaluation using the ARC-2 definition.

6.2 Secondary endpoints

All secondary endpoints are exploratory and predefined.

Procedural safety endpoints (of index procedure)

- Technical success: deployment of main branch (MB) stent with <20% residual stenosis and a patent side branch at end of procedure upon corelab analysis
- Total procedure time
- Acute gain in main and side branches as measured with a dedicated quantitative coronary analysis (QCA) software
- Fluoroscopy time
- Total radiation dose (air kerma)
- Total contrast volume
- Need to cross over (any intervention on SB in the test arm or failure to intervene on the SB in the control arm)
- Highest <48h post-procedure level of troponin expressed as multiple of the upper limit of normal

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Clinical endpoints (at 1 month and 12 months)

- Target Lesion Failure (TLF). Composite of:
    - Cardiac Death
    - Target vessel Myocardial Infarction
    - Clinically driven Target lesion revascularization (ci-TLR)
- All-cause mortality
- Main Branch TLR
- Side Branch TLR
- Definite or Probable stent thrombosis (ARC-2 criteria)
- Angina status according to the Canadian Cardiovascular Society (CCS) classification (Grade 1 to 4)

7. Statistical Methods

Statistical analysis will be programmed using the SAS® system version 9.4 or higher.

7.1 General principles

For all endpoints, the time will start counting at the day of the procedure. That is, the day of the procedure is set as day 0. Hence, e.g. Day 30 = Day of procedure + 30.

For all time-to-event endpoints, the event rates and the 95% confidence intervals (CIs) in both treatment groups are calculated by complementing the survival rates. The 95% CIs are calculated using a log-log transformation.

For all time-to-event endpoints, when there are no events present in one or both arms, the CIs for the events rate for those arms and the risk difference will not be calculated. In addition, also the p-value will not be calculated in this case.

The standard error in competing risk analyses will be determined using the delta method.

7.2 Patient, procedural and imaging characteristics

Comparison between the two treatment groups will be performed on demographical, clinical, procedural and imaging characteristics. In general, continuous variables will be presented as mean ± standard deviation or median and interquartile range depending on their distribution. Categorical variables will be stated as number and proportion. For independent data, characteristics will be compared between the two

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

treatment groups by means of unpaired t-tests or Wilcoxon test for continuous variables and chi-square tests or (Fisher's) exact tests for categorical variables, whichever is appropriate. The reported p-values will be two-sided. For clustered data like characteristics of multiple procedures per patient, characteristics will be compared between the two treatment groups by means of a generalized estimating equation (GEE) model (Liang and Zeger, 1986) or a Wilxocon test for clustered data (Rosner et al., 2003), whichever is appropriate. For the GEE model, an independent working correlation matrix will be taken.

7.3 Analysis of primary endpoint

The hypothesis with regard to the primary clinical endpoint is formulated as follows:

H_0: $p_A - p_B \geq \delta$

H_1: $p_A - p_B < \delta$

where p_A and p_B stand for the risks of peri-procedural MI at discharge or 48h post-procedure without side branch intervention and with side branch ballooning, respectively. δ represents the non-inferiority margin, and is set to 7.5%.

The null hypothesis (H_0) states that the risk of peri-procedural MI at discharge or 48h post-procedure is higher without side branch intervention versus with side branch ballooning by at least δ. Hence, the alternative hypothesis (H_1) postulates that the risk of peri-procedural MI at discharge or 48h post-procedure without side branch intervention will be no higher than δ in comparison to treatment with side branch ballooning. Rejection of the null hypothesis is needed to conclude non-inferiority.

The hypothesis test will be performed using a Farrington-Manning Score Test. The test statistic is calculated as follows:

$$z = \frac{\hat{p}_A - \hat{p}_B - \delta}{SE(\hat{p}_A - \hat{p}_B)} = \frac{\hat{p}_A - \hat{p}_B - \delta}{\sqrt{\tilde{p}_A(1-\tilde{p}_A)/n_A + \tilde{p}_B(1-\tilde{p}_B)/n_B}}$$

In this formula, $\tilde{p}_A$ and $\tilde{p}_B$ are the maximum likelihood estimates of the risks under the restriction that the risk difference is equal to δ. Further details on these calculations can be found in Farrington and Manning, 1990. The *p*-value is calculated assuming that the test statistic has a standard normal distribution. The risk difference and its associated 1-sided 95% confidence interval will be presented with the z-test. Non-inferiority will be concluded if the upper limit of this interval is lower than δ, for both the ITT and the PP analysis. In case non-inferiority is met for the primary endpoint, a superiority analysis will be performed using a two-tailed significance level of 0.05.

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

In case multiple imputation has been performed, the procedure of Li, Raghunathan and Rubin (1991), as implemented in PROC MIANALYZE, will be performed in order to test the null hypothesis. A sensitivity analysis in which patients who died in-hospital within the first 48h will be considered too.

Note that the analysis of the primary endpoint deviates from the suggested analysis in the clinical investigation plan.

7.4 Analysis of secondary endpoints

Secondary analyses will be performed observationally. To this aim, differences between the two arms regarding the secondary endpoints will be evaluated for hypotheses generating purposes and future studies development.

The differences between the two arms regarding TLF, main branch TLR, side branch TLR and definite or probable stent thrombosis are analysed as follows: the events rates are determined from the cumulative incidence functions. For TLF, non-cardiovascular mortality is taken into account as a competing risk. For main branch TLR, side branch TLR and definite or probable stent thrombosis, mortality (cardiovascular and non-cardiovascular) is taken into account as competing risk. The between group comparisons will be done by presenting the risk differences with a two-sided 95% CI and a Z-test. These analyses are performed at 1-month and 12-months.

The difference between the two arms regarding the proportion improvers in angina status from baseline to 1-month and 12-months, the technical success and need to cross over will be analysed using a chi square test, or Fisher's exact test if there are cells with expected counts lower than 5 patients. In addition, an (exact) risk difference with a 95% confidence interval will be reported.

The difference between the two arms regarding the total procedure time, acute gain in main and side branches, fluoroscopy time, total radiation dose and total contrast dose will be analysed using a 2-sample t-test, or a Wilcoxon rank-sum test if the data shows serious deviations from a normal distribution. The mean difference or Hodges-Lehman estimate of the median difference between the two treatment groups with 95% confidence interval will be reported.

7.5 Subgroup Analyses

Subgroup analyses will be done for the primary endpoint only for the subgroups listed below. The Farrington-Manning Score Test to show non-inferiority will be applied for

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

each subgroup level. A test for interaction between subgroup and treatment group will be performed using a logistic regression model.

- Sex (Male, Female)
- Age (<70, ≥ 70)
- Target vessel type (LAD vs non-LAD)
- Medina class (XX1 vs XX0)
- Angiographic diameter of side branch (<2.25 vs ≥2.25 by corelab)
- Side branch residual stenosis (< 50% versus ≥ 50% by corelab QCA)

Within the SB intervention arm, additional analyses will be performed to compare SB ballooning vs Kissing Balloon Technique (

8. References

Farrington, C.P. and Manning, G., 1990. Test statistics and sample size formulae for comparative binomial trials with null hypothesis of nonzero risk difference or nonentity relative risk. Stat. Med., 9: 1447-1454. Doi: 10.1002/sim.4780091208

Li KH, Raghunathan TE and Rubin DB (1991). Large-Sample Significance Levels from Multiply Imputed Data Using Moment-Based Statistics and an F Reference Distribution. Journal of the American Statistical Association, Vol. 86, No. 416, 1065-1073

Liang, KY and Zeger SB, Longitudinal data analysis using generalized linear models, *Biometrika*, Volume 73, Issue 1, April 1986, Pages 13–22

Rosner B, Glynn RJ, Lee ML. Incorporation of clustering effects for the Wilcoxon rank sum test: a large-sample approach. Biometrics. 2003 Dec;59(4):1089-98. doi: 10.1111/j.0006-341x.2003.00125.x.

9. Appendix

Tables and graph templates, to be used in the statistical Report

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 1 Subject Disposition

Patient Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	Total (N=XX)
Total Number of Subjects	N			XX
Randomised	N	xx	xx	xx
Treatment				
Randomised Treatment Received	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Crossover Treatment Received	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
No Treatment Received	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Included in Intent-to-Treat Population (ITT)	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Included in Per Protocol Population (PP)	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Included in As Treated Population (AT)$	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Excluded from PP	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Protocol Violation 1	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Protocol Violation 2	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Protocol Violation 3	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Number of Index Procedures	N	xx	xx	xx
Number of Staged Procedures	N	xx	xx	xx
Number of Lesions	N	xx	xx	xx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Patient Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	Total (N=XX)
Hospital Status (Index Hospitalisation)				
Discharged	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Death	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Unknown	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Day 30 Visit Status				
Done	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Death Prior to Visit	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Termination Study	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Lost to Follow-UP	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Month 12 Visit Status				
Done	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Death Prior to Visit	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Termination Study	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Lost to Follow-UP	N (%)	xx (xx.x)	xx (xx.x)	xx (xx.x)
Total Length of Follow-Up [months]	[n] Median (Q1; Q3)	[xx] xx.x (xx.x; xx.x)	[xx] xx.x (xx.x; xx.x)	[xx] xx.x (xx.x; xx.x)

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 2.1 Baseline Characteristics: Demography (ITT)

Patient Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Gender				0.xxxx
Male	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Female	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Age (yrs)	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
BMI (kg/m²)	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Heart Rate (beats/min)	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Systolic blood pressure (mmHg)	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Patient Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Diastolic blood pressure (mmHg)	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Angina status				0.xxxx
Silent Ischemia	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Stable angina	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Unstable Angina	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
NSTEMI	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Angina class				0.xxxx
CCS 1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CCS 2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CCS 3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CCS 4	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
EF [%]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 3.1 Medical history and Diagnosis (ITT)

Medical History	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
History of Cerebrovascular Disease	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Peripheral Vascular Disease	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Previous Coronary Intervention	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Previous Myocardial Infarction	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Previous CABG				0.xxxx
History of Congestive Heart Failure	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
History of Renal Insufficiency	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Any Risk Factors	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Current Smoker	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Former Smoker				
Diabetes	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Hypertension	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Hypercholesterolemia	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Family History of Coronary Disease	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 4.1 Pre-Procedural Characteristics (ITT)

Pre-Procedural Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Vital Signs				
Heart Rate (beats/min)	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Systolic blood pressure (mmHg)	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Diastolic blood pressure (mmHg)	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Angina Status				
Angina status				0.xxxx
Silent Ischemia	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Stable angina	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Unstable Angina	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Pre-Procedural Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
NSTEMI	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Angina class				0.xxxx
CCS 1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CCS 2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CCS 3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CCS 4	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
EF [%]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Medications Before Procedure				
Any routine medication before procedure	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Aspirine	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
P2/Y12 inhibitors	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Clopidogrel	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Ticagrelor	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Prasugrel	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
OAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
VKA	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Warfarine	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Acenocoumarol	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
phenprocoumon	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
fluindione	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Pre-Procedural Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
NOAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Dabigatran	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Rivaroxaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Apixaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Endoxaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Ticlopidine	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Oral anticoagulation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Statins	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Other Important Medication	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
ECG				
Cardiac Rhythm				
Sinusal	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Atrial Fibrillation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Pacemaker	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Right Bundle Branch Block	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Left Bundle Branch Block	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Q-Waves	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
ST-T Abnormalities	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Blood Samples				
Creatinine [µmol/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Pre-Procedural Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Haemoglobine [mg/dL]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Platelets [10^9/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Cardiac Enzymes				
Troponin hsT [µg/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Troponin hsT [µg/L]				0.xxxx
Within range	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range, but not Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range and Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Troponin hsI [µg/L]	N	XX	XX	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Pre-Procedural Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Troponin hsI [µg/L]				0.xxxx
Within range	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range, but not Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range and Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
CK [IU/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
CK [IU/L]				0.xxxx
Within range	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range, but not Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range and Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CK-MB [µg/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Pre-Procedural Characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
CK-MB [µg/L]				0.xxxx
Within range	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range, but not Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range and Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 5.1.1 Index Procedure: Procedural Characteristics (ITT)

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Total Number of Procedures	N	xxx	xxx	
Approach				0.xxxx
Right Femoral Artery	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Left Femoral Artery	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Right Radial Artery	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Left Radial Artery	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Sheath Size				0.xxxx
6F	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
7F	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
8F	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Other	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Periprocedural Medication				0.xxxx
Heparin	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
LMWH	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
ACT Value [second]	N	xx	xx	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
NTL Successfully Treated Before Treatment of Targeted Bifurcation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Total Number of Study Stents Placed	Sum	xx	xx	
Total Number of Non-Study Stents Placed	Sum	xx	xx	
Radiation Exposure (Dose Area Product) [Gy.cm²]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 6.2.1 Index Procedure: Bifurcation Lesion (ITT)

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Total Number of Procedures	N	xxx	xxx	
Medina Classification				0.xxxx
1,1,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Tortuosity				0.xxxx
None	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Mild	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Moderate	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Severe	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Main Branch Lesion Length [mm]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Main Branch (MB) TIMI Flow Pre-Procedure				0.xxxx
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Proximal MB Reference Vessel Diameter [mm]	N	XX	XX	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Proximal MB Stenosis Pre-Procedure [%]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Distal MB Reference Vessel Diameter [mm]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Distal MB Stenosis Pre-Procedure [%]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Side Branch (SB) TIMI Flow Pre-Procedure				0.xxxx
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Side Branch Lesion Length [mm]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Side Branch Reference Vessel Diameter [mm]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Side Branch Stenosis Pre-Procedure [%]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Preparation of Main Branch				0.xxxx
Balloon	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Cutting Balloon	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Rotablator	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Shock-Wave	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Other	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Not Done	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Total Number of Study Stents Implanted in MB	Sum	Xx	Xx	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Non-Study Stents Implanted in MB	N	Xx	xx	
POT (Proximal Optimization Technique)				
Maximum Post-Dilatation Balloon Diameters [mm]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Maximum Post-Dilatation Balloon Length[mm]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Maximum Pressure in Main Branch [atm]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 7.3.1 Index Procedure: Side Branch (ITT)

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Total Number of Procedures	N	xxx	xxx	
If Randomised to SB, Need to Cross-Over	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Failure to Wire the SB	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Failure to Place the Balloon in the SB	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Other	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
If Randomised to No SB, Need to Cross-Over	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
SB Occlusion	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
SB Dissection	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
SB Low Flow	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Ischaemia	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
SB Intervention Done	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch Intervention				0.xxxx
Kissing Balloon	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
SB Ballooning	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Kissing Balloon				0.xxxx
Inflation Successful	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
MB: Balloons Used for Kissing Inflation				
Compliant	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Non-Compliant	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Maximum Kissing Balloon Diameter [mm] (MB)	[n] Mean (SD)	[xx] xx.x (xx.xx)	[xx] xx.x (xx.xx)	0.xxxx
	Median (Q1; Q3)	xx.x (xx.x; xx.x)	xx.x (xx.x; xx.x)	
Maximum Kissing Balloon Length [mm] (MB)	[n] Mean (SD)	[xx] xx.x (xx.xx)	[xx] xx.x (xx.xx)	0.xxxx
	Median (Q1; Q3)	xx.x (xx.x; xx.x)	xx.x (xx.x; xx.x)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Maximum Kissing Balloon Pressure in MB [atm]	[n] Mean (SD)	[xx] xx.x (xx.xx)	[xx] xx.x (xx.xx)	0.xxxx
	Median (Q1; Q3)	xx.x (xx.x; xx.x)	xx.x (xx.x; xx.x)	
SB: Balloons Used for Kissing Inflation				
Compliant	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Non-Compliant	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Maximum Kissing Balloon Diameter [mm] (SB)	[n] Mean (SD)	[xx] xx.x (xx.xx)	[xx] xx.x (xx.xx)	0.xxxx
	Median (Q1; Q3)	xx.x (xx.x; xx.x)	xx.x (xx.x; xx.x)	
Maximum Kissing Balloon Length [mm] (SB)	[n] Mean (SD)	[xx] xx.x (xx.xx)	[xx] xx.x (xx.xx)	0.xxxx
	Median (Q1; Q3)	xx.x (xx.x; xx.x)	xx.x (xx.x; xx.x)	
Maximum Kissing Balloon Pressure in SB [atm]	[n] Mean (SD)	[xx] xx.x (xx.xx)	[xx] xx.x (xx.xx)	0.xxxx
	Median (Q1; Q3)	xx.x (xx.x; xx.x)	xx.x (xx.x; xx.x)	
SB Ballooning				0.xxxx
Inflation Successful	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Balloons Used for SB Ballooning				
Compliant	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Non-Compliant	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Maximum SB Balloon Diameter [mm] (MB)	[n] Mean (SD)	[xx] xx.x (xx.xx)	[xx] xx.x (xx.xx)	0.xxxx
	Median (Q1; Q3)	xx.x (xx.x; xx.x)	xx.x (xx.x; xx.x)	
Maximum SB Balloon Length [mm] (MB)	[n] Mean (SD)	[xx] xx.x (xx.xx)	[xx] xx.x (xx.xx)	0.xxxx
	Median (Q1; Q3)	xx.x (xx.x; xx.x)	xx.x (xx.x; xx.x)	
Maximum Balloon Pressure in SB [atm]	[n] Mean (SD)	[xx] xx.x (xx.xx)	[xx] xx.x (xx.xx)	0.xxxx
	Median (Q1; Q3)	xx.x (xx.x; xx.x)	xx.x (xx.x; xx.x)	
Further Treatment to SB Needed	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Stent Implanted	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Total Number of Study Stents Implanted in SB	Sum	Xx	xx	
Technique Used				0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
T-Stenting	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
TAP	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Culotte	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Inverted Crush	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Other	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Non-Study Stent Implanted	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch Occlusion	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Side Branch Low Flow Associated with Ischaemia	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Main Branch Post TIMI Flow				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Main Branch Stenosis Post-Procedure [mm]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Side Branch Post TIMI Flow				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch Stenosis Post-Procedure [mm]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 8.4.1 Index Procedure: Procedure Results (ITT)

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Total Number of Procedures	N	xxx	xxx	
Procedural Success	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Complete Revascularisation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Non-Complete Revascularisation: Therapeutic Decision	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Medical Treatment	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Surgical	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Staged Procedure	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
None	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Procedural Complication	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Main Branch	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Coronary Dissection	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Perforation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Decrease of TIMI Flow <3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Acute Occlusion	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Thrombus Formation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Significant Air Embolism	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Severe Arrhythmia	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Other	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Coronary Dissection	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Perforation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Decrease of TIMI Flow <3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Acute Occlusion	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Thrombus Formation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Significant Air Embolism	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Severe Arrhythmia	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Other	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Main Branch Complication				
Resolved	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Resolved with Sequelae	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Ongoing	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Death	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch Complication				
Resolved	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Resolved with Sequelae	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Ongoing	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Death	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Global Outcome Death	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 5.1 Post-Procedure (ITT)

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Angina Status				
Angina status				0.xxxx
Asymptomatic	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Angina	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Angina class				0.xxxx
CCS 1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CCS 2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CCS 3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CCS 4	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
ECG				
Cardiac Rhythm				
Sinusal	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Atrial Fibrillation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Pacemaker	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Right Bundle Branch Block	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Left Bundle Branch Block	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Q-Waves	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
ST-T Abnormalities	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Blood Samples				
Creatinine [µmol/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Haemoglobine [mg/dL]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Platelets [10^9/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Cardiac Enzymes				
Troponin hsT [µg/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Troponin hsT [µg/L]				0.xxxx
Within range	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Out-of-range, but not Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range and Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Troponin hsI [µg/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
Troponin hsI [µg/L]				0.xxxx
Within range	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range, but not Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range and Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CK [IU/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
CK [IU/L]				0.xxxx
Within range	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range, but not Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range and Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
CK-MB [µg/L]	N	XX	XX	0.xxxx
	Mean	xx.x	xx.x	
	Median	xx.x	xx.x	
	Std	xx.xx	xx.xx	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	(Q1, Q3)	(xx.x,xx.x)	(xx.x,xx.x)	
	Range	(xx.x,xx.x)	(xx.x,xx.x)	
CK-MB [µg/L]				0.xxxx
Within range	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range, but not Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Out-of-range and Clin. Sign.	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 6.1 Antithrombotic regimen during follow-up (ITT)

Antithrombotic regimen	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
At 1 month				
Aspirine	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
P2/Y12 inhibitors				
Clopidogrel	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Ticagrelor	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Prasugrel	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
OAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
VKA	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Warfarine	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Acenocoumarol	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
phenprocoumon	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
fluindione	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
NOAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Dabigatran	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Rivaroxaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Apixaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Endoxaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Aspirine + P2Y12	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Aspirine or P2Y12 + VKA or NOAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
At 6 months				
Aspirine	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
P2/Y12 inhibitors				
Clopidogrel	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Ticagrelor	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Antithrombotic regimen	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Prasugrel	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
OAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
VKA	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Warfarine	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Acenocoumarol	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
phenprocoumon	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
fluindione	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
NOAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Dabigatran	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Rivaroxaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Apixaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Endoxaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Aspirine + P2Y12	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Aspirine or P2Y12 + VKA or NOAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
At 1 year				
Aspirine	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
P2/Y12 inhibitors				
Clopidogrel	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Ticagrelor	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Prasugrel	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
OAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
VKA	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Warfarine	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Acenocoumarol	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
phenprocoumon	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Antithrombotic regimen	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
fluindione	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
NOAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Dabigatran	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Rivaroxaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Apixaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Endoxaban	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Aspirine + P2Y12	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Aspirine or P2Y12 + VKA or NOAC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 7.1 Primary endpoint (ITT)

	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	Risk Difference (95% CI)	P-value
Peri-Procedural MI at Discharge or 48h Post-Procedure	n/N	xx/XX	xx/XX		
	% (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
				xx.x (xx; xx.x)	0.xxxx£
Sensitivity analysis*	n/N	xx/XX	xx/XX		
	% (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
				xx.x (xx; xx.x)	0.xxxx£

n: number of events; N= number of patients in the analysis; $: one-sided non-inferiority test with a margin of 7.56%.

£: two-sided superiority test

Missing data were accounted for by performing 100 multiple imputations.

*Patients who died in-hospital within the first 48 hours will be considered to have the primary endpoint.

Note: superiority test will only be performed is non-inferiority has been shown.

Table 8.1 Secondary Procedural Safety Endpoints (ITT)

Endpoint	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	Difference (95% CI)	P-value
Technical Success$	n	xx	xx		
	% (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
Procedure Time [min]@	n	xx	xx		
	% (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
Acute Gain in Main Branch@	n	xx	xx		
	Mean (SD)	xx.x (xx.xx)	xx.x (xx.xx)	xx.x (xx.x; xx.x)	0.xxxx
Acute Gain in Side Branch@	n	xx	xx		
	Mean (SD)	xx.x (xx.xx)	xx.x (xx.xx)	xx.x (xx.x; xx.x)	0.xxxx
Fluoroscopic Time [min]@	n	xx	xx		
	Mean (SD)	xx.x (xx.xx)	xx.x (xx.xx)	xx.x (xx.x; xx.x)	0.xxxx
Total Radiation Dose (Air Kerma) [mGy]@	n	xx	xx		
	Mean (SD)	xx.x (xx.xx)	xx.x (xx.xx)	xx.x (xx.x; xx.x)	0.xxxx
Total Contrast Volume [mL]@	n	xx	xx		
	Mean (SD)	xx.x (xx.xx)	xx.x (xx.xx)	xx.x (xx.x; xx.x)	0.xxxx
Need to Cross-Over$	n	xx	xx		
	% (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx

n/N: number of events on number of patients with data; n : number of patients with data; CI:2-sided confindence interval; SD: Standard Deviation

$ Analysed using a chi-square test.

OR $ Analysed using Fisher's exact test. Confidence intervals are exact confidence intervals.

@ Analysed using 2-sample t-test.

OR @ Analysed using Wilcoxon rank-sum test. Differences (95% CI) are estimated by the Hodges-Lehman estimator for the mean difference.

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 9.1 Secondary Clinical Endpoints (ITT)

Endpoint	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	Risk Difference (95% CI)	P-value
Target Lesion Failure					
At 1 month	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
At 1 year	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
Cardiac Death					
At 1 month	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
At 1 year	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
Target Vessel Myocardial Infarction					
At 1 month	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
At 1 year	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
Clinically Driven Target Lesion Revascularization (TLR)					
At 1 month	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
At 1 year	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
All-cause mortality					

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Endpoint	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	Risk Difference (95% CI)	P-value
At 1 month	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
At 1 year	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
Main Branch TLR					
At 1 month	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
At 1 year	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
Side Branch TLR					
At 1 month	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
At 1 year	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
Definite or Probable Stent Thrombosis					
At 1 month	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
At 1 year	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
Improvement in Angina Status					
At 1 month	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx
At 1 year	n	xx	xx		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (xx.x; xx.x)	0.xxxx

n : number of events;CI: confindence interval.

*: as defined by Consensus statement from the European Association of Percutaneous Cardiovascular Interventions (EAPCI) (European Journal of Cardio-Thoracic Surgery 52 (2017) 408–417)

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 10.1 Primary endpoint by subgroup (ITT)

Peri-Procedural MI at Discharge or 48h Post-Procedure	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	Risk difference (95% CI)[£]	P-value
Overall	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
Sex (Interaction p=0.xxxx)					
Male	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
Female	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
Age (Interaction p=0.xxxx)					
< 70 years	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
≥ 70 years	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
Target Vessel Type (Interaction p=0.xxxx)					
LAD	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
Non-LAD	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
Medina Class (Interaction p=0.xxxx)					
1,1,1 & 1,0,1 & 0,1,1	n/N	xx/XX	xx/XX		

Peri-Procedural MI at Discharge or 48h Post-Procedure	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	Risk difference (95% CI)£	P-value
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
1,1,0 & 1,0,0 & 0,1,0	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
Angiographic Diameter of Side Branch by Corelab (Interaction p=0.xxxx)					
≤ 2.25 mm	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
> 2.25 mm	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
Side Branch Residual Stenosis by Corelab QCA (Interaction p=0.xxxx)					
≤ 50%	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$
> 50%	n/N	xx/XX	xx/XX		
	%, (95% CI)	xx.x% (xx.x; xx.x)	xx.x% (xx.x; xx.x)	xx.x (-∞; xx.x)	0.xxxx$

Percentages are Kaplan-Meier estimates.

£: One-sided confidence interval

$: Non-inferiority test with a margin of 6%.

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 11.1 CORELAB BASELINE QCA Statistical- Bifurcation analysis

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Stenosis localisation in lesion (% diameter stenosis)				
Proximal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Distal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Side branch	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Stenosis localisation in lesion (% stenosis >=50%)				
Proximal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Distal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Side branch	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Max- stenosis in lesion (Segm- 2 + Segm- 3 + Segm- 5) PRE				
MLD (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis (%)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Main- Branch (Segm- 2 + Segm- 3 + Segm- 5)				
MLD (mm)- PRE	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
MLD (mm)- Post	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx	xx	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis - PRE	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis - Post	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Acute gain (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Prox Main- Branch (Segm- 2))				
MLD (mm)- PRE	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
MLD (mm)- Post	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx	xx	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis - PRE	N	xx	xx	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis - Post	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Reference diameter (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Obstruction length pre (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Acute gain (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Distal Main- Branch (Segm- 3)				
MLD (mm)- PRE	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
MLD (mm)- Post	N	xx	xx	0.xxxx
	Mean	xx	xx	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Sd	xx	xx	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis - PRE	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis - Post	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Reference diameter (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Obstruction length pre (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Acute gain (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Side- Branch (Segm- 5;6)				
MLD (mm)- PRE	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
MLD (mm)- Post	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx	xx	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis - PRE	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis - Post	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Reference diameter (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Obstruction length pre (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Acute gain (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Side- Branch (Segm- 5;6)				

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
MLD (mm)- PRE	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
MLD (mm)- Post	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx	xx	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis – PRE	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis – Post	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Reference diameter (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Obstruction length pre (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Acute gain (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 11.2 CORELAB EVENT QCA Statistical- Bifurcation analysis

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Stenosis localisation in lesion (Max Diam Stenosis location : Segm 2 , Segm 3 or Segm 5)				
Proximal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Distal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Side branch	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Stenosis localisation in lesion (% stenosis >=50%)				
Proximal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Distal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Side branch	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Max- stenosis in lesion (Segm- 2 + Segm- 3 + Segm- 5)				
MLD (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis (%)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Main- Branch (Segm- 2 + Segm- 3 + Segm- 5)				
MLD(mm)-	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis –(%)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Min (mm)	xx.x	xx.x	
Prox Main- Branch (Segm- 2))				
MLD(mm)-	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis (%)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Obstruction length event (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Min (mm)	xx.x	xx.x	
Distal Main-(Segm- 3)				
MLD (mm)-	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis -	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Obstruction length event (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Side- Branch (Segm- 5;6)				
MLD (mm)-	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis –(%)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Obstruction length event (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Side- Branch (Segm- 5;6)				
MLD (mm)-	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
% Diam- Stenosis -	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	
	Range	(xx.x; xx.x)	(xx.x; xx.x)	
Obstruction length event (mm)	N	xx	xx	0.xxxx
	Mean	xx	xx	
	Sd	xx.x	xx.x	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Range	(xx.x; xx.x)	(xx.x; xx.x)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 11.3 CORELAB Qualitative statistical analysis

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Bifurcation Core CASS Segment				
Proximal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
12- LAD1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
13- LAD2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
18- LCX1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
19- LCX2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3-RCA3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Distal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
13- LAD2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
14- LAD3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
19- LCX2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
20- Marg1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
21- Marg2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
4- PDA	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
5-PLV	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side branch 2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
15- Diag 1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
16- Diag 23	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
20- Marg1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
21- Marg2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
4- PDA	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
5-PLV	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Medina Visual PM/DM/SB				0.xxxx
1,1,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
1,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,0,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,0,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Medina Visual PM/DM/SB/SB2				0.xxxx
1,1,1,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,1,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,1,0,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,1,0,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,1,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,0,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,0,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,1,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,0,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Angulation(Pre- Angle DM/SB (°))				0.xxxx
<=70	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
>120	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
>70 and <=120	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Timi Flow pre				
Proximal main				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Distal main				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch 2				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Timi Flow post				
Proximal main				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Distal main				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch 2				0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Thrombus pre				
Proximal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Distal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Side Branch	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Side Branch 2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Calcification	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Proximal main-				0.xxxx
	Non/Mild	xx/XX (xx.x%)	xx/XX (xx.x%)	
	Mod	xx/XX (xx.x%)	xx/XX (xx.x%)	
	Severe	xx/XX (xx.x%)	xx/XX (xx.x%)	
Distal main-				0.xxxx
	Non/Mild	xx/XX (xx.x%)	xx/XX (xx.x%)	
	Mod	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Severe	xx/XX (xx.x%)	v	
Side Branch-				0.xxxx
	Non/Mild	xx/XX (xx.x%)	xx/XX (xx.x%)	
	Mod	xx/XX (xx.x%)	xx/XX (xx.x%)	
	Severe	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch 2-				0.xxxx
	Non/Mild	xx/XX (xx.x%)	xx/XX (xx.x%)	
	Mod	xx/XX (xx.x%)	xx/XX (xx.x%)	
	Severe	xx/XX (xx.x%)	xx/XX (xx.x%)	
Dissection/ Staining (Dissection Post)				
Proximal main				0.xxxx
	0	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE A	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE B	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE C	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE D	xx/XX (xx.x%)	xx/XX (xx.x%)	
Distal main				0.xxxx
	0	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE A	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE B	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE C	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE D	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch				0.xxxx
	0	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE A	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE B	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	TYPE C	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE D	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side branch 2				0.xxxx
	0	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE A	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE B	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE C	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE D	xx/XX (xx.x%)	xx/XX (xx.x%)	
Dissection/ Staining (Others)				0.xxxx
Abrupt Closure	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
No reflow	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Distal embolization	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Perforation	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Overlap-Gap				
Overlap	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Gap	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
If overlap: localisation				0.xxxx
	1	xx	xx	
	2	xx.x	xx.x	
Techniques / Stents Position				
Proximal main				0.xxxx
	No TTT	xx/XX (xx.x%)	xx/XX (xx.x%)	
	PTCA	xx/XX (xx.x%)	xx/XX (xx.x%)	
	1 stent	xx/XX (xx.x%)	xx/XX (xx.x%)	
	>1 stent	xx/XX (xx.x%)	xx/XX (xx.x%)	
Distal main				0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	No TTT	xx/XX (xx.x%)	xx/XX (xx.x%)	
	PTCA	xx/XX (xx.x%)	xx/XX (xx.x%)	
	1 stent	xx/XX (xx.x%)	xx/XX (xx.x%)	
	>1 stent	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side branch				0.xxxx
	No TTT	xx/XX (xx.x%)	xx/XX (xx.x%)	
	PTCA	xx/XX (xx.x%)	xx/XX (xx.x%)	
	1 stent	xx/XX (xx.x%)	xx/XX (xx.x%)	
	>1 stent	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side branch 2				0.xxxx
	No TTT	xx/XX (xx.x%)	xx/XX (xx.x%)	
	PTCA	xx/XX (xx.x%)	xx/XX (xx.x%)	
	1 stent	xx/XX (xx.x%)	xx/XX (xx.x%)	
	>1 stent	xx/XX (xx.x%)	xx/XX (xx.x%)	
Technique				0.xxxx
Kissing Balloon	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Side Branch Ballon	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
POT	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Second POT (Final)	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
Analysable film	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Table 11.4 CORELAB EVENT Qualitative statistical analysis

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
Bifurcation Core CASS Segment				0.xxxx
11-LM	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
12-LAD1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
13- LAD2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
14-LAD3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
15-Diag 1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
16-Diag 2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
18 - LCX1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
19 - LCX2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
20- Marg1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
21- Marg2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1- RCA1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2- RCA2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3- RCA3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
4-PDA	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
5-PLV	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Medina				0.xxxx
1,1,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1,0,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
0,0,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,0,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,1,0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
0,1,0,1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Timi Flow				0.xxxx
Proxiaml main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Distal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch 2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
0	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
1	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Stant trombosis	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	Yes	xx/XX (xx.x%)	xx/XX (xx.x%)	
	No(%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
Dissection				0.xxxx
Proximal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
	0	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE A	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE B	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE C	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE D	xx/XX (xx.x%)	xx/XX (xx.x%)	
Distal main	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
	0	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE A	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE B	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE C	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE D	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side Branch	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
	0	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE A	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE B	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE C	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE D	xx/XX (xx.x%)	xx/XX (xx.x%)	
Side branch 2	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	0.xxxx
	0	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE A	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE B	xx/XX (xx.x%)	xx/XX (xx.x%)	
	TYPE C	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Procedural characteristic	Statistic	No SB intervention (N=XX)	SB ballooning (N=XX)	P-value
	TYPE D	xx/XX (xx.x%)	xx/XX (xx.x%)	
Classification ISR (Classification Mehran)				0.xxxx
1 = Type IA	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
2 = Type IB	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
3 = Type IC	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
4 = Type ID	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
5 = Type II	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
6 = Type III	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	
7 = Type IV	n/N (%)	xx/XX (xx.x%)	xx/XX (xx.x%)	

DocuSign Envelope ID: DF58D057-D056-468D-925A-C386A8B5FE24

Figure 12.1 Kaplan-meier curve for *cardiac mortality* (ITT)

Repeat tables 1 to 9 for SB group only, comparing SB ballooning vs Kissing Balloon Technique (KBT)

DocuSign

Certificate Of Completion

Envelope Id: DF58D057D056468D925AC386A8B5FE24
Subject: Correction: KISS_SAP_version 2.0_2023_03_15
Source Envelope:
Document Pages: 71
Certificate Pages: 5
AutoNav: Enabled
EnvelopeId Stamping: Enabled
Time Zone: (UTC+01:00) Brussels, Copenhagen, Madrid, Paris

Status: Completed

Signatures: 4
Initials: 0

Envelope Originator:
Patricia HERNANDEZ ORTIZ
7 rue du Théâtre
Massy, Essonne 91300
phernandez-ortiz@cerc-europe.org
IP Address: 92.186.219.143

Record Tracking

Status: Original
3/21/2023 2:44:38 PM

Holder: Patricia HERNANDEZ ORTIZ
phernandez-ortiz@cerc-europe.org

Location: DocuSign

Signer Events	Signature	Timestamp
Bernard CHEVALIER bchevalier@cerc-europe.org Director CERC Security Level: Email, Account Authentication (None) **Electronic Record and Signature Disclosure:** Not Offered via DocuSign	DocuSigned by: 524718D9E9AF4E5... Signature Adoption: Drawn on Device Using IP Address: 80.215.246.131 Signed using mobile	Sent: 3/21/2023 2:56:55 PM Resent: 3/23/2023 4:27:22 PM Viewed: 3/23/2023 5:20:47 PM Signed: 3/23/2023 5:21:03 PM
Kris Bogaerts kris.bogaerts@kuleuven.be Security Level: Email, Account Authentication (None) **Electronic Record and Signature Disclosure:** Accepted: 10/3/2022 8:21:20 AM ID: b84f56fc-b9cc-40f9-ac0a-264af7464665	DocuSigned by: Kris Bogaerts B352E3687BCE477... Signature Adoption: Pre-selected Style Using IP Address: 81.11.244.128	Sent: 3/21/2023 2:56:56 PM Viewed: 3/22/2023 12:49:17 PM Signed: 3/22/2023 12:51:26 PM
Peter Smits psmits@cerc-europe.org Medical Director CERC Security Level: Email, Account Authentication (None) **Electronic Record and Signature Disclosure:** Not Offered via DocuSign	DocuSigned by: Peter Smits 0945CA73022D46A... Signature Adoption: Pre-selected Style Using IP Address: 81.175.122.2	Sent: 3/21/2023 2:56:56 PM Viewed: 3/23/2023 2:41:00 PM Signed: 3/23/2023 2:41:11 PM
Stephane Cook scook@ceric-europe.com CERIC Security Level: Email, Account Authentication (None) **Electronic Record and Signature Disclosure:** Accepted: 3/22/2023 10:31:35 AM ID: b16b31e4-824d-4602-807a-5db006662a5e	DocuSigned by: C4981411F64E4D6... Signature Adoption: Drawn on Device Using IP Address: 194.230.158.33 Signed using mobile	Sent: 3/21/2023 2:56:54 PM Viewed: 3/22/2023 10:31:35 AM Signed: 3/22/2023 10:31:49 AM

In Person Signer Events	Signature	Timestamp

Editor Delivery Events	Status	Timestamp

Agent Delivery Events	Status	Timestamp

Intermediary Delivery Events	Status	Timestamp

Certified Delivery Events	Status	Timestamp

Carbon Copy Events	Status	Timestamp
Maelys RIAUDEL mriaudel@cerc-europe.org CPL CERC Security Level: Email, Account Authentication (None) **Electronic Record and Signature Disclosure:** Not Offered via DocuSign	COPIED	Sent: 3/21/2023 2:56:55 PM
Patricia HERNANDEZ ORTIZ phernandez-ortiz@cerc-europe.org CPL CERC Security Level: Email, Account Authentication (None) **Electronic Record and Signature Disclosure:** Not Offered via DocuSign	COPIED	Sent: 3/21/2023 2:56:57 PM Resent: 3/23/2023 5:21:20 PM Viewed: 3/22/2023 12:21:19 PM

Witness Events	Signature	Timestamp

Notary Events	Signature	Timestamp

Envelope Summary Events	Status	Timestamps
Envelope Sent	Hashed/Encrypted	3/21/2023 2:56:57 PM
Certified Delivered	Security Checked	3/22/2023 10:31:35 AM
Signing Complete	Security Checked	3/22/2023 10:31:49 AM
Completed	Security Checked	3/23/2023 5:21:03 PM

Payment Events	Status	Timestamps

Electronic Record and Signature Disclosure

ELECTRONIC RECORD AND SIGNATURE DISCLOSURE

From time to time, CERC Europe (we, us or Company) may be required by law to provide to you certain written notices or disclosures. Described below are the terms and conditions for providing to you such notices and disclosures electronically through the DocuSign system. Please read the information below carefully and thoroughly, and if you can access this information electronically to your satisfaction and agree to this Electronic Record and Signature Disclosure (ERSD), please confirm your agreement by selecting the check-box next to 'I agree to use electronic records and signatures' before clicking 'CONTINUE' within the DocuSign system.

Getting paper copies

At any time, you may request from us a paper copy of any record provided or made available electronically to you by us. You will have the ability to download and print documents we send to you through the DocuSign system during and immediately after the signing session and, if you elect to create a DocuSign account, you may access the documents for a limited period of time (usually 30 days) after such documents are first sent to you. After such time, if you wish for us to send you paper copies of any such documents from our office to you, you will be charged a $0.00 per-page fee. You may request delivery of such paper copies from us by following the procedure described below.

Withdrawing your consent

If you decide to receive notices and disclosures from us electronically, you may at any time change your mind and tell us that thereafter you want to receive required notices and disclosures only in paper format. How you must inform us of your decision to receive future notices and disclosure in paper format and withdraw your consent to receive notices and disclosures electronically is described below.

Consequences of changing your mind

If you elect to receive required notices and disclosures only in paper format, it will slow the speed at which we can complete certain steps in transactions with you and delivering services to you because we will need first to send the required notices or disclosures to you in paper format, and then wait until we receive back from you your acknowledgment of your receipt of such paper notices or disclosures. Further, you will no longer be able to use the DocuSign system to receive required notices and consents electronically from us or to sign electronically documents from us.

All notices and disclosures will be sent to you electronically

Unless you tell us otherwise in accordance with the procedures described herein, we will provide electronically to you through the DocuSign system all required notices, disclosures, authorizations, acknowledgements, and other documents that are required to be provided or made available to you during the course of our relationship with you. To reduce the chance of you inadvertently not receiving any notice or disclosure, we prefer to provide all of the required notices and disclosures to you by the same method and to the same address that you have given us. Thus, you can receive all the disclosures and notices electronically or in paper format through the paper mail delivery system. If you do not agree with this process, please let us know as described below. Please also see the paragraph immediately above that describes the consequences of your electing not to receive delivery of the notices and disclosures electronically from us.

How to contact CERC Europe:

You may contact us to let us know of your changes as to how we may contact you electronically, to request paper copies of certain information from us, and to withdraw your prior consent to receive notices and disclosures electronically as follows:
To contact us by email send messages to: ccorsaro@cerc-europe.org

To advise CERC Europe of your new email address

To let us know of a change in your email address where we should send notices and disclosures electronically to you, you must send an email message to us at ccorsaro@cerc-europe.org and in the body of such request you must state: your previous email address, your new email address. We do not require any other information from you to change your email address.

If you created a DocuSign account, you may update it with your new email address through your account preferences.

To request paper copies from CERC Europe

To request delivery from us of paper copies of the notices and disclosures previously provided by us to you electronically, you must send us an email to ccorsaro@cerc-europe.org and in the body of such request you must state your email address, full name, mailing address, and telephone number. We will bill you for any fees at that time, if any.

To withdraw your consent with CERC Europe

To inform us that you no longer wish to receive future notices and disclosures in electronic format you may:

i. decline to sign a document from within your signing session, and on the subsequent page, select the check-box indicating you wish to withdraw your consent, or you may;

ii. send us an email to ccorsaro@cerc-europe.org and in the body of such request you must state your email, full name, mailing address, and telephone number. We do not need any other information from you to withdraw consent.. The consequences of your withdrawing consent for online documents will be that transactions may take a longer time to process..

Required hardware and software

The minimum system requirements for using the DocuSign system may change over time. The current system requirements are found here: https://support.docusign.com/guides/signer-guide-signing-system-requirements.

Acknowledging your access and consent to receive and sign documents electronically

To confirm to us that you can access this information electronically, which will be similar to other electronic notices and disclosures that we will provide to you, please confirm that you have read this ERSD, and (i) that you are able to print on paper or electronically save this ERSD for your future reference and access; or (ii) that you are able to email this ERSD to an email address where you will be able to print on paper or save it for your future reference and access. Further, if you consent to receiving notices and disclosures exclusively in electronic format as described herein, then select the check-box next to 'I agree to use electronic records and signatures' before clicking 'CONTINUE' within the DocuSign system.

By selecting the check-box next to 'I agree to use electronic records and signatures', you confirm that:

- You can access and read this Electronic Record and Signature Disclosure; and
- You can print on paper this Electronic Record and Signature Disclosure, or save or send this Electronic Record and Disclosure to a location where you can print it, for future reference and access; and
- Until or unless you notify CERC Europe as described above, you consent to receive exclusively through electronic means all notices, disclosures, authorizations, acknowledgements, and other documents that are required to be provided or made available to you by CERC Europe during the course of your relationship with CERC Europe.