

# Affine Objects

Mohamed Hashi

March 2, 2016

# Abstract

Short summary of the contents of your thesis.

# Acknowledgements

Put your acknowledgements here.

# 1 Introduction

Hoi, dit is de introductie.

## 2 Preliminaries

Let  $Y = (X, \mathcal{T}, \mathfrak{D})$  be a ringed site. Let  $R = \Gamma(1; \mathfrak{D})$ . Let  $a, b \in X$ .

### 2.0.1 basic categorical definitions

**Notation 1.** Let  $C$  be a category.

**Definition 2** (Presheaf categories and yoneda functors). Let  $C$  be a category. Then we define

$$C^+ := [C^{\text{op}}, \text{Set}],$$

$$C^- := [C^{\text{op}}, \text{Set}^{\text{op}}] \cong [C, \text{Set}],$$

and the functors

$$h_C : C \rightarrow C^+ := X \mapsto \text{Hom}(-, X),$$

$$h^C : C \rightarrow C^- := X \mapsto \text{Hom}(X, -).$$

Both these functors are fully faithful by the Yoneda lemma.

**Definition 3** (Over/Under categories). Let  $C$  and  $C'$  be categories. Let  $F : C \rightarrow C'$  and  $Z \in C'$ . Define the category  $C_Z$  and  $C^Z$  as follows

$$\text{Obj}(C_Z) := \{(X, u) \mid X \in C, u : F(X) \rightarrow Z\},$$

$$\text{Hom}((X, u), (Y, v)) := \{f : X \rightarrow Y \mid v \circ F(f) = u\},$$

and

## 2 Preliminaries

$$\begin{aligned}\text{Obj}(C^Z) &:= \{(X, u) \mid X \in C, u : Z \rightarrow F(X)\}, \\ \text{Hom}((X, u), (Y, v)) &:= \{f : X \rightarrow Y \mid F(f)u = v\}.\end{aligned}$$

We get faithful functors  $C_Z \rightarrow C : (X, u) \rightarrow X$  and  $C^Z \rightarrow C : (X, u) \rightarrow X$ .

### 2.0.2 Presheaves

**Definition 4** (Presheaves). Let  $A$  be any category. An  $A$ -valued presheaf  $\mathfrak{F}$  is a functor  $C^{\text{op}} \rightarrow A$ . The category  $[C^{\text{op}}, A]$  of all  $A$ -valued presheaves is denoted  $C^+(A)$ . If  $A = \text{Set}$ , we will use  $C^+$ .

When it is obvious which presheaf is under consideration, then  $\mathfrak{F}(f)$  we be denoted as  $f^*$ .

**Definition 5** (Sections functor). For any  $X \in C$  define the functor

$$\Gamma(X; -) : C^+(A) \rightarrow A$$

by

$$\mathfrak{F} \rightarrow \mathfrak{F}(X).$$

Let  $L : I \rightarrow C$  be diagram and assume that  $\text{colim}_{h \circ L}$  exists in  $C^+(A)$ . Define

$$\Gamma(\text{colim}_{i \in I} L(i); -) : C^+(A) \rightarrow A$$

by

$$\mathfrak{F} \rightarrow \text{Hom}(\text{colim}_{i \in I} L(i), \mathfrak{F}) = \lim_{i \in I} \text{Hom}(L(i), \mathfrak{F}).$$

By definition of the colimit these definitions coincide when the colimits exists in  $C$ .

**Example 6.** As we will see,  $\mathcal{C}$  is (co)complete so even if  $C$  does not have a terminal object, we can still compute the 'global sections'.

**Lemma 7** (Complete). *If  $A$  complete (or cocomplete) then  $C^+(A)$  is and all sections functors commute with arbitrary limits and colimits.*

## 2 Preliminaries

*Proof.* Let  $I \rightarrow C^+(A)$  be an diagram, with  $i \mapsto \mathfrak{F}_i$ . Then the presheaves  $U \mapsto \lim_{i \in I} F_i(U)$  and  $U \mapsto \operatorname{colim}_{i \in I} F_i(U)$  are the limit and colimit of this diagram. ■

**Lemma 8** (Abelian/Grothendieck). *If  $A$  abelian (or Grothendieck) then  $C^+(A)$  is.*

*Proof.* Let  $f, g : \mathfrak{F} \rightarrow \mathfrak{G}$ . Let  $U \in C$ . The sum  $f + g$  will be defined such that all section functors are additive. Hence  $(f + g)_U = f_U + g_U$ , this completely determines  $f + g$ . Note that this makes  $f + g$  into an actual morphism of presheaves because composition is bilinear in  $A$ .

Define  $\operatorname{Ker}(f) \rightarrow \mathfrak{F}$  to be  $U \mapsto \operatorname{Ker}(f_U)$  and  $\operatorname{Coker}(f) \rightarrow \mathfrak{F}$  to be  $U \mapsto \operatorname{Coker}(f_U)$ . Since  $A$  is abelian we have  $\operatorname{Im}(f) = \operatorname{Coim}(f)$ . By the previous lemma  $C^+(A)$  has direct sums. It also has a zero object which is the presheaf  $U \mapsto 0$ .

Hence  $C^+(A)$  is abelian as defined in tag 0109 of stacks.

Assume  $A$  is also Grothendieck. Note that the section functions  $\Gamma(U; -)$  have adjoints : Then using this the family  $\{c_! G : c \in C\}$  is a small generating family. Taking the colimit over this family provides us with a generator in  $C^+(A)$ .

Exactness and taking colimits both are determined pointwise, so directed colimits aka direct limits are exact because they are in  $A$ . ■

*Remark.* So  $C^+$  is a Grothendieck category.

**Definition 9** (direct image). Let  $f : C \rightarrow D$  and  $X \in C$ . Define the direct image  $f_*$  of  $\mathfrak{F} \in C^+(A)$  to be

$$f_*(F) = \mathfrak{F} \circ f.$$

**Lemma 10** (direct image commutes with limits).  *$f_*$  commutes with limits.*

*Proof.* Let  $\mathfrak{G} = \lim_{i \in I} \mathfrak{F}_i$  be a limit of presheaves. Let  $X \in C$ . Then  $f_* \mathfrak{G}(X) = \mathfrak{G}(f(X)) = \lim_{i \in I} \mathfrak{F}_i(f(X)) = (\lim_{i \in I} f_* \mathfrak{F}_i)(X)$ . Hence  $f_*$  commutes with limits and this also holds for colimits. ■

## 2 Preliminaries

**Definition 11** (Inverse image, direct image & push-forward). Define the inverse image  $f^*$  of  $\mathfrak{F} \in D^+(A)$  to be

$$f^*(F)(X) = \operatorname{colim}_{(Y,u) \in C_X} \mathfrak{F}(Y).$$

Define the pushforward  $f_!$  of  $\mathfrak{F} \in D^+(A)$  to be

$$f_!(F)(X) = \lim_{(Y,u) \in C^X} \mathfrak{F}(Y).$$

Now  $f^*$  is left adjoint to  $f_*$  and  $f_!$  is right adjoint to  $f^*$  by construction. This is a general construction to get adjoints, which works now because the indexing categories are small and the target contains all small (co)limits.

See Stacks Tag 09YX for a different existence lemma for the push-forward.

Assume  $C$  has binary products.

**Definition 12.** Let  $j_X : C_X \rightarrow C$  be the projection. Let  $i_X : C \rightarrow C_X$  be defined by

$$V \mapsto (U \times V, p_0)$$

We will use  $F|_X = j_{X*}\mathfrak{F}$ .

**Lemma 13.** Let  $\mathfrak{F} \in C^+(A)$  and let  $\mathfrak{G} \in X^+(A)$

$$\begin{aligned} j_{X*}\mathfrak{F}(V \rightarrow X) &= \mathfrak{F}(V), \\ j_X^*\mathfrak{G}(V) &= \bigoplus_{s \in \operatorname{Hom}(V,X)} \mathfrak{F}(V \xrightarrow{s} X) \\ j_{X!}\mathfrak{G}(V) &= \mathfrak{F}(X \times V \rightarrow X) \end{aligned}$$

**Definition 14** (Internal Hom). Let  $\mathfrak{F}, \mathfrak{G} \in C^+(A)$ . Define  $F^G$  to be the presheaf  $X \mapsto \operatorname{Hom}(\mathfrak{F}|_X, \mathfrak{G}|_X)$ . For  $A = \operatorname{Set}$  or  $A = \operatorname{R-Mod}$ , we have  $F^G \in C^+(A)$ .

If we assume that the representable sheaves are  $A$ -valued, then we can also define

$$F^G(X) = \operatorname{Hom}(\mathfrak{F} \times h_X, \mathfrak{G}).$$



## 2 Preliminaries

These two definitions are equal because the following functions are inverses of each other:

$$r \mapsto (\mathcal{U} : \mathfrak{a} \mapsto r(\mathfrak{a}, k)) \text{ where } \mathcal{U} \xrightarrow{k} X,$$

$$s \mapsto (\mathcal{U} : (\mathfrak{a}, b) \mapsto s(\mathfrak{a})).$$

**Definition 15** (Monoidal structure). Let  $\mathfrak{F}, \mathfrak{G} \in C^+(A)$ . Let  $A$  have a monoidal structure. Define  $\mathfrak{F} \otimes \mathfrak{G}$  as  $X \mapsto \mathfrak{F}(X) \otimes \mathfrak{G}(X)$

**Lemma 16** (Adjunction/monoidal closed structure). *Let  $\mathfrak{F}$  be fixed. Then  $- \otimes \mathfrak{F}$  is left adjoint to  $\text{Hom}(\mathfrak{F}, -)$ .*

### 2.0.3 Topology

**Definition 17** (Sieve). A sieve on  $X \in C$  is a subpresheaf (or subobject or subfunctor) of the representable presheaf  $h_X$ . The maximal sieve will be denoted  $\max(C)$ .

**Definition 18** (Grothendieck Topology). A Grothendieck topology  $\mathcal{T}$  is a family  $\mathcal{T}(X)$  of 'covering' sieves for every  $X \in C$  with the following conditions:

- $\max(X) \in \mathcal{T}(X)$
- $f^*R \in \mathcal{T}(X')$  if  $R \in \mathcal{T}(X)$  for any  $f : X' \rightarrow X$
- if  $f^*R \in \mathcal{T}(X')$  for all  $f \in S$  with  $S \in \mathcal{T}(X)$  then  $R \in \mathcal{T}(X)$

Note that if  $f \in R$  then  $f^*R = \max(X')$ . So if  $R \subset S$  and  $R$  is covering then  $S$  is covering. Also  $R \cap S$  is covering if and only if  $R$  and  $S$  are covering.

**Definition 19** (Basis). Let  $C$  have pullbacks. A Grothendieck pretopology  $\mathcal{B}$  is a collection  $\mathcal{B}(X)$  of families  $\{f_i : X_i \rightarrow X\}$  of 'covering' morphisms for every  $X \in C$  with the following conditions:

- every isomorphism is a covering singleton family.
- (Stability) The pullback of a covering family is covering. If  $\{f_i : X_i \rightarrow X\}$  is covering and  $g : Y \rightarrow X$ , then  $\{f'_i : X_i \times_X Y \rightarrow Y\}$  is covering.
- (Transitivity) If  $\{f_i : X_i \rightarrow X\}$  is covering and  $\{f_{ij} : X_{ij} \rightarrow X_i\}$  for every  $i$ , then  $\{f_{ij} : X_{ij} \rightarrow X\}$  is covering.

## 2 Preliminaries

Generating a real topology: take any sieve containing a covering family to be a covering sieve. Any sieve is generated by itself as covering family, in this way any topology can be interpreted as a pretopology. This enables one to use the pullbacks in proofs.

**Definition 20.** A site  $(C, \mathcal{T})$  is a category  $C$  with the Grothendieck topology  $\mathcal{T}$ . If  $C$  has pullbacks, then we consider  $\mathcal{T}$  always as a pretopology.

**Definition 21** (Cocontinuous functor).

**Lemma 22.** Let  $(C, \mathcal{T}) \xrightarrow{g} (D, \mathcal{S})$ . Let  $\mathfrak{F}$  be a presheaf on  $D$ . If  $g$  is cocontinuous, then

$$g_! \mathfrak{F}^+ \cong g_! \mathfrak{F}^+.$$

*Proof.* Let  $X$  be an object. The two presheafs reduce to

$$\lim_{R \in \mathcal{S}(g(X))} \text{Hom}(R, F) \rightarrow \lim_{K \in \mathcal{T}(X)} \text{Hom}(g(K), F).$$

The poset of covering sieves on  $X$  is sent to a dense poset of  $g(X)$  so the limits are isomorphic and this isomorphism is natural. ■

### 2.0.4 Ringed sites

### 2.0.5 Sheaves

**Definition 23** (Sheaves of sets). Let  $(C, \mathcal{T})$  be a site. Let  $\mathfrak{F} \in C^+$ . A compatible family on  $X$  is a family of elements  $x_f \in \mathfrak{F}(X_f)$  indexed by a sieve  $R$  on  $X$ , where  $X_f = \text{Dom}(f)$  and such that  $g^*(x_f) = x_{fg}$ . This is the same as a morphism  $R \rightarrow \mathfrak{F}$  as presheaves.

An amalgamation of a compatible family  $(x_f)_R$  on  $X$  is an element  $x \in \mathfrak{F}(X)$  such that  $f^*(x) = x_f$ . Hence given an morphism  $X \rightarrow \mathfrak{F}$  that extends the morphism  $R \rightarrow \mathfrak{F}$  defined by the compatible family.

A presheaf that admits a unique amalgamation of every compatible family is called a sheaf. The category  $\text{Shv}(C)$  is the full subcategory on these sheaves. Let  $i$  be the inclusion functor  $\text{Shv}(C) \rightarrow C^+$ .

## 2 Preliminaries

**Definition 24** (Sheaves #2). Let  $(C, \mathcal{T})$  be a site. Let  $\mathfrak{F} \in C^+A$ . Let  $A$  be (small) complete. Define  $\mathfrak{F}(R)$  for a sieve  $R$  on  $X$  to be

$$\text{Hom}(R, F).$$

We call  $\mathfrak{F}$  a sheaf if the map

$$\mathfrak{F}(X) \rightarrow \mathfrak{F}(R)$$

is an isomorphism.

This is just shorthand notation for the above definition. There is a bijection between  $\text{Hom}(R, F)$  and matching families and the map sends a section to the unique matching family indexed by  $R$  it is an amalgamation of.

**Definition 25** (Plus construction Shapiro). Let  $(C, \mathcal{T})$  be a site. Let  $\mathfrak{F} \in C^+$ . Define the category  $I$  whose objects are pairs  $(R, \varphi)$  with  $R \in \mathcal{T}(X)$  and  $R \xrightarrow{\varphi} F$ . A morphism between  $(R, \varphi) \rightarrow (S, \phi)$  are inclusions  $R \rightarrow S$  such that  $\varphi = \phi$  restricted to  $R$ .

Then

$$\mathfrak{F}^+(X) = \lim_{(R, \varphi) \in I} F(R).$$

More concretely,  $\mathfrak{F}^+(X)$  is the set of all objects  $(R, \varphi)$  with the equivalence relation that  $(R, \varphi) \sim (S, \phi)$  if  $\varphi = \phi$  on  $R \cap S$ . Or equivalently the set all compatible families with the equivalence relation that  $(x_f)_R \sim (y_g)_S$  if  $(x_f)_{R \cap S} = (y_g)_{R \cap S}$ .

We have the map

$$\begin{aligned} \eta : \mathfrak{F}(X) &\rightarrow \prod_{f \in R\mathfrak{F}(\text{Dom}(f))} f \\ x &\mapsto (X, x). \end{aligned}$$

This defines a natural transformation from  $\text{Id}$  to  $-^+$ .

**Definition 26** (Plus construction Moerdijk).

**Lemma 27.**  $\mathfrak{F}^+$  is separated

**Lemma 28.** If  $\mathfrak{F}$  is separated then  $\mathfrak{F}^+$  is a sheaf.

**Lemma 29.** Let  $(C, S) \xrightarrow{g} (D, \mathcal{T})$ . Let  $\mathfrak{F}$  be a presheaf on  $D$ . Then

$$\mathfrak{F}^+g \cong \mathfrak{F}g^+.$$

## 2 Preliminaries

*Proof.* ■

**Definition 30** (Sheafification). Define  $\alpha(\mathfrak{F}) = \mathfrak{F}^{++} : C^+ \rightarrow \text{Shv}(C)$ . This is a left adjoint to the inclusion functor.

The functor  $\alpha(-)$  commutes with finite limits. There are two different proofs: Moerdijk and stack+Shapiro. Shapiro defines the plus with as a colimit over a directed set, hence this commutes with limits.

**Theorem 31.** *The following this are equivalent for a category  $C$ .*

- *A Grothendieck topology*
- *A full subcategory  $E \subset C^+$  such that the inclusion functor has a left adjoint that preserves finite limits.*

*Remark* (Properties).      • The restriction of a sheaf is a sheaf

### 3 Restrictive

**Definition 32** (Restrictive functor). A functor  $f : (\mathcal{C}, \mathcal{T}, \mathfrak{D}) \rightarrow (\mathcal{D}, \mathcal{S}, \mathfrak{U})$  between ringed sites is called restrictive if for every quasi-coherent module  $\mathfrak{G}$  the co-unit induces an isomorphism

$$\begin{aligned} \mathfrak{G} &\rightarrow f_* f^* \mathfrak{G}, \\ \Gamma(1; \mathfrak{G}) &\rightarrow \Gamma(1; f_* f^* \mathfrak{G}) \cong \Gamma(1; f^* \mathfrak{G}) \\ \Gamma(1; \mathfrak{G}) \otimes_{\Gamma(1; \mathfrak{D})} \Gamma(1; \mathfrak{U}) &\rightarrow f_* f^* \mathfrak{G}. \end{aligned}$$

Note that

**Definition 33** (Restrictive morphism). A morphism  $f : X \rightarrow Y \in \mathcal{C}$  is called restrictive if the induced functor

$$C_X \rightarrow C_Y$$

is restrictive.

**Example 34.** In  $\text{Sch}$ , the morphism  $\text{Spec } A_f \rightarrow \text{Spec } A$  is restrictive.

**Non-Example 35.** The open immersion  $\text{Spec } R^2 \setminus 0 \rightarrow \text{Spec } R^2$  is not restrictive. The quasi-coherent sheaf  $\wedge(\frac{R[x,y]}{xy})$  fails to satisfy the condition from the definition.

**Lemma 36** (Restrictive to affines). *If  $f : X \rightarrow \text{Spec } R$  is a restrictive open immersion, then  $X$  is affine.*

*Proof.* ■

**Non-Example 37** (Affine non-restrictive map). Both canonical inclusions  $\mathbb{A}^1 \rightarrow \mathbb{P}^1$  are not restrictive. Look at the quasi-coherent module  $\mathfrak{O}(-1)$ . There are no global sections but on every affine chart this invertible sheaf is trivial.

**Non-Example 38.** Any inclusion  $\text{Spec } \kappa(\mathfrak{p}) \rightarrow \mathbb{P}^1$  is not restrictive. Look at  $\mathfrak{O}(-1)$ .

### 3 Restrictive

**Lemma 39.** *The composition of two restrictive functors is restrictive. If the composition  $gf$  is restrictive, then  $g$  is restrictive*

*Proof.*

■

## 4 Caffine objects

### 4.0.1 introduction

Let  $(C, \mathcal{T}, \mathfrak{D})$  be a ringed site.

**Definition 40** (Caffine). Let  $a \in C$  be an object. We call  $a$  *caffine* if the adjunction  $\Gamma(1; -) \dashv \wedge(-)$  is an equivalence of categories. Or equivalently that the unit  $\eta$  and co-unit  $\epsilon$  of this adjunction are natural isomorphisms.

**Example 41** (Examples of caffine objects). The main example to keep in mind is  $\text{Spec } R \in \text{Sch}$ .

Let  $(*, \mathfrak{R})$  be a ringed space. This space is always caffine, because all presheaves are sheaves. If  $R$  is non-local, then this space is not a scheme. This is an example of a non-scheme caffine ringed space

### 4.0.2 Restrictive maps between caffine objects

**Lemma 42** (Morphism between caffines is restrictive). *Let  $b \xrightarrow{f} a \in C$  be a morphism between caffine objects, then  $f$  is restrictive.*

*Proof.* Let  $\mathfrak{F}$  be a quasi-coherent module on  $C_a$ . We have to show that  $\Gamma(a; \mathfrak{F}) \otimes_{\Gamma(a; \mathfrak{D})} \Gamma(b; \mathfrak{D}) \xrightarrow{k} \Gamma(b; \mathfrak{F})$  is an isomorphism. The map  $k$  is the adjunct of  $\mathfrak{F}(f)$  with respect to the adjunction between restricting scalars and extending scalars along the map  $\Gamma(a; \mathfrak{D}) \rightarrow \Gamma(b; \mathfrak{D})$ . More concretely, this map is

$$k : x \otimes m \mapsto \mathfrak{F}(f)(x)m.$$

#### 4 Affine objects

The argument will go as follows. First we observe that the morphism  $\epsilon_{\mathfrak{F}} : \mathfrak{F} \rightarrow \Lambda(\Gamma(1; \mathfrak{F}))$  is an isomorphism because  $a$  is affine. Second  $i_a : \Gamma(1; \mathfrak{F})(a) \rightarrow \Lambda(\Gamma(1; \mathfrak{F}))(a)$  is an isomorphism by lemma 2.1. This holds for any affine objects, so also for  $b$ . The consequence is that  $\Gamma(1; \mathfrak{F})(f) = i_b^{-1} \mathfrak{F}(f) \circ i_a$ , by naturality of the transformation  $i : \Gamma(1; \mathfrak{F}) \rightarrow \mathfrak{F}$ . Third, show that  $\Gamma(1; \mathfrak{F})(f)$  has an isomorphism as adjunct along the same extension/restriction adjunction. Call this adjunct  $k'$ . Fourth, use naturality of adjunction bijections to conclude that  $k$  must also be an isomorphism.

Since  $a$  is affine,  $\mathfrak{F} = \Lambda(\Gamma(1; \mathfrak{F}))$ . Since  $\Lambda(-) = a(-) \circ -$ , we know that  $\mathfrak{F}$  is the sheafification of the presheaf

$$\Gamma(1; \mathfrak{F}) = c \mapsto \Gamma(a; \mathfrak{F}) \otimes_{\Gamma(a; \mathfrak{D})} \Gamma(c; \mathfrak{D}).$$

Set  $M = \Gamma(a; \mathfrak{F})$ .

Define  $k' : \Gamma(a; \mathfrak{M}) \otimes_{\Gamma(a; \mathfrak{D})} \Gamma(b; \mathfrak{D}) \rightarrow \Gamma(b; \mathfrak{M})$  By  $k' : x \otimes m \mapsto M(f)(x)m$ . If you unfold the constructions, it follows that

$$k'(x \otimes m) = x \otimes m \in \Gamma(b; \mathfrak{M}) = M \otimes \Gamma(b; \mathfrak{D})$$

is actually the identity.

We will prove that A) The component at a affine object of the universal sheafification morphism is an iso  $\epsilon_{T,a}$ . Hence when  $a$  is affine then  $\Gamma(1; i)$  on is an iso. Note that  $\Gamma(1; i) : C/x \rightarrow \mathfrak{D}(x)\text{-Mod}$  is equal to  $\Gamma(x; i) : C_y \rightarrow$

B) Adjunction bijection respects composition with isos We have now that  $M(f) = i_b^{-1} \circ \Lambda(F)(f) \circ i_a$ . Let  $F$  be the bijection from the adjunction  $\Gamma(1; -) \dashv \Lambda(-)$ . Then

C) Hence  $k$  is also an iso.

Note that restricting and sheafification commute. We can first restrict our presheaf to  $C/b$  and then sheafify. The global sections component of the universal sheafification morphism will be

$$M \otimes \Gamma(b; \mathfrak{D}) \rightarrow \Gamma(b; \mathfrak{F}),$$

$$m \otimes r \mapsto mr$$

because the triangle



#### 4 Caffeine objects

$$\begin{array}{ccc}
 \Gamma(a; \mathfrak{F}) & \xrightarrow{\text{restr}} & \Gamma(b; \mathfrak{F}) \\
 \downarrow & \nearrow & \\
 \Gamma(a; \mathfrak{F}) \otimes \Gamma(b; \mathfrak{D}) & & 
 \end{array}$$

must commute by naturality. This is exactly the component of the unit  $\eta$  of the  $\Gamma(1; -) \dashv \Lambda(-)$  on  $(C, \mathcal{T})\mathcal{O}/b$  for  $\Gamma(a; \mathfrak{F}) \otimes \Gamma(b; \mathfrak{D})$ . Since  $b$  is caffeine, this is an isomorphism by assumption.

Let  $(C, \mathcal{T}, \mathfrak{D})$  be a ringed site. Let  $R = \Gamma(1; \mathfrak{D})$ . Let  $a \in C$  and  $f : x \rightarrow y \in C$ . ■

**Lemma 43** (Co-unit is iso when locally iso). *Let  $a \in C$  be an object. Let  $\{b_i \rightarrow a\}$  be a restrictive caffeine cover, then  $\epsilon$  is a natural isomorphism.*

*Proof.* Assume we have a restrictive caffeine cover  $\{b_i \rightarrow a\}$ . Let  $\mathfrak{F}$  be a quasi-coherent sheaf module. Set  $M = \Gamma(a; \mathfrak{F})$ . Set  $M_i = \Gamma(b_i; \mathfrak{F})$  on  $C/a$ .

Consider the co-unit at  $\mathfrak{F}$

$$\epsilon_a(\mathfrak{F}) : \Lambda(\Gamma(a; \mathfrak{F})) \rightarrow \mathfrak{F}.$$

This morphism restricted gives

$$\epsilon_{a, b_i}(\mathfrak{F}) : \Lambda(\Gamma(a; \mathfrak{F}))|_{b_i} \rightarrow \mathfrak{F}|_{b_i},$$

which is the same map as  $\epsilon_b(\mathfrak{F}|_b)$ . We only need to establish that  $\Lambda(\Gamma(a; \mathfrak{F}))|_{b_i}$ .

Because  $b_i$  is caffeine, the canonical morphism given by sheafification

$$M \otimes_{\Gamma(a; \mathfrak{D})} \Gamma(b_i; \mathfrak{D}) \rightarrow M_i$$

is an isomorphism.

Hence the component

$$\epsilon_{b_i}(\mathfrak{F}|_{b_i}) : \Lambda(\Gamma(b_i; \mathfrak{F})) \rightarrow \mathfrak{F}|_{b_i}.$$

is an isomorphism, because  $\epsilon_{b_i}$  is a natural isomorphism because  $b_i$  is caffeine. The subscript  $b_i$  signifies that we are working in  $C/b_i$ .

#### 4 Caffine objects

But over an caffine object, a map is an isomorphism if and only if it is an isomorphism on global sections. In this case, using naturality of  $\epsilon$ ,

$$\epsilon_{b_i}(b_i) : \Gamma(1; \mathfrak{F}) \otimes_{\Gamma(a; \mathfrak{D})} \Gamma(b_i; \mathfrak{D}) \rightarrow \Gamma(b_i; \mathfrak{F}),$$

$$m \otimes r \rightarrow mr.$$

By restrictiveness of the map  $b_i \rightarrow a$ , this map is an isomorphism. A local isomorphism between sheaves is an isomorphism. ■

#### 4.0.3 Caffine = Affine for schemes

Let  $(X, \mathfrak{D})$  be a caffine scheme. Let  $X \xrightarrow{F} \text{Spec } \Gamma(1; \mathfrak{D}) = Y$  be the adjunct of the identity map via the adjunction  $(\text{Spec } -, \Gamma(1; -))$ . Equivalently, let  $F$  be the component at  $X$  of the unit from this adjunction.

Let's introduce our variables. Let  $x, y \in X$ . Let  $\mathfrak{p}, I, J \subset \mathfrak{D}(X)$  be ideals with  $\mathfrak{p}$  prime. Let  $a, b \in \mathfrak{D}(X)$  be global sections.

**Definition 44.** Define

$$\ker(x) = \ker(\mathfrak{D}(X) \rightarrow \kappa(x)),$$

$$D_X(a) = \{x \in X \mid a \notin \ker(x)\},$$

Define

$$V_X(I) = \text{Supp}(\wedge_X(\frac{\mathfrak{D}(X)}{I})).$$

*Remark.* Recall that  $F(x) = \ker(x)$  for  $x \in X$ . I will use  $D_Y(a)$  for the distinguished open defined by  $a$  in the affine  $Y$ . Note that  $D_X(ab) = D_X(a) \cap D_X(b)$  since  $\ker(x)$  is a prime ideal.

*Remark.* If the support of a sheaf  $\mathfrak{G}$  is empty, then locally all sections are zero. Hence all sections are equal to the zero section and  $\mathfrak{G} = 0$ .

**Lemma 45.** *The set  $V_X(I)$  is closed.*

#### 4 Affine objects

*Proof.* Let  $z \in X$  and  $M$  a  $\mathfrak{D}$ -module. Assume  $z$  is in the support of  $M$ , then  $g \neq 0$  for any generating element  $g \in M_z$ .

Consider the exact sequence

$$\mathfrak{D}(X) \rightarrow \frac{\mathfrak{D}(X)}{I} \rightarrow 0.$$

The functor  $\Lambda_X$  is a left adjoint hence right exact so

$$\mathfrak{D} \xrightarrow{f} \Lambda_X\left(\frac{\mathfrak{D}(X)}{I}\right) \rightarrow 0$$

is exact. Hence the sequence

$$\mathfrak{D}_x \xrightarrow{f_x} \Lambda_X\left(\frac{\mathfrak{D}(X)}{I}\right)_x \rightarrow 0$$

is exact. The global section  $f(1)$  must generate  $\Lambda_X(\frac{\mathfrak{D}(X)}{I})$  as a module by surjectivity of  $f$ . Similarly  $f_x(1_x)$  generates  $\Lambda_X(\frac{\mathfrak{D}(X)}{I})_x$ .

Note that  $f_x(1_x) = f(1)_x$  by definition of  $f_x$ , hence  $f(1)_x$  is a generating element. Hence  $\Lambda_X(\frac{\mathfrak{D}(X)}{I})_x \neq 0$  if and only if  $f(1)_x \neq 0$ .

This implies  $V_X(I) = \text{Supp}(f(1))$  which makes  $V_X(I)$  closed as the support of a global section. ■

The functor  $\Lambda_X$  is exact, so it commutes with quotients. So

$$\Lambda_X\left(\frac{\mathfrak{D}(X)}{I}\right) = \frac{\mathfrak{D}}{\Lambda_X(I)}$$

and

$$\Lambda_X\left(\frac{\mathfrak{D}(X)}{I}\right)_x = \frac{\mathfrak{D}_x}{\Lambda_X(I)_x} = \frac{\mathfrak{D}_x}{I \otimes \mathfrak{D}_x}$$

$\frac{\mathfrak{D}_x}{\Lambda_X(I)_x} \neq 0$ , which is the same as saying that  $\Lambda_X(I)_x$  is a proper ideal of  $\mathfrak{D}_x$ . The sheaf  $\Lambda_X(I)_x$  is the sheafification of the presheaf  $(U \mapsto I \otimes \mathfrak{D}(U))$ , hence the stalk at  $x$  of the sheaf is  $\text{colim}_{x \in U} I \otimes \mathfrak{D}(U)$ . The functor  $I \otimes -$  is a left adjoint, hence commutes with colimits. So the stalk is isomorphic to  $I \otimes \text{colim}_{x \in U} \mathfrak{D}(U) = I \otimes \mathfrak{D}_x$ . See Stacks[01BH].

**Lemma 46.** *For  $x \in X$  TFAE:*

#### 4 Caffine objects

1.  $x \in V_X(I)$
2.  $I\mathfrak{O}_x \neq \mathfrak{O}_x$
3.  $I \subset \ker(x)$ .

*Proof.* 1  $\Rightarrow$  2:

Assume  $x \in V_X(I)$ . Then  $\Lambda_X(\frac{\mathfrak{O}_x}{I})_x = \frac{\mathfrak{O}_x}{I\mathfrak{O}_x} \neq 0$ . Hence  $I\mathfrak{O}_x \neq \mathfrak{O}_x$ .

2  $\Rightarrow$  3:

Assume  $I\mathfrak{O}_x \neq \mathfrak{O}_x$ . Then  $I\mathfrak{O}_x$  is proper hence contained in the unique maximal ideal of the local ring  $\mathfrak{O}_x$ , therefore  $I \mapsto 0$  in  $k(x)$  or equivalently  $I \subset \ker(x)$ .

3  $\Rightarrow$  1:

Assume  $I \subset \ker(x)$ . Then  $I$  maps into  $\mathfrak{m}_x$ , hence  $I\mathfrak{O}_x \subset \mathfrak{m}_x$ . Therefore

$$\frac{\mathfrak{O}_x}{\Lambda_X(I)_x} = \frac{\mathfrak{O}_x}{I\mathfrak{O}_x} \neq 0.$$

■

**Corollary 47.** *If  $y \in I$  then  $D_X(y) \cap V_X(I) = \emptyset$*

*Proof.* Assume  $y \in I$ . Let  $z \in V_X(I)$ , then  $y \in \ker(z)$  by the previous lemma. This implies  $z \notin D_X(y)$  ■

**Corollary 48.**  $V_X(I) \cup V_X(J) = V_X(IJ)$

*Proof.* Let  $z \in V_X(I) \cup V_X(J)$ . Then  $I \subset \ker(z)$  and  $J \subset \ker(z)$  by the lemma, hence  $IJ \subset \ker(z)$ . Apply the lemma again to get  $z \in V_X(IJ)$ . Let  $z \in V_X(IJ)$ . Then  $IJ \subset \ker(z)$  by the lemma. The ideal  $\ker(z)$  is prime, so  $I \subset \ker(z)$  or  $J \subset \ker(z)$ . Invoke the lemma again to get  $z \in V_X(I) \cup V_X(J)$ . ■

**Lemma 49.** *Every closed set  $W$  can be written as  $V_X(I)$  for some ideal  $I$ .*

*Proof.* Let  $\mathfrak{I}$  be some ideal sheaf inducing a closed subscheme structure on  $W$ . Let  $\mathfrak{O}_W$  be the structure sheaf of this closed subscheme. By construction  $V_X(I)$  is the support of the push-forward of  $\mathfrak{O}_W$ , hence  $V_X(I) = W$ . ■

#### 4 Affine objects

**Lemma 50.** *The sets  $D_X(a)$  form a basis for the topology of  $X$ .*

*Proof.* Let  $U \subset X$  be any open. Let  $x \in U$ . By the previous lemma we get  $I$  such that  $V_X(I) = U^c$ . It follows that  $x \notin V_X(I)$  and  $I \not\subset \ker(x)$ . So we get a  $g \in I$  with  $g \notin \ker(x)$ . We get  $x \in D_X(g)$  and by corollary ..  $D_X(g) \subset U$ . As stated earlier,  $D_X(ab) = D_X(a) \cap D_X(b)$  since  $\ker(x)$  is a prime ideal. So  $D_X(a)$  form a basis. ■

**Lemma 51.** *The map  $F$  is surjective.*

*Proof.* Let  $p \in Y$  be a point in the target of  $F$ . Then  $\Lambda_X(\kappa(p))$  is a quasi-coherent sheaf of modules. In fact  $\kappa(p) \otimes_{\Gamma} \mathfrak{O}(U)$  is a  $\mathfrak{O}(U)$  algebra, hence  $\Lambda_X(\kappa(p))$  is a quasi-coherent sheaf of algebras. Hence we can compute the relative spec  $\text{Rspec}(\Lambda_X(\kappa(p))) \rightarrow X$ . The adjunct of the map

$$\text{Rspec}(\Lambda_X(\kappa(p))) \rightarrow Y$$

is the canonical morphism  $g : \text{R} \rightarrow \kappa(p)$ . This morphism is also the adjunct of the composition

$$\text{Rspec}(\Lambda_X(\kappa(p))) \rightarrow \text{Spec } \kappa(p) \rightarrow X,$$

so both maps must be equal. This gives us a commutative square

$$\begin{array}{ccc} X & \xrightarrow{F} & Y \\ \uparrow & & \uparrow \\ \text{Rspec}(\Lambda_X(\kappa(p))) & \longrightarrow & \text{Spec } \kappa(p) \end{array}$$

By lemma .., we know that  $\Lambda_X(\kappa(p))$  is not the zero sheaf hence the structure sheaf of  $\text{Rspec}(\Lambda_X(\kappa(p)))$  non-zero. This implies that the scheme is not the empty scheme. Therefore the point  $p$  is in the image of  $F$ . ■

**Lemma 52.** *The closed set  $V_X(p)$  is irreducible. This implies that  $F$  is injective.*

#### 4 Affine objects

*Proof.* Let  $F(z) = p$  for some  $z \in X$ . By lemma .. this is possible. Let  $y \in V_X(p)$ . Then  $\ker(z) \subset \ker(y)$ , hence if  $y \in D_X(a)$  then  $x \in D_X(a)$ . Therefore  $y$  specialises to  $z$ , which thus must be  $V_X(p)$ . This shows that it is irreducible. Uniqueness of generic points of closed irreducible subsets of schemes implies injectivity of  $F$ . ■

**Lemma 53.** *The function  $F$  is open, hence a homeomorphism.*

*Proof.* Note that  $F(D_X(a)) = \{F(x) \mid a \notin F(x)\} = F(X) \cap D_Y(a) = D_Y(a)$ . Our map  $F$  is continuous and open, so a homeomorphism. ■

**Lemma 54.** *If  $F$  is a homeomorphism, then  $X$  is affine.*

*Proof.* Let  $\text{Spec } A_i = U_i \subset X$  be open and let  $\bigcup_i U_i = X$ . Assume it is a finite affine cover. Using our base, we get a cover of  $U_i = \bigcup_j D_X(a_{ij})$  with  $a_{ij}$  global sections. Observe that  $D_X(a_{ij}) \subset U_i$ , hence  $D(a_{ij}|_{U_i}) = D_X(a_{ij})$  which makes them affine. Continuing like this, we get a finite cover of affines  $D_X(a_{ij})$  of  $X$ . since  $F(X) = F(\bigcup_{ij} D_X(a_{ij})) = \bigcup_{ij} D_Y(a_{ij}) = Y$ , we have  $(a_{ij}) = (1)$ . Affine-ness satisfies the two requirements for the affine communication lemma[HAG II Ex.2.17], hence  $X$  is affine. ■

#### 4.0.4 P1

Quasi-coherent modules on a scheme  $X$  can be defined for affine schemes first and as module sheaves that are  $\tilde{M}$  for some  $f^*\mathcal{O}$ -module  $M$  pulled back to every affine  $\text{Spec } R \xrightarrow{f} X$ . Let's consider the following conjecture.

**Conjecture 55 (P1).** Let  $a \in C$ . A sheaf module  $\mathfrak{F}$  on  $(C, \mathcal{T}, \mathcal{O})/a$  is quasi-coherent if and only if  $\mathfrak{F}$  is quasi-coherent on  $(C, \mathcal{T}, \mathcal{O})/b$  for any affine  $b \rightarrow a$

*Remark.* The only if direction always holds.

**Definition 56.** We say that a ringed site  $(C, \mathcal{O}, \mathcal{T})$  has *enough affines* if any object has an affine covering  $\{b_i \rightarrow a\}$ .

**Lemma 57** (P1 holds with enough affines). *P1 holds for any ringed site with enough affines.*

#### 4 Affine objects

*Proof.* Quasi-coherence is a local property. So if every object admits a affine cover P1 holds. ■

**Lemma 58** (Finite poset has enough affines). *Any finite poset has enough affines.*

*NA-chain proof.* Let  $x_0 \in C$ . If  $x_0$  is covered by the maximal sieve only or the maximal sieve and the empty sieve, it is affine and we are done. Assume otherwise. Let  $S = \{y_i \rightarrow x_0\}$  be a non-maximal, non-empty cover of  $x_0$ . Then  $S$  does not contain isomorphisms.

We can associate to any non-maximal non-empty covering sieve  $S$  of an element  $x_0$ , the set of all NA-chains  $x_0 \leftarrow x_1 \leftarrow \dots \leftarrow x_n$ . An NA-chain, associated to  $R$ , is a chain of maps ending in  $x_0$  such that  $x_i \leftarrow x_{i+1}$  is contained in a non-maximal, non-empty cover of  $x_i$ , where  $x_0 \leftarrow x_1$  is contained in  $R$ .

By finiteness of  $C$ , any chain of maps is bounded by the size of  $C$  or contains a cycle. If a chain contains a cycle, it contains isomorphisms. By construction, no isomorphism can be present in a NA-chain. Therefore the length of any NA-chain is bounded by  $\|C\|$ .

Let  $H$  be a NA-chain associated to  $S$  of maximal length  $m$ . Then the last map  $\dots \leftarrow h \leftarrow g$  in  $H$  has an affine object  $g$  as domain, because  $H$  cannot be increased and so  $g$  has no non-maximal, non-empty coverings which makes it affine. Also the non-maximal, non-empty covering of  $h$  where this map appears must be an affine covering by applying the same reasoning to the other objects occurring in it. Hence all objects occurring at the  $(m-1)$ th place in any NA-chain admits an affine cover. Let  $i \leq m-1$ . Assume all elements at the  $(i-1)$ th place admit an affine cover. Let  $b$  be a object occurring at the  $(i-1)$ th place in a chain. It is either affine or all objects in any non-maximal, non-empty cover occur at the  $i$ th place in some chain hence admit an affine cover. Therefore any non maximal, non empty cover on  $b$  can be refined to an affine cover. This provides us with an affine cover of  $b$ . By reversed induction,  $x_0$  admits a affine cover. ■

**Lemma 59** (Non-quasi-coherent sheaf). *Any category admits a non-quasi-coherent sheaf.*

*Proof.* Let  $C$  be a ringed site with no affines. Therefore no object can have the empty sieve as a covering sieve, because that would make all sheafs trivial localized at this object. Let  $\mathcal{O}$  be its structure sheaf.

Let  $a$  be an object of  $C$ . Let  $b$  be an object of  $C$  such that  $\text{Hom}(b, a) = \emptyset$ .

#### 4 Affine objects

The following situation, the commuting square with conditions on the maps and objects, will be called S1. Note that  $a, b$  are fixed and not variables in S1.

$$\begin{array}{ccc} e & \xrightarrow{L} & b \\ \uparrow K & & \uparrow F \\ d & \xrightarrow{G} & c \end{array}$$

With  $\mathfrak{D}(d) \neq 0$ ,  $\text{Hom}(e, a) = \emptyset$  and  $\text{Hom}(c, a) \neq \emptyset$ .

Assume that for any  $S$  a covering sieve on  $b$ .

- 1) every map  $F \in S$  as in S1, so  $F$  has codomain  $b$  and its domain maps to  $a$ , we can find maps  $G, K, L$  to complete to S1 with  $L \in S$ .
- 2) For every  $L \in S$  as in S1, so  $L$  has codomain  $b$  and its domain does not map to  $a$ , we can complete the to get S1.

Consequences: Every non-empty covering sieve of  $b$  contains maps  $L$  and  $F$  that fit in S1. 'Objects under  $a$  get under every object under  $b$ '. Call this assumption A1 en A2.

Define the presheaf  $\mathfrak{F}$  as

$$\begin{aligned} x &\mapsto \mathfrak{D}(x) \text{ if } \text{Hom}(x, a) = \emptyset, \\ x &\mapsto \mathfrak{D}(x)[y] \text{ otherwise ,} \end{aligned}$$

$$\begin{aligned} u \xrightarrow{f} v &\mapsto \mathfrak{D}(f) \text{ if } \text{Hom}(x, a) = \emptyset, \\ (u \xrightarrow{f} v) &\mapsto (\mathfrak{D}(u) \rightarrow \mathfrak{D}(u)[y]) \circ \mathfrak{D}(f) \text{ otherwise .} \end{aligned}$$

Let  $G = \mathfrak{F}^{++}$ .

Let  $S$  be a covering sieve on  $b$ . If  $S$  is empty, then  $G(b) = 0 = \mathfrak{D}(b)$ . Assume otherwise. Let  $(x_f)$  be a matching family indexed by  $S$ . Let  $x_f \in G(u) \setminus O(u)$  for  $u \xrightarrow{f} b$  such that  $\text{Hom}(u, a) \neq \emptyset$ , which is possible by A2. Set  $F = f$  and complete to S1. Then



#### 4 Affine objects

$\mathfrak{F}(G)(x_f) = \mathfrak{F}(K)(x_L) = x_{KL}$  since it is a matching family, which is impossible because  $\mathfrak{F}(G)(x_f)$  is not in the image of  $\mathfrak{F}(K)$ . So all matching families  $(x_f)$  have components that are elements of  $O(\text{Dom}(f)) \subset G(\text{Dom}(f))$ , which already have unique amalgamations in  $\mathfrak{D}(b)$ . Hence  $G(b) = \mathfrak{D}(b)$ .

Let  $\mathcal{U}$  be a global cover of the category. By A1,  $\mathcal{U}(b)$  contains an element  $g : e \rightarrow b$  with  $\text{Hom}(e, a) = \emptyset$ . Set  $g = L$  and complete S1. The element  $y \in G(d)$  is not generated locally by sections of  $G(e)$ . Hence  $G$  is not locally presentable.

*Examples that satisfy A1&A2*

- Open category of any irreducible space.
- Neighbourhood space of any point in any topological space.
- categories with pullbacks, terminal object and are irreducible.

■

**Example 60** (Stacks 01BL example). Let  $L = (\mathbb{R}, O_{\mathbb{R}})$  be the real line with the euclidean topology and the sheaf of continuous real valued functions as structure sheaf. Let

$$X = \frac{\bigcup_{i=0}^{\infty} L_i}{\sim}$$

with  $[i, x] \sim [j, y]$  if and only if  $i = j$  and  $x = y$  or  $y = x = 0$ . The real lines are glued to each other at zero. Define the open  $U_n \subset X$  as  $U_n \cap L_i = (-\frac{1}{n}, \frac{1}{n})$ . These opens form a basis of neighbourhoods of 0. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be any continuous function such that  $f(x) = 0$  if  $x \in (-1, 1)$  and  $f(x) = 1$  if  $x \in (-\infty, -2) \cup (2, \infty)$ . Let  $f_n(x) = f(nx)$ .

Define the sheaf map

$$\bigoplus_i O_{\mathbb{R}} \xrightarrow{\alpha} \bigoplus_{ij} O_{\mathbb{R}},$$

$$e_i \mapsto \sum_j f_i 1_{L_j} e_{ij}.$$

To proof that this is well-defined, we need to show that the sum  $\sum_j f_i 1_{L_j} e_{ij}$  is locally finite for every  $i$ . Let  $[k, y] \in X$ . If  $y \neq 0$ , then

$$W_{[k,y]} = \{[k, z] \in X \mid z \in (y - \delta, y + \delta) \subset L_k\}$$

#### 4 Affine objects

is open in  $X$  and  $\alpha_{W_{[k,y]}}(e_i) = f_i e_{ik}$  for any  $\delta < |\frac{y}{2}|$ . If  $y = 0$ , then  $\alpha_{U_n}(e_i) = 0$  if  $n > i$  because  $f_i$  is zero on  $U_n$ . Hence we found a cover on which our sum is locally finite, which makes  $\alpha$  well-defined.

Let  $U$  be any open of any topological space  $X$ . Let  $\mathfrak{F}$  be any presheaf. Consider the map of presheafs

$$\mathfrak{F}|_U \xrightarrow{+} \mathfrak{F}^+|_U$$

defined by the components

$$g_V : \operatorname{colim}_{S \in \operatorname{Cov}(V)} \operatorname{Match}(S, F) \xrightarrow{\operatorname{id}} \operatorname{colim}_{S \in \operatorname{Cov}(V)} \operatorname{Match}(S, F).$$

Every component is a isomorphism, hence  $g$  is an isomorphism.

The adjunction  $(\Lambda(-), \Gamma(X, -))$  implies that  $\Lambda(-)$  commutes with arbitrary colimits. Moreover

$$O_X \cong \Lambda(\Gamma(X, O_X))$$

so

$$\bigoplus_i O_R \cong \Lambda(\bigoplus_i \Gamma(X, O_X)).$$

This shows that  $\alpha$  is a morphism between associated sheafs. Let  $\beta : \bigoplus_i \Gamma(U, O_X) \rightarrow \bigoplus_{ij} \Gamma(U, O_X)$  for some open  $U$ . Then  $\Lambda(\beta)(e_i) = \sum_{j \in J_i} a_{ij} e_{ij}$  where  $J_i$  is finite for every  $i$ .

Assume that  $\alpha = \Lambda(\beta)$  over some neighbourhood  $U$  of 0. Then there exists a  $m$  such that  $U_m \subset U$ . Let  $k > 2m$ . Then  $f_k \neq 0$  on  $U_m$ , hence  $f_k 1_{L_j} \neq 0$  on  $U_m$  for every  $j$  and so no coefficients vanish of  $\alpha_{U_m}(e_k) = \sum_j f_k 1_{L_j} e_{kj}$ . This contradicts  $\alpha = \Lambda(\beta)$ .

**Example 61** (Category without enough affines #1). Let  $X, f_j$  and  $U_n$  be as in the previous example. Define the full subcategory  $N(y) \xrightarrow{i} \operatorname{Open}((X))$  of all opens  $U$  that contain the point  $y \in X$ .

This category has all fibre products. Let  $U \rightarrow V \leftarrow W$  be two morphisms. Then  $U \leftarrow U \cap W \rightarrow W$  is the pullback.

#### 4 Affine objects

On this category  $N(y)$ , let a family  $\{f_i : U_i \rightarrow U\}$  be covering if  $\bigcup_i f_i(U_i) = U$ .

Let  $V \xrightarrow{f} U$  be an isomorphism, then  $f(V) = U$  so  $\{f\}$  is a covering family.

Let  $\{U_i \xrightarrow{f_i} U\}$  be a covering family. For every  $i$ , let  $\{U_{ij} \xrightarrow{f_{ij}} U_i\}$  be a covering family. By definition this gives that  $\bigcup_i f_i(U_i) = U$  and  $\bigcup_j f_{ij}(U_{ij}) = U_i$  for every  $i$ . Hence

$$\bigcup_{i,j} (f_i \circ f_{ij})(U_{ij}) = \bigcup_i f_i(U_i) = U$$

and so the family  $\{U_{ij} \xrightarrow{f_i \circ f_{ij}} U\}$  is covering.

Let  $V \rightarrow U$  be a morphism in  $N(y)$  and  $\{U_i \xrightarrow{f_i} U\}$  be a covering family on  $U$ . This tells us that  $\bigcup_i f_i(U_i) = U$ , hence also  $\bigcup_i g_i(U_i \cap V) = V$  where  $g_i : U_i \cap V \rightarrow V$  is the pullback of  $f_i$ . Hence  $\{U_i \cap V \xrightarrow{g_i} V\}$  is a covering family of  $V$ .

All criteria for a pretopology are established. Let  $\tau$  be the generated Grothendieck topology.

Let  $\mathfrak{F}$  be a sheaf on  $\text{Open}(X)$ . Let  $\hat{\mathfrak{F}} = \mathfrak{F} \circ i$ . Let  $\{U_i \rightarrow V\}$  be a covering family on  $V$  in  $N(y)$ . Let  $(x)_i$  be a matching family of  $\hat{\mathfrak{F}}$  indexed by  $\{U_i \rightarrow V\}$ , so  $x_i \in \hat{\mathfrak{F}}(U_i) = \mathfrak{F}(U_i)$ . Note that  $\{U_i \rightarrow V\}$  is also a covering family on  $V$  in  $\text{Open}(X)$ , hence  $(x)_i$  is also a matching family of  $\mathfrak{F}$  on  $V$ . Since  $\mathfrak{F}$  is a sheaf, there exists a unique amalgamation  $x \in \mathfrak{F}(V) = \hat{\mathfrak{F}}(V)$  such that  $x = x_i$  in  $\mathfrak{F}(U_i) = \hat{\mathfrak{F}}(U_i)$ . This shows that  $\hat{\mathfrak{F}}$  is a sheaf, hence  $i$  is continuous.

Let  $\mathfrak{O}_{X,y} = \mathfrak{O}_X \circ \tau$ . This is a sheaf of rings by the previous. We constructed a ringed site  $(N(y), \tau, \mathfrak{O}_{X,y})$ .

Let  $F$  be the inclusion functor from the category of sheafs to the category of presheafs. We have the adjunctions:

1.  $(\wedge(-), \Gamma(X, -))$ ,
2.  $((-)^{++}, F)$ .

The structure sheaf  $\mathfrak{O}_{X,y}$  is, trivially, isomorphic to  $\wedge(\Gamma(X, \mathfrak{O}_{X,y}))$ . By adjunction (2)

$$\bigoplus_i \mathfrak{O}_{X,y} \cong (\bigoplus_i \mathfrak{O}_{X,y})^{++},$$

#### 4 Affine objects

where the coproduct on the left hand side is in the category of sheafs and the coproduct on the right hand side in the category of presheafs.

By adjunction (1)

$$\bigoplus_i \Lambda(\Gamma(X, \mathcal{O}_{X,y})) \cong \Lambda(\bigoplus_i \Gamma(X, \mathcal{O}_{X,y})).$$

Combine these 3 observation to get

$$\bigoplus_i \mathcal{O}_{X,y} \cong \Lambda(\bigoplus_i \Gamma(X, \mathcal{O}_{X,y})),$$

which shows that  $\bigoplus_i \mathcal{O}_{X,y}$  is quasi-coherent.

Set  $y = 0 \in X$ . Define the sheaf map

$$\begin{aligned} \bigoplus_i \mathcal{O}_{X,y} &\xrightarrow{\alpha} \bigoplus_{ij} \mathcal{O}_{X,y}, \\ e_i &\mapsto \sum_j f_i 1_{L_j} e_{ij}. \end{aligned}$$

Fix  $i$ . We will prove that  $\alpha_X(e_i)$  is a well-defined global section. Let  $m > i$ . Let  $V_k = U_k \cup U_m$  and  $\{V \rightarrow \bullet\} \{V_k\}$ . By construction  $f_i$  is zero on  $U_m$ , hence  $f_i 1_{L_j}$  is zero on  $V_k$  if  $k \neq j$  and so  $\sum_j f_i 1_{L_j} e_{ij} = f_i 1_{L_k} e_{ik}$  on  $V_k$ . This shows that  $\alpha_X(e_i)$  is a well-defined section on any element of the cover  $\{U_i \rightarrow V\}$  and this family is matching since the sections are functions and the 'restriction' maps are actual restriction.

Assume there exists  $\beta : \bigoplus_i \Gamma(V, \mathcal{O}_{X,y}) \rightarrow \bigoplus_{ij} \Gamma(V, \mathcal{O}_{X,y})$  such that  $\Lambda(\beta) = \alpha_V$ . Then  $\alpha_V(e_i) = \sum_j f_i 1_{L_j} e_{ij}$  is not just locally finite over some cover, but actually finite globally on  $V$  for all  $i$ . So almost all  $f_i 1_{L_j}$  are zero on  $V$ . Note that  $y \in V$ , so  $U_d \subset V$  for some  $d$ . Let  $i > 2d$ , then  $f_i \neq 0$  on  $(-\frac{1}{d}, \frac{1}{d})$  and so  $f_i 1_{L_j} \neq 0$  on  $U_d$  for any  $j$ . Hence  $\alpha_V(e_i) = \sum_j f_i 1_{L_j} e_{ij}$  is not a finite sum for  $i > 2d$ . This contradicts our assumption.

Let  $U \subset X$  be  $U \cap L_j = U_j$ . Fix  $i$ . Then  $f_i 1_{L_j} = 0$  if  $i < j$ , hence

$$\sum_j f_i 1_{L_j} e_{ij} = \sum_{j \leq i} f_i 1_{L_j} e_{ij}$$

is a finite sum.

#### 4 Affine objects

The restriction of any quasi-coherent sheaf is quasi-coherent. Observe that  $\alpha$ , and its restrictions, is a morphism between quasi-coherent sheafs but does not come from a map of modules. Therefore  $\Lambda(-)_V : \Gamma(V, \mathcal{O}_{X,Y}) - \text{Mod} \rightarrow \text{Qcoh}(\mathfrak{Y})$  is not full for any  $V$  and no object  $V$  is affine in  $N(Y)$ .

**Example 62** (Category without enough affines #1).

**Example 63** (General P1 does not hold). The category  $C$  is  $\mathbb{Z} \times \mathbb{Z}$  with the usual ordering. An element  $(i, j)$  is only non-trivially covered by  $\{(i, j-1) \rightarrow (i, j), (i-1, j) \rightarrow (i, j)\}$ . Let  $k$  be any field. Let  $R = k[x_{ij} | i, j \in \mathbb{Z}]$ . Define the structure sheaf as  $\mathcal{O}(i, j) = R[x_{kl}^{-1} | i \leq k \text{ \& } j \leq l]$ .

Fix  $(a, b) \in C$ . Consider the over category  $C \downarrow (a, b)$  at this point. Let  $(i, j) \rightarrow (a, b)$  be an object of  $C \downarrow (a, b)$ . Define the presheaf of modules  $F(i, j) = \mathcal{O}(i, j) / (x_{a-1, b} x_{a, b-1})$  on  $X$ . Then  $a > i$  or  $b > j$  or  $(i = a \text{ and } j = b)$ . If  $a > i$  or  $b > j$ , then  $x_{a-1, b}$  or  $x_{a, b-1}$  is invertible in  $\mathcal{O}(i, j)$ , hence  $F(i, j) = 0$  in both cases. This presheaf is zero everywhere except at  $(a, b)$ , hence sheafifies to the zero sheaf. In other words:  $\Lambda(\frac{\mathcal{O}(a, b)}{(x_{a-1, b} x_{a, b-1})}) = 0$ , where  $\Lambda$  is the 'tilde' functor. Hence  $(a, b)$  is not affine, which shows that  $C$  has no affine objects.

Consider  $G = \mathcal{O}(i, j)[y_{kl} | k \leq i \text{ \& } l \leq j]$ . Let  $\bigoplus_{k \in I} \mathcal{O} \xrightarrow{\alpha} G$  be any sheaf map. Let  $\alpha_{00}(e_k)$  be the image of the generators  $e_k \in \bigoplus_{k \in I} \mathcal{O}$  in the global sections. The section  $y_{1,1} \in G(1, 1)$  cannot be written as a finite sum  $\sum_k \lambda_k \alpha_{00}(e_k)$  for scalars  $\lambda_k \in \mathcal{O}(i, j)$  for any  $(i, j)$ . This shows that  $\alpha$  is not surjective hence  $G$  is not quasi-coherent (locally presentable).

## 5 Cohomology