SOEN 6011 SEP

Problem 1 Function 6: B(x,y) Beta Function Mahavir Patel 40198619

Date: 05/07/2022

Introduction 1

The beta function is a special function that belongs to the first category of Euler's integrals. The beta function is denoted by the symbol "B". B(x,y) refers to the beta function, where x and y are real-valued parameters.

The formula of B(x, y) is¹:

- $B(x,y) = \frac{(x-1)!(y-1)!}{(x+y-1)!}$ For positive integers
- $B(x,y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt$ For positive real numbers

2 **Properties of Beta Function**

- Beta function is symmetric which means its beta value is independent of the order of its parameters: B(x,y) = B(y,x)
- B(x,y) = B(x,y+1) + B(x+1,y)
- $B(x, y + 1) = B(x, y) \cdot \left[\frac{y}{x+y}\right]$ $B(x + 1, y) = B(x, y) \cdot \left[\frac{x}{x+y}\right]$
- $B(x,y).B(x+y,1-y) = \frac{\pi}{x}\sin \pi y$
- Beta function in terms of Gamma functions as: $B(x,y) = \frac{\Gamma x \Gamma y}{\Gamma(x+y)}$, When x and y are positive whole number then it follows the form of gamma function.
- The beta function can be extended to a function with more than two arguments: $B(X_1, X_2, ... X_n) = \frac{\Gamma X_1 \Gamma X_2 ... \Gamma X_n}{\Gamma (X_1 + X_2 + ... + X_n)}$

3 Domain and Co-Domain

The domains of real numbers are where the beta function is defined. The limitations of the integral function determine the beta function's co-domain. the beta function is defined For real values that are positive and greater than zero. nonetheless, there are other ways to write a beta function, for example,

- $B(x,y) = \int_0^{\frac{\pi}{2}} (\sin \theta)^{2x-1} (\cos \theta)^{2x-1}$ where x > 0 and y > 0
- $B(x,y) = \int_0^\infty \frac{t^{x-1}}{(1+t)^{x+y}} dt$ where x > 0 and y > 0

for the variables domain is $(0,\infty]$ and based on the beta function's integral limits codomain can be defined.

¹https://en.wikipedia.org/wiki/Beta_function

Context of Use Model




Figure 1: Context of use model for a user.