

Lecture 13 - Planar Graphs and Graph coloring

April 18, 2019

Recap

- Graphs – Connectivity
- Euler and Hamilton Circuits and Paths

Planar Graphs

Planar Graphs

- The motivation – to design connections where the connecting edges (wires?) do not criss cross each other.

Planar Graphs

- The motivation – to design connections where the connecting edges (wires?) do not criss cross each other.
- Important in designing electronic circuits!

Planar Graphs

- The motivation – to design connections where the connecting edges (wires?) do not criss cross each other.
- Important in designing electronic circuits!
- Examples –

Planar Graphs

- The motivation – to design connections where the connecting edges (wires?) do not criss cross each other.
- Important in designing electronic circuits!
- Examples –
- A graph is called **planar** if it can be drawn in the plane without any edges crossing (where a crossing of edges is the **intersection of the lines or arcs representing them at a point other than their common endpoint**).

Planar Graphs

- The motivation – to design connections where the connecting edges (wires?) do not criss cross each other.
- Important in designing electronic circuits!
- Examples –
- A graph is called **planar** if it can be drawn in the plane without any edges crossing (where a crossing of edges is the **intersection of the lines or arcs representing them at a point other than their common endpoint**).
- Such a drawing is called planar representation of the graph.

Planar Graphs

- The motivation – to design connections where the connecting edges (wires?) do not criss cross each other.
- Important in designing electronic circuits!
- Examples –
- A graph is called **planar** if it can be drawn in the plane without any edges crossing (where a crossing of edges is the **intersection of the lines or arcs representing them at a point other than their common endpoint**).
- Such a drawing is called planar representation of the graph.
- We just saw K_4 and Q_3 are planar. What about $K_{3,3}$ and K_5 ?

Planar Graphs

- The motivation – to design connections where the connecting edges (wires?) do not criss cross each other.
- Important in designing electronic circuits!
- Examples –
- A graph is called **planar** if it can be drawn in the plane without any edges crossing (where a crossing of edges is the **intersection of the lines or arcs representing them at a point other than their common endpoint**).
- Such a drawing is called planar representation of the graph.
- We just saw K_4 and Q_3 are planar. What about $K_{3,3}$ and K_5 ?
No.

Planar Graphs

- The motivation – to design connections where the connecting edges (wires?) do not criss cross each other.
- Important in designing electronic circuits!
- Examples –
- A graph is called **planar** if it can be drawn in the plane without any edges crossing (where a crossing of edges is the **intersection of the lines or arcs representing them at a point other than their common endpoint**).
- Such a drawing is called planar representation of the graph.
- We just saw K_4 and Q_3 are planar. What about $K_{3,3}$ and K_5 ?
No.
- We can use arguments that look at regions divided and then argue that they cannot be planar.

Euler's Formula

A planar representation of a graph splits the plane into regions, including an unbounded region.

Theorem

Let G be a connected planar simple graph with e edges and v vertices. Let r be the number of regions in a planar representation of G . Then $r = e - v + 2$.

Euler's Formula

A planar representation of a graph splits the plane into regions, including an unbounded region.

Theorem

Let G be a connected planar simple graph with e edges and v vertices. Let r be the number of regions in a planar representation of G . Then $r = e - v + 2$.

- First we specify a planar representation of G .

Euler's Formula

A planar representation of a graph splits the plane into regions, including an unbounded region.

Theorem

Let G be a connected planar simple graph with e edges and v vertices. Let r be the number of regions in a planar representation of G . Then $r = e - v + 2$.

- First we specify a planar representation of G .
- We construct a sequence of subgraphs $G_1, G_2, \dots, G_e = G$ by adding an edge at each stage (it has to be incident with a vertex in G_{n-1}).

Euler's Formula

A planar representation of a graph splits the plane into regions, including an unbounded region.

Theorem

Let G be a connected planar simple graph with e edges and v vertices. Let r be the number of regions in a planar representation of G . Then $r = e - v + 2$.

- First we specify a planar representation of G .
- We construct a sequence of subgraphs $G_1, G_2, \dots, G_e = G$ by adding an edge at each stage (it has to be incident with a vertex in G_{n-1}).
- G_1 will have one edge (some arbitrary edge!)

Euler's Formula

A planar representation of a graph splits the plane into regions, including an unbounded region.

Theorem

Let G be a connected planar simple graph with e edges and v vertices. Let r be the number of regions in a planar representation of G . Then $r = e - v + 2$.

- First we specify a planar representation of G .
- We construct a sequence of subgraphs $G_1, G_2, \dots, G_e = G$ by adding an edge at each stage (it has to be incident with a vertex in G_{n-1}).
- G_1 will have one edge (some arbitrary edge!)
- Let r_n, e_n, v_n represent the number of regions, edges, and vertices of the planar representation of G_n .

Euler's Formula - Proof by induction

- $r_1 = e_1 - v_1 + 2$ is true for G_1 since $e_1 = r_1 = 1$ and $v_1 = 2$.

Euler's Formula - Proof by induction

- $r_1 = e_1 - v_1 + 2$ is true for G_1 since $e_1 = r_1 = 1$ and $v_1 = 2$.
- Assume $r_k = e_k - v_k + 2$.

Euler's Formula - Proof by induction

- $r_1 = e_1 - v_1 + 2$ is true for G_1 since $e_1 = r_1 = 1$ and $v_1 = 2$.
- Assume $r_k = e_k - v_k + 2$.
- Let $\{a_{k+1}, b_{k+1}\}$ be the edge that is added to G_k to obtain G_{k+1} .

Euler's Formula - Proof by induction

- $r_1 = e_1 - v_1 + 2$ is true for G_1 since $e_1 = r_1 = 1$ and $v_1 = 2$.
- Assume $r_k = e_k - v_k + 2$.
- Let $\{a_{k+1}, b_{k+1}\}$ be the edge that is added to G_k to obtain G_{k+1} .
- Case i: The vertices were already in G_{k+1} .

Euler's Formula - Proof by induction

- $r_1 = e_1 - v_1 + 2$ is true for G_1 since $e_1 = r_1 = 1$ and $v_1 = 2$.
- Assume $r_k = e_k - v_k + 2$.
- Let $\{a_{k+1}, b_{k+1}\}$ be the edge that is added to G_k to obtain G_{k+1} .
- Case i: The vertices were already in G_{k+1} .
- They must have been on the boundary of a common region R or else $\{a_{k+1}, b_{k+1}\}$ would have caused crossing!

Euler's Formula - Proof by induction

- $r_1 = e_1 - v_1 + 2$ is true for G_1 since $e_1 = r_1 = 1$ and $v_1 = 2$.
- Assume $r_k = e_k - v_k + 2$.
- Let $\{a_{k+1}, b_{k+1}\}$ be the edge that is added to G_k to obtain G_{k+1} .
- Case i: The vertices were already in G_{k+1} .
- They must have been on the boundary of a common region R or else $\{a_{k+1}, b_{k+1}\}$ would have caused crossing!
- The new edge splits R into two regions $\Rightarrow r_{k+1} = r_k + 1$,
 $e_{k+1} = e_k + 1$, $v_{k+1} = v_k$ - formula holds!

Euler's Formula - Proof by induction

- Case (ii) : One of the two vertices is not already in G_k .

Euler's Formula - Proof by induction

- Case (ii) : One of the two vertices is not already in G_k .
- Assume a_{k+1} is in G_k and b_{k+1} is not.

Euler's Formula - Proof by induction

- Case (ii) : One of the two vertices is not already in G_k .
- Assume a_{k+1} is in G_k and b_{k+1} is not.
- Adding this new edge does not produce new regions – b_{k+1} must be in a region that has a_{k+1} on its boundary.

Euler's Formula - Proof by induction

- Case (ii) : One of the two vertices is not already in G_k .
- Assume a_{k+1} is in G_k and b_{k+1} is not.
- Adding this new edge does not produce new regions – b_{k+1} must be in a region that has a_{k+1} on its boundary.
- $r_{k+1} = r_k$, $e_{k+1} = e_k + 1$ and $v_{k+1} = v_k + 1$ - formula still holds

Euler's Formula - Proof by induction

- Case (ii) : One of the two vertices is not already in G_k .
- Assume a_{k+1} is in G_k and b_{k+1} is not.
- Adding this new edge does not produce new regions – b_{k+1} must be in a region that has a_{k+1} on its boundary.
- $r_{k+1} = r_k$, $e_{k+1} = e_k + 1$ and $v_{k+1} = v_k + 1$ - formula still holds
- Completes the induction argument!

Euler's Formula - Very Important Corollaries

Corollary

If G is a connected planar simple graph with e edges and v vertices, where $v \geq 3$, then $e \leq 3v - 6$.

Corollary

If G is a connected planar simple graph then G has a vertex of degree not exceeding five.

Corollary

If a connected planar simple graph has e edges and v vertices with $v \geq 3$ and no circuits of length 3 then $e \leq 2v - 4$.

Euler's Formula - Very Important Corollaries

- Second corollary is easy to prove. It is used to prove Corollary 1. (Try!)

Euler's Formula - Very Important Corollaries

- Second corollary is easy to prove. It is used to prove Corollary 1. (Try!)
- First corollary shows that K_5 is nonplanar - It has five vertices and ten edges. $e \leq 3v - 6$ is not satisfied!

Euler's Formula - Very Important Corollaries

- Second corollary is easy to prove. It is used to prove Corollary 1. (Try!)
- First corollary shows that K_5 is nonplanar - It has five vertices and ten edges. $e \leq 3v - 6$ is not satisfied!
- $K_{3,3}$ satisfies Corollary 1 but it is not a sufficient condition – it violates Corollary 3. $e = 9, 2v - 4 = 8$

Graph coloring

Idea of Graph coloring

- Coloring of maps - regions with common border different colors.

Idea of Graph coloring

- Coloring of maps - regions with common border different colors.
- One way - a new color every time! -

Idea of Graph coloring

- Coloring of maps - regions with common border different colors.
- One way - a new color every time! - Very inefficient! We would prefer smaller number of colors!

Idea of Graph coloring

- Coloring of maps - regions with common border different colors.
- One way - a new color every time! - Very inefficient! We would prefer smaller number of colors!
- Consider the problem of determining the least number of colors that can be used to color a map so that adjacent regions never have the same color.

Idea of Graph coloring

- Coloring of maps - regions with common border different colors.
- One way - a new color every time! - **Very inefficient! We would prefer smaller number of colors!**
- Consider the problem of determining the **least number of colors** that can be used to color a map so that adjacent regions never have the same color.
- A **coloring** of a simple graph is the assignment of a color to each vertex of the graph so that no two adjacent vertices are assigned the same color.

Idea of Graph coloring

- Coloring of maps - regions with common border different colors.
- One way - a new color every time! - **Very inefficient! We would prefer smaller number of colors!**
- Consider the problem of determining the **least number of colors** that can be used to color a map so that adjacent regions never have the same color.
- A **coloring** of a simple graph is the assignment of a color to each vertex of the graph so that no two adjacent vertices are assigned the same color.
- The **chromatic number** $\chi(G)$ of a graph is the least number of colors needed for a coloring of this graph.

Four Color Theorem

- Asking for chromatic number of planar graph - same as asking for the minimum number of colors required to color a planar map so that no two adjacent regions are assigned the same color.

Four Color Theorem

- Asking for chromatic number of planar graph - same as asking for the minimum number of colors required to color a planar map so that no two adjacent regions are assigned the same color.
- An old question - more than 100 years old!

Theorem (Four Color Theorem)

The chromatic number of a planar graph is no greater than four.

Four Color Theorem

- Asking for chromatic number of planar graph - same as asking for the minimum number of colors required to color a planar map so that no two adjacent regions are assigned the same color.
- An old question - more than 100 years old!

Theorem (Four Color Theorem)

The chromatic number of a planar graph is no greater than four.

- Posed as a conjecture in 1850s.

Four Color Theorem

- Asking for chromatic number of planar graph - same as asking for the minimum number of colors required to color a planar map so that no two adjacent regions are assigned the same color.
- An old question - more than 100 years old!

Theorem (Four Color Theorem)

The chromatic number of a planar graph is no greater than four.

- Posed as a conjecture in 1850s.
- Many incorrect proofs were published, many with hard to find errors that they were thought to be true!

Four Color Theorem

- Asking for chromatic number of planar graph - same as asking for the minimum number of colors required to color a planar map so that no two adjacent regions are assigned the same color.
- An old question - more than 100 years old!

Theorem (Four Color Theorem)

The chromatic number of a planar graph is no greater than four.

- Posed as a conjecture in 1850s.
- Many incorrect proofs were published, many with hard to find errors that they were thought to be true!
- Finally proved in 1976 - Appel and Haken.

Four Color Theorem

Four Color Theorem

- The proof uses case-by-case analysis by computer.

Four Color Theorem

- The proof uses case-by-case analysis by computer.
- First assume the theorem is false, then show that there can be ≈ 2000 types of possible counterexamples.
- Then they show none of these types exist.

Four Color Theorem

- The proof uses case-by-case analysis by computer.
- First assume the theorem is false, then show that there can be ≈ 2000 types of possible counterexamples.
- Then they show none of these types exist.
- Simpler proofs have come but everything relies on computer.

Four Color Theorem

- The proof uses case-by-case analysis by computer.
- First assume the theorem is false, then show that there can be ≈ 2000 types of possible counterexamples.
- Then they show none of these types exist.
- Simpler proofs have come but everything relies on computer.
- Doesn't work for nonplanar graphs – they can have large chromatic number.

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.
- Examples –

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.
- Examples –
- Chromatic color of K_n –

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.
- Examples –
- Chromatic color of K_n –
 - It can be colored by n colors.

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.
- Examples –
- Chromatic color of K_n –
 - It can be colored by n colors.
 - Can there be fewer colors? -

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.
- Examples –
- Chromatic color of K_n –
 - It can be colored by n colors.
 - Can there be fewer colors? - No! since every vertex is connected to every other vertex and therefore needs a different color.

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.
- Examples –
- Chromatic color of K_n –
 - It can be colored by n colors.
 - Can there be fewer colors? - No! since every vertex is connected to every other vertex and therefore needs a different color.
 - $\chi(K_n) = n$.

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.
- Examples –
- Chromatic color of K_n –
 - It can be colored by n colors.
 - Can there be fewer colors? - No! since every vertex is connected to every other vertex and therefore needs a different color.
 - $\chi(K_n) = n$.
 - Note K_n is not planar for $n \geq 5$ so four color theorem holds!

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.
- Examples –
- Chromatic color of K_n –
 - It can be colored by n colors.
 - Can there be fewer colors? - No! since every vertex is connected to every other vertex and therefore needs a different color.
 - $\chi(K_n) = n$.
 - Note K_n is not planar for $n \geq 5$ so four color theorem holds!
- What is the chromatic number of the complete bipartite graph $K_{m,n}$?

Coloring of graphs

- To show chromatic number is k – first color with k colors and then show it cannot be colored with fewer than k colors.
- Examples –
- Chromatic color of K_n –
 - It can be colored by n colors.
 - Can there be fewer colors? - No! since every vertex is connected to every other vertex and therefore needs a different color.
 - $\chi(K_n) = n$.
 - Note K_n is not planar for $n \geq 5$ so four color theorem holds!
- What is the chromatic number of the complete bipartite graph $K_{m,n}$?
- $\chi(K_{m,n}) = 2$ since its bipartite.

Chromatic Numbers

- Chromatic number of cycle C_n , $n \geq 3$
 - Consider C_n when n is even.

Chromatic Numbers

- Chromatic number of cycle C_n , $n \geq 3$
 - Consider C_n when n is even.
 - Pick a vertex, color it say with red.

Chromatic Numbers

- Chromatic number of cycle C_n , $n \geq 3$
 - Consider C_n when n is even.
 - Pick a vertex, color it say with red.
 - Proceed along in a clockwise direction with a planar representation of the graph – coloring the second vertex blue, the third vertex red and so on.

Chromatic Numbers

- Chromatic number of cycle C_n , $n \geq 3$
 - Consider C_n when n is even.
 - Pick a vertex, color it say with red.
 - Proceed along in a clockwise direction with a planar representation of the graph – coloring the second vertex blue, the third vertex red and so on.
 - The first and $n - 1$ vertices are both colored red and therefore n th vertex can be colored blue.

Chromatic Numbers

- Chromatic number of cycle C_n , $n \geq 3$
 - Consider C_n when n is even.
 - Pick a vertex, color it say with red.
 - Proceed along in a clockwise direction with a planar representation of the graph – coloring the second vertex blue, the third vertex red and so on.
 - The first and $n - 1$ vertices are both colored red and therefore n th vertex can be colored blue.
 - What if n is odd?

Chromatic Numbers

- Chromatic number of cycle C_n , $n \geq 3$
 - Consider C_n when n is even.
 - Pick a vertex, color it say with red.
 - Proceed along in a clockwise direction with a planar representation of the graph – coloring the second vertex blue, the third vertex red and so on.
 - The first and $n - 1$ vertices are both colored red and therefore n th vertex can be colored blue.
 - What if n is odd? In that case, first and $n - 1$ st vertex are of different colors - red and blue and therefore n th vertex has to be a third color.

Chromatic Numbers

- Chromatic number of cycle C_n , $n \geq 3$
 - Consider C_n when n is even.
 - Pick a vertex, color it say with red.
 - Proceed along in a clockwise direction with a planar representation of the graph – coloring the second vertex blue, the third vertex red and so on.
 - The first and $n - 1$ vertices are both colored red and therefore n th vertex can be colored blue.
 - What if n is odd? In that case, first and $n - 1$ st vertex are of different colors - red and blue and therefore n th vertex has to be a third color.
 - $\chi(C_n) = 2$ if n is an even positive integer, $\chi(C_n) = 3$, if n is odd, $n \geq 3$.

Five Color Theorem – Heawood's Theorem

Theorem

A planar graph G can be assigned a proper vertex k -coloring, $k \geq 5$.

- Induction on the number of vertices.

Five Color Theorem – Heawood's Theorem

Theorem

A planar graph G can be assigned a proper vertex k -coloring, $k \geq 5$.

- Induction on the number of vertices. $P(n) : G_n$ can be assigned a proper k -coloring.

Five Color Theorem – Heawood's Theorem

Theorem

A planar graph G can be assigned a proper vertex k -coloring, $k \geq 5$.

- Induction on the number of vertices. $P(n) : G_n$ can be assigned a proper k -coloring.
- Basis : $P(r)$ is trivially true, $1 \leq r \leq 5$.

Five Color Theorem – Heawood's Theorem

Theorem

A planar graph G can be assigned a proper vertex k -coloring, $k \geq 5$.

- Induction on the number of vertices. $P(n) : G_n$ can be assigned a proper k -coloring.
- Basis : $P(r)$ is trivially true, $1 \leq r \leq 5$.
- Hypothesis: Assume that $P(r)$, $r \geq 5$ is true.

Five Color Theorem – Heawood's Theorem

Theorem

A planar graph G can be assigned a proper vertex k -coloring, $k \geq 5$.

- Induction on the number of vertices. $P(n) : G_n$ can be assigned a proper k -coloring.
- Basis : $P(r)$ is trivially true, $1 \leq r \leq 5$.
- Hypothesis: Assume that $P(r)$, $r \geq 5$ is true.
- T.S.T. G_{r+1} can be assigned a k -vertex coloring if G_r can be assigned a k -vertex coloring.

Five Color Theorem – Heawood's Theorem

Theorem

A planar graph G can be assigned a proper vertex k -coloring, $k \geq 5$.

- Induction on the number of vertices. $P(n) : G_n$ can be assigned a proper k -coloring.
- Basis : $P(r)$ is trivially true, $1 \leq r \leq 5$.
- Hypothesis: Assume that $P(r)$, $r \geq 5$ is true.
- T.S.T. G_{r+1} can be assigned a k -vertex coloring if G_r can be assigned a k -vertex coloring.
- Lemma : Let G be a simple connected planar graph, then the minimum degree of a graph is less than 5.

Proof by Induction

- Let that vertex be v .

Proof by Induction

- Let that vertex be v .
- Remove v and incident edges to get a graph G' – it is five-colorable.

Proof by Induction

- Let that vertex be v .
- Remove v and incident edges to get a graph G' – it is five-colorable.
- Case(i) : Suppose the 5 colors of G' are not connected to v - then just give the missing color to v and G_{r+1} is five-colorable.

Proof by Induction

- Let that vertex be v .
- Remove v and incident edges to get a graph G' – it is five-colorable.
- Case(i) : Suppose the 5 colors of G' are not connected to v - then just give the missing color to v and G_{r+1} is five-colorable.
- Case(ii): Suppose all 5 colors are connected to v :

Proof by Induction

- Let that vertex be v .
- Remove v and incident edges to get a graph G' – it is five-colorable.
- Case(i) : Suppose the 5 colors of G' are not connected to v - then just give the missing color to v and G_{r+1} is five-colorable.
- Case(ii): Suppose all 5 colors are connected to v :
- Let the five vertices adjacent to v be called u_1, u_2, u_3, u_4, u_5 and colored c_1, c_2, c_3, c_4, c_5 .

Proof by Induction

- $H_{i,j}$ a subgraph of G' induced by vertices that have the colors c_i and c_j .

Proof by Induction

- $H_{i,j}$ a subgraph of G' induced by vertices that have the colors c_i and c_j .
- In $H_{1,3}$ consider two cases:

Proof by Induction

- $H_{i,j}$ a subgraph of G' induced by vertices that have the colors c_i and c_j .
- In $H_{1,3}$ consider two cases:
- Case (a): There is no path between u_1 and u_3 , i.e. two disconnected components in $H_{1,3}$.

Proof by Induction

- $H_{i,j}$ a subgraph of G' induced by vertices that have the colors c_i and c_j .
- In $H_{1,3}$ consider two cases:
- Case (a): There is no path between u_1 and u_3 , i.e. two disconnected components in $H_{1,3}$.
- Interchange c_1 and c_3 in the component that is connected to u_1 .

Proof by Induction

- $H_{i,j}$ a subgraph of G' induced by vertices that have the colors c_i and c_j .
- In $H_{1,3}$ consider two cases:
- Case (a): There is no path between u_1 and u_3 , i.e. two disconnected components in $H_{1,3}$.
- Interchange c_1 and c_3 in the component that is connected to u_1 .
- Then v is no longer adjacent to a vertex of color c_1 and v can be given c_1 .

Proof by Induction

- Case(b): There is a path between u_1 and u_3 in $H_{1,3}$.

Proof by Induction

- Case(b): There is a path between u_1 and u_3 in $H_{1,3}$.
- Include v in this path, we get a circuit C .

Proof by Induction

- Case(b): There is a path between u_1 and u_3 in $H_{1,3}$.
- Include v in this path, we get a circuit C .
- Since u_i s are numbered clockwise this implies exactly u_2 or u_4 is inside C .

Proof by Induction

- Case(b): There is a path between u_1 and u_3 in $H_{1,3}$.
- Include v in this path, we get a circuit C .
- Since u_i s are numbered clockwise this implies exactly u_2 or u_4 is inside C .
- This implies u_2 and u_4 are in different connected components of $H_{2,4}$.

Proof by Induction

- Case(b): There is a path between u_1 and u_3 in $H_{1,3}$.
- Include v in this path, we get a circuit C .
- Since u_i s are numbered clockwise this implies exactly u_2 or u_4 is inside C .
- This implies u_2 and u_4 are in different connected components of $H_{2,4}$.
- Then do the same as Case (a) - switch colors c_2 and c_4 in $H_{2,4}$ that is connected to u_2 .

Proof by Induction

- Case(b): There is a path between u_1 and u_3 in $H_{1,3}$.
- Include v in this path, we get a circuit C .
- Since u_i s are numbered clockwise this implies exactly u_2 or u_4 is inside C .
- This implies u_2 and u_4 are in different connected components of $H_{2,4}$.
- Then do the same as Case (a) - switch colors c_2 and c_4 in $H_{2,4}$ that is connected to u_2 .
- v is not adjacent to vertex of color c_2 and can be given c_2 . -

Proof by Induction

- Case(b): There is a path between u_1 and u_3 in $H_{1,3}$.
- Include v in this path, we get a circuit C .
- Since u_i s are numbered clockwise this implies exactly u_2 or u_4 is inside C .
- This implies u_2 and u_4 are in different connected components of $H_{2,4}$.
- Then do the same as Case (a) - switch colors c_2 and c_4 in $H_{2,4}$ that is connected to u_2 .
- v is not adjacent to vertex of color c_2 and can be given c_2 . - Done!

Algorithms for finding Chromatic Numbers

- The best algorithms have exponential worst-case time complexity (in the number of vertices of the graph).
- Finding an approximation to the chromatic number of a graph is difficult.

Algorithms for finding Chromatic Numbers

- The best algorithms have exponential worst-case time complexity (in the number of vertices of the graph).
- Finding an approximation to the chromatic number of a graph is difficult.
- Applications - scheduler of exams, no students have two exams at the same time.

Algorithms for finding Chromatic Numbers

- The best algorithms have exponential worst-case time complexity (in the number of vertices of the graph).
- Finding an approximation to the chromatic number of a graph is difficult.
- Applications - scheduler of exams, no students have two exams at the same time.
- Frequency assignments for TV stations so that within a certain distance two stations won't have the same channel.

Algorithms for finding Chromatic Numbers

- The best algorithms have exponential worst-case time complexity (in the number of vertices of the graph).
- Finding an approximation to the chromatic number of a graph is difficult.
- Applications - scheduler of exams, no students have two exams at the same time.
- Frequency assignments for TV stations so that within a certain distance two stations won't have the same channel.
- Compilers - assigning index registers for a loop.