

# EP 1027: Maxwell's Equations and Electromagnetic Waves

Instructor: Shubho Roy<sup>1</sup>  
(Dept. of Physics)

Lecture 12

April 20, 2019

---

<sup>1</sup>Email: [sroy@iith.ac.in](mailto:sroy@iith.ac.in), Office: C 313 D, Office hrs: Walk in or by Email Appointment

# Agenda

# Agenda

- ▶ Resonant Cavity and Guided Waves

# References/Readings

# References/Readings

- ▶ Griffiths, D.J., **Introduction to Electrodynamics, Ch. 9**

# “Resonant” EM Waves

# “Resonant” EM Waves

- ▶ Analogous to organ pipes (sonic/acoustic resonant cavities)

# “Resonant” EM Waves

- ▶ Analogous to organ pipes (sonic/acoustic resonant cavities)
- ▶ Longitudinal pressure/density oscillations of the air column (stimulated by a tuning fork): standing waves; resonant frequencies w/ large amplitudes



# “Resonant” EM Waves

- ▶ Analogous to organ pipes (sonic/acoustic resonant cavities)
- ▶ Longitudinal pressure/density oscillations of the air column (stimulated by a tuning fork): standing waves; resonant frequencies w/ large amplitudes
- ▶ “Resonant cavity” for EM waves : closed container made up of conducting walls

# “Resonant” EM Waves

- ▶ Analogous to organ pipes (sonic/acoustic resonant cavities)
- ▶ Longitudinal pressure/density oscillations of the air column (stimulated by a tuning fork): standing waves; resonant frequencies w/ large amplitudes
- ▶ “Resonant cavity” for EM waves : closed container made up of conducting walls
- ▶ EM Excitations can be stimulated by an antenna (emitter) inside :  
Can achieve resonant (large amplitude) standing EM waves

# “Resonant” EM Waves

- ▶ Analogous to organ pipes (sonic/acoustic resonant cavities)
- ▶ Longitudinal pressure/density oscillations of the air column (stimulated by a tuning fork): standing waves; resonant frequencies w/ large amplitudes
- ▶ “Resonant cavity” for EM waves : closed container made up of conducting walls
- ▶ EM Excitations can be stimulated by an antenna (emitter) inside : Can achieve resonant (large amplitude) standing EM waves
- ▶ Perfect conducting walls ( $\sigma \rightarrow \infty$ ), waves last forever; realistic (Ohmic) conductors, Ohmic heat loss causes waves to dissipate and disappear after a while

# EM waves in Resonant Cavity

# EM waves in Resonant Cavity

- ▶ Must solve for **E** and **B** fields oscillating in the cavity

# EM waves in Resonant Cavity

- ▶ Must solve for **E** and **B** fields oscillating in the cavity
- ▶ More complicated than sound waves;
  - a) These are *vector* fields,
  - b) **E** and **B** field equations are coupled thru Maxwell's eqn.s,
  - c) Both must satisfy their own boundary conditions

# EM waves in Resonant Cavity

- ▶ Must solve for **E** and **B** fields oscillating in the cavity
- ▶ More complicated than sound waves;
  - a) These are *vector* fields,
  - b) **E** and **B** field equations are coupled thru Maxwell's eqn.s,
  - c) Both must satisfy their own boundary conditions
- ▶ Assume: Empty cavity (easy to amend this to cavity filled with dielectric)

# EM waves in Resonant Cavity

- ▶ Must solve for **E** and **B** fields oscillating in the cavity
- ▶ More complicated than sound waves;
  - a) These are *vector* fields,
  - b) **E** and **B** field equations are coupled thru Maxwell's eqn.s,
  - c) Both must satisfy their own boundary conditions
- ▶ Assume: Empty cavity (easy to amend this to cavity filled with dielectric)
- ▶ Maxwell equations

$$\nabla \cdot \mathbf{E} = 0,$$

$$\nabla \cdot \mathbf{B} = 0,$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t},$$

$$\nabla \times \mathbf{B} = \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t}$$



# EM waves in Resonant Cavity

- ▶ Must solve for **E** and **B** fields oscillating in the cavity
- ▶ More complicated than sound waves;
  - a) These are *vector* fields,
  - b) **E** and **B** field equations are coupled thru Maxwell's eqn.s,
  - c) Both must satisfy their own boundary conditions
- ▶ Assume: Empty cavity (easy to amend this to cavity filled with dielectric)
- ▶ Maxwell equations

$$\nabla \cdot \mathbf{E} = 0,$$

$$\nabla \cdot \mathbf{B} = 0,$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t},$$

$$\nabla \times \mathbf{B} = \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t}$$

- ▶ Wave equations!

$$\nabla^2 \mathbf{E} - \frac{1}{c^2} \frac{\partial^2 \mathbf{E}}{\partial t^2} = 0 \quad \nabla^2 \mathbf{B} - \frac{1}{c^2} \frac{\partial^2 \mathbf{B}}{\partial t^2} = 0$$

# EM waves in Resonant Cavity: Eigenvalue problem

# EM waves in Resonant Cavity: Eigenvalue problem

- ▶ Oscillating ansatz:

$$\mathbf{E}(t, \mathbf{x}) = \mathbf{E}(\mathbf{x})e^{-i\omega t}, \quad \mathbf{B}(t, \mathbf{x}) = \mathbf{B}(\mathbf{x})e^{-i\omega t}$$

# EM waves in Resonant Cavity: Eigenvalue problem

- ▶ Oscillating ansatz:

$$\mathbf{E}(t, \mathbf{x}) = \mathbf{E}(\mathbf{x})e^{-i\omega t}, \quad \mathbf{B}(t, \mathbf{x}) = \mathbf{B}(\mathbf{x})e^{-i\omega t}$$

- ▶ Wave equation then becomes an eigenvalue problem:

$$\nabla^2 \mathbf{E}(\mathbf{x}) = -\frac{\omega^2}{c^2} \mathbf{E}(\mathbf{x}), \quad \nabla^2 \mathbf{B}(\mathbf{x}) = -\frac{\omega^2}{c^2} \mathbf{B}(\mathbf{x})$$

# EM waves in Resonant Cavity: Eigenvalue problem

- ▶ Oscillating ansatz:

$$\mathbf{E}(t, \mathbf{x}) = \mathbf{E}(\mathbf{x})e^{-i\omega t}, \quad \mathbf{B}(t, \mathbf{x}) = \mathbf{B}(\mathbf{x})e^{-i\omega t}$$

- ▶ Wave equation then becomes an eigenvalue problem:

$$\nabla^2 \mathbf{E}(\mathbf{x}) = -\frac{\omega^2}{c^2} \mathbf{E}(\mathbf{x}), \quad \nabla^2 \mathbf{B}(\mathbf{x}) = -\frac{\omega^2}{c^2} \mathbf{B}(\mathbf{x})$$

- ▶ Our job: Solve the eigenvalue problem subject to some boundary condition

# EM waves in Resonant Cavity: Eigenvalue problem

- ▶ Oscillating ansatz:

$$\mathbf{E}(t, \mathbf{x}) = \mathbf{E}(\mathbf{x})e^{-i\omega t}, \quad \mathbf{B}(t, \mathbf{x}) = \mathbf{B}(\mathbf{x})e^{-i\omega t}$$

- ▶ Wave equation then becomes an eigenvalue problem:

$$\nabla^2 \mathbf{E}(\mathbf{x}) = -\frac{\omega^2}{c^2} \mathbf{E}(\mathbf{x}), \quad \nabla^2 \mathbf{B}(\mathbf{x}) = -\frac{\omega^2}{c^2} \mathbf{B}(\mathbf{x})$$

- ▶ Our job: Solve the eigenvalue problem subject to some boundary condition
- ▶ However: Wave equation is not everything, not all 6 components of  $\mathbf{E}$  and  $\mathbf{B}$  are independent

# EM waves in Resonant Cavity: Eigenvalue problem

- ▶ Oscillating ansatz:

$$\mathbf{E}(t, \mathbf{x}) = \mathbf{E}(\mathbf{x})e^{-i\omega t}, \quad \mathbf{B}(t, \mathbf{x}) = \mathbf{B}(\mathbf{x})e^{-i\omega t}$$

- ▶ Wave equation then becomes an eigenvalue problem:

$$\nabla^2 \mathbf{E}(\mathbf{x}) = -\frac{\omega^2}{c^2} \mathbf{E}(\mathbf{x}), \quad \nabla^2 \mathbf{B}(\mathbf{x}) = -\frac{\omega^2}{c^2} \mathbf{B}(\mathbf{x})$$

- ▶ Our job: Solve the eigenvalue problem subject to some boundary condition
- ▶ However: Wave equation is not everything, not all 6 components of  $\mathbf{E}$  and  $\mathbf{B}$  are independent
- ▶ To simplify life: assume cylindrical cavities and wave guides from now on

# “Resonant” EM Waves: Boundary conditions



# “Resonant” EM Waves: Boundary conditions

- ▶ Waves confined in some finite cavity with *perfectly conducting walls*

$$\mathbf{E}_{\parallel} = 0,$$

$$B_{\perp} = 0.$$

# “Resonant” EM Waves: Boundary conditions

- ▶ Waves confined in some finite cavity with *perfectly conducting walls*

$$\mathbf{E}_{\parallel} = 0,$$

$$B_{\perp} = 0.$$

- ▶ What about other two b.c.'s ?

$$E_{\perp} = \frac{\sigma}{\varepsilon_0},$$

$$\mathbf{B}_{\parallel} = \mu_0 \mathbf{K} \times \hat{\mathbf{n}}$$

# “Resonant” EM Waves: Boundary conditions

- ▶ Waves confined in some finite cavity with *perfectly conducting walls*

$$\mathbf{E}_{\parallel} = 0,$$

$$B_{\perp} = 0.$$

- ▶ What about other two b.c.'s ?

$$E_{\perp} = \frac{\sigma}{\varepsilon_0},$$

$$\mathbf{B}_{\parallel} = \mu_0 \mathbf{K} \times \hat{\mathbf{n}}$$

- ▶  $\sigma, \mathbf{k}$  in conducting walls will adjust themselves to satisfy above, no new b.c.!

# Cylindrical Cavities

# Cylindrical Cavities

- ▶ Infinitely long axis, but arbitrary cross-section shape

# Cylindrical Cavities

- ▶ Infinitely long axis, but arbitrary cross-section shape
- ▶ Axial/Cylindrical Ansatz:

$$\mathbf{E}(t, \mathbf{x}) = \mathbf{E}(x, y) e^{-i\omega t \pm ikz}, \quad \mathbf{B}(t, \mathbf{x}) = \mathbf{B}(x, y) e^{-i\omega t \pm ikz}$$

# Cylindrical Cavities

- ▶ Infinitely long axis, but arbitrary cross-section shape
- ▶ Axial/Cylindrical Ansatz:

$$\mathbf{E}(t, \mathbf{x}) = \mathbf{E}(x, y) e^{-i\omega t \pm ikz}, \quad \mathbf{B}(t, \mathbf{x}) = \mathbf{B}(x, y) e^{-i\omega t \pm ikz}$$

- ▶ Maxwell equations (coupled linear diff. eqn.)

$$\begin{aligned} \frac{\partial B_z}{\partial y} \mp ikB_y &= -\frac{i\omega}{c^2} E_x \\ \pm ikB_x - \frac{\partial B_z}{\partial x} &= -i\frac{\omega}{c^2} E_y \\ \frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} &= -i\frac{\omega}{c^2} E_z \\ &\vdots \end{aligned}$$

# Cylindrical Cavities

- ▶ Infinitely long axis, but arbitrary cross-section shape
- ▶ Axial/Cylindrical Ansatz:

$$\mathbf{E}(t, \mathbf{x}) = \mathbf{E}(x, y) e^{-i\omega t \pm ikz}, \quad \mathbf{B}(t, \mathbf{x}) = \mathbf{B}(x, y) e^{-i\omega t \pm ikz}$$

- ▶ Maxwell equations (coupled linear diff. eqn.)

$$\begin{aligned} \frac{\partial B_z}{\partial y} \mp ikB_y &= -\frac{i\omega}{c^2} E_x \\ \pm ikB_x - \frac{\partial B_z}{\partial x} &= -i\frac{\omega}{c^2} E_y \\ \frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} &= -i\frac{\omega}{c^2} E_z \\ &\vdots \end{aligned}$$

- ▶ Solve  $E_x, E_y, B_x, B_y$  in terms of  $E_z, B_z$ .



# Cylindrical Cavities

# Cylindrical Cavities

► For instance,

$$\begin{aligned} E_x &= \frac{1}{\omega^2/c^2 - k^2} \left( \pm ik \frac{\partial E_z}{\partial x} + i\omega \frac{\partial B_z}{\partial y} \right) \\ &= \frac{1}{\omega^2/c^2 - k^2} \left( \frac{\partial^2 E_z}{\partial x \partial z} + i\omega \frac{\partial B_z}{\partial y} \right) \end{aligned}$$

# Cylindrical Cavities

- For instance,

$$\begin{aligned} E_x &= \frac{1}{\omega^2/c^2 - k^2} \left( \pm ik \frac{\partial E_z}{\partial x} + i\omega \frac{\partial B_z}{\partial y} \right) \\ &= \frac{1}{\omega^2/c^2 - k^2} \left( \frac{\partial^2 E_z}{\partial x \partial z} + i\omega \frac{\partial B_z}{\partial y} \right) \end{aligned}$$

- Similar for  $E_y, B_x, B_y$

$$\begin{aligned} E_y &= \frac{1}{\omega^2/c^2 - k^2} \left( \frac{\partial^2 E_z}{\partial y \partial z} - i\omega \frac{\partial B_z}{\partial x} \right) \\ B_x &= \frac{1}{\omega^2/c^2 - k^2} \left( \frac{\partial^2 B_z}{\partial x \partial z} - i \frac{\omega}{c^2} \frac{\partial E_z}{\partial y} \right) \\ B_y &= \frac{1}{\omega^2/c^2 - k^2} \left( \frac{\partial^2 B_z}{\partial y \partial z} + i \frac{\omega}{c^2} \frac{\partial E_z}{\partial x} \right) \end{aligned}$$

# Cylindrical Cavities

- ▶ For instance,

$$\begin{aligned} E_x &= \frac{1}{\omega^2/c^2 - k^2} \left( \pm ik \frac{\partial E_z}{\partial x} + i\omega \frac{\partial B_z}{\partial y} \right) \\ &= \frac{1}{\omega^2/c^2 - k^2} \left( \frac{\partial^2 E_z}{\partial x \partial z} + i\omega \frac{\partial B_z}{\partial y} \right) \end{aligned}$$

- ▶ Similar for  $E_y, B_x, B_y$

$$\begin{aligned} E_y &= \frac{1}{\omega^2/c^2 - k^2} \left( \frac{\partial^2 E_z}{\partial y \partial z} - i\omega \frac{\partial B_z}{\partial x} \right) \\ B_x &= \frac{1}{\omega^2/c^2 - k^2} \left( \frac{\partial^2 B_z}{\partial x \partial z} - i \frac{\omega}{c^2} \frac{\partial E_z}{\partial y} \right) \\ B_y &= \frac{1}{\omega^2/c^2 - k^2} \left( \frac{\partial^2 B_z}{\partial y \partial z} + i \frac{\omega}{c^2} \frac{\partial E_z}{\partial x} \right) \end{aligned}$$

- ▶  $E_z, B_z$  determine everything! Themselves determined by EV equations

$$\nabla^2 E_z = -\frac{\omega^2}{c^2} E_z; \quad \nabla^2 B_z = -\frac{\omega^2}{c^2} B_z$$

# Cylindrical Cavities: TE and TM modes

# Cylindrical Cavities: TE and TM modes

- ▶ Cavity oscillations are not transverse!  $E_z$  and  $B_z$  can't simultaneously vanish for then no waves at all

# Cylindrical Cavities: TE and TM modes

- ▶ Cavity oscillations are not transverse!  $E_z$  and  $B_z$  can't simultaneously vanish for then no waves at all
- ▶ However can have  $E_z = 0$  ( $B_z \neq 0$ ) : **TE** waves or can have  $B_z = 0$  ( $E_z \neq 0$ ) : **TM** waves

# Cylindrical Cavities: TE and TM modes

- ▶ Cavity oscillations are not transverse!  $E_z$  and  $B_z$  can't simultaneously vanish for then no waves at all
- ▶ However can have  $E_z = 0$  ( $B_z \neq 0$ ) : **TE** waves or can have  $B_z = 0$  ( $E_z \neq 0$ ) : **TM** waves
- ▶ Rectangular cavities: Cross-section rectangular ( $a, b$ )

$$E_z(x, y, z) = E_0 \sin\left(\frac{m\pi x}{a}\right) \sin\left(\frac{n\pi y}{b}\right) \sin kz$$

- ▶ w/ dispersion

$$\frac{\omega_{kmn}^2}{c^2} = \left(\frac{m\pi}{a}\right)^2 + \left(\frac{n\pi}{b}\right)^2 + k^2$$

- ▶ For cavity of depth  $d$ :

$$k = \frac{l\pi}{d}, \omega_{lmn} = c \sqrt{\left(\frac{m\pi}{a}\right)^2 + \left(\frac{n\pi}{b}\right)^2 + \left(\frac{l\pi}{d}\right)^2}$$



# Cylindrical Waveguides

# Cylindrical Waveguides

- ▶ waveguides: infinite depth,  $d \rightarrow \infty$

# Cylindrical Waveguides

- ▶ waveguides: infinite depth,  $d \rightarrow \infty$
- ▶ Wave velocity,

$$v = \frac{\omega}{k} = c \sqrt{\left(\frac{m\pi}{ak}\right)^2 + \left(\frac{n\pi}{bk}\right)^2 + 1} > c!$$

# Cylindrical Waveguides

- ▶ waveguides: infinite depth,  $d \rightarrow \infty$
- ▶ Wave velocity,

$$v = \frac{\omega}{k} = c \sqrt{\left(\frac{m\pi}{ak}\right)^2 + \left(\frac{n\pi}{bk}\right)^2 + 1} > c!$$

- ▶ Group velocity

$$v_g = \frac{d\omega}{dk} = \frac{c}{\sqrt{\left(\frac{m\pi}{ak}\right)^2 + \left(\frac{n\pi}{bk}\right)^2 + 1}} < c!$$

- ▶ Coaxial Cables: **TEM** modes exist! Cylindrical symmetry reduces the problem to electrostatics and magnetostatics in **2** space dimensions instead of **3**. The solution:

$$\mathbf{E} = \frac{A \cos(kz - \omega t)}{\rho} \hat{\rho}$$
$$\mathbf{B} = \frac{A \cos(kz - \omega t)}{c\rho} \hat{\phi}$$

Full analysis in Sec. 9.5.2 of Griffiths' text