

EP 1027: Maxwell's Equations and Electromagnetic Waves

Instructor: Shubho Roy¹
(Dept. of Physics)

Lecture 13

April 23, 2019

¹Email: sroy@iith.ac.in, Office: C 313 D, Office hrs: Walk in or by email appointment

Agenda

Agenda

- ▶ Radiation emitted from point charges: Retarded potential, Lienard-Wiechert Potential, Radiation fields and pattern, Dielectrics (Mach-Cerenkov Effect)

References/Readings

References/Readings

- ▶ Griffiths, D.J., **Introduction to Electrodynamics, Ch. 10, 11**

Radiation from a point charge: Retarded potential

Radiation from a point charge: Retarded potential

- ▶ When point charge is at rest or uniform motion, **E** and **B** fields are dragged along as if rigidly attached to the charge

Radiation from a point charge: Retarded potential

- ▶ When point charge is at rest or uniform motion, \mathbf{E} and \mathbf{B} fields are dragged along as if rigidly attached to the charge
- ▶ When a point charge accelerates, \mathbf{E} and \mathbf{B} fields can't keep up and pieces of EM fields “break away” in the form of spherical wave-pulse (radiation)

Radiation from a point charge: Retarded potential

- ▶ When point charge is at rest or uniform motion, **E** and **B** fields are dragged along as if rigidly attached to the charge
- ▶ When a point charge accelerates, **E** and **B** fields can't keep up and pieces of EM fields “break away” in the form of spherical wave-pulse (radiation)
- ▶ The radial Coulomb piece falls off $1/R^2$; but the transverse radiation piece falls off as $1/R$ and survives to large distances

Radiation from a point charge: Retarded potential

- ▶ When point charge is at rest or uniform motion, **E** and **B** fields are dragged along as if rigidly attached to the charge
- ▶ When a point charge accelerates, **E** and **B** fields can't keep up and pieces of EM fields “break away” in the form of spherical wave-pulse (radiation)
- ▶ The radial Coulomb piece falls off $1/R^2$; but the transverse radiation piece falls off as $1/R$ and survives to large distances
- ▶ Retarded potentials: Solution to Maxwell equations for charge distribution w/ arbitrary motion

Radiation from a point charge: Retarded potential

- ▶ When point charge is at rest or uniform motion, \mathbf{E} and \mathbf{B} fields are dragged along as if rigidly attached to the charge
- ▶ When a point charge accelerates, \mathbf{E} and \mathbf{B} fields can't keep up and pieces of EM fields “break away” in the form of spherical wave-pulse (radiation)
- ▶ The radial Coulomb piece falls off $1/R^2$; but the transverse radiation piece falls off as $1/R$ and survives to large distances
- ▶ Retarded potentials: Solution to Maxwell equations for charge distribution w/ arbitrary motion
- ▶ Notation: Combine Φ, \mathbf{A} into a four-component vector $A^\mu = (A^0, A^1, A^2, A^3) = (\frac{\Phi}{c}, \mathbf{A})$ and combine ρ, \mathbf{j} into $j^\mu = (\rho c, \mathbf{j})$

Radiation from a point charge: Retarded potential

Radiation from a point charge: Retarded potential

- Maxwell equation in terms of potentials in Lorenz gauge

$$\square A^\mu(t, \mathbf{x}) = \frac{j^\mu(t, \mathbf{x})}{c^2 \epsilon_0}, \quad \square = \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2$$

Radiation from a point charge: Retarded potential

- ▶ Maxwell equation in terms of potentials in Lorenz gauge

$$\square A^\mu(t, \mathbf{x}) = \frac{j^\mu(t, \mathbf{x})}{c^2 \epsilon_0}, \quad \square = \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2$$

- ▶ Solution: *Retarded* potentials

$$A^\mu(t, \mathbf{x}) = \frac{1}{4\pi\epsilon_0} \frac{1}{c^2} \int d^3\mathbf{x}' \frac{j^\mu(t', \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}, \quad t' = t - \frac{|\mathbf{x} - \mathbf{x}'|}{c}.$$

Radiation from a point charge: Retarded potential

- ▶ Maxwell equation in terms of potentials in Lorenz gauge

$$\square A^\mu(t, \mathbf{x}) = \frac{j^\mu(t, \mathbf{x})}{c^2 \epsilon_0}, \quad \square = \frac{1}{c^2} \frac{\partial^2}{\partial t^2} - \nabla^2$$

- ▶ Solution: *Retarded* potentials

$$A^\mu(t, \mathbf{x}) = \frac{1}{4\pi\epsilon_0} \frac{1}{c^2} \int d^3\mathbf{x}' \frac{j^\mu(t', \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}, \quad t' = t - \frac{|\mathbf{x} - \mathbf{x}'|}{c}.$$

- ▶ Point charge case: $\rho(t, \mathbf{x}) = q\delta^3(\mathbf{x} - \mathbf{y}(t))$, where $\mathbf{y}(t)$ is the trajectory,

$$A^\mu(t, \mathbf{x}) = \frac{q}{4\pi\epsilon_0} \frac{v'^\mu}{c^2} \frac{1}{1 - \frac{\hat{\mathbf{n}} \cdot \mathbf{v}}{c}} \frac{1}{|\mathbf{x} - \mathbf{x}'|} \Big|_{\mathbf{x}'=\mathbf{y}(t')},$$

$$v^\mu(t') = (c, \dot{\mathbf{y}}(t')), \quad \hat{\mathbf{n}}(t) = \frac{\mathbf{x} - \mathbf{x}'}{|\mathbf{x} - \mathbf{x}'|} \Big|_{\mathbf{x}'=\mathbf{y}(t')}$$

(Liénard-Wiechert potentials, refer to supplementary material)

Radiation from a point charge: Radiation fields

Radiation from a point charge: Radiation fields

► EM fields,

$$\mathbf{E}(t, \mathbf{x}) = \underbrace{\frac{1}{4\pi\epsilon_0} q \frac{\hat{\mathbf{n}} \times [(\hat{\mathbf{n}} - \mathbf{v}'/c) \times \mathbf{a}']}{\left(1 - \frac{\hat{\mathbf{n}} \cdot \mathbf{v}}{c}\right)^3} \frac{1}{|\mathbf{x} - \mathbf{x}'(t)|}}_{\mathbf{E}_{rad}} + \text{Coulomb}$$

$$\mathbf{B}(t, \mathbf{x}) = [\hat{\mathbf{n}}] \times \mathbf{E}_{rad}(t, \mathbf{x}) + \text{Biot-Savart}$$

Radiation from a point charge: Radiation fields

- ▶ EM fields,

$$\mathbf{E}(t, \mathbf{x}) = \underbrace{\frac{1}{4\pi\epsilon_0} q \frac{\hat{\mathbf{n}} \times [(\hat{\mathbf{n}} - \mathbf{v}'/c) \times \mathbf{a}']}{(1 - \frac{\hat{\mathbf{n}} \cdot \mathbf{v}}{c})^3}}_{\mathbf{E}_{rad}} \frac{1}{|\mathbf{x} - \mathbf{x}'(t)|} + \text{Coulomb}$$

$$\mathbf{B}(t, \mathbf{x}) = [\hat{\mathbf{n}}] \times \mathbf{E}_{rad}(t, \mathbf{x}) + \text{Biot-Savart}$$

- ▶ Energy Flux: Poynting's vector (for $|\mathbf{v}| \ll c$)

$$\mathbf{S} = \frac{\mathbf{E}_{rad} \times \mathbf{B}_{rad}}{2\mu_0} \propto q^2 a'^2 \sin^2 \theta \frac{\hat{\mathbf{n}}}{|\mathbf{x} - \mathbf{x}'(t)|^2}; \theta = \theta(\hat{\mathbf{n}}, \mathbf{a})$$

Radiation from a point charge: Radiation fields

- ▶ EM fields,

$$\mathbf{E}(t, \mathbf{x}) = \underbrace{\frac{1}{4\pi\epsilon_0} q \frac{\hat{\mathbf{n}} \times [(\hat{\mathbf{n}} - \mathbf{v}'/c) \times \mathbf{a}']}{(1 - \frac{\hat{\mathbf{n}} \cdot \mathbf{v}}{c})^3}}_{\mathbf{E}_{rad}} \frac{1}{|\mathbf{x} - \mathbf{x}'(t)|} + \text{Coulomb}$$

$$\mathbf{B}(t, \mathbf{x}) = [\hat{\mathbf{n}}] \times \mathbf{E}_{rad}(t, \mathbf{x}) + \text{Biot-Savart}$$

- ▶ Energy Flux: Poynting's vector (for $|\mathbf{v}| \ll c$)

$$\mathbf{S} = \frac{\mathbf{E}_{rad} \times \mathbf{B}_{rad}}{2\mu_0} \propto q^2 a'^2 \sin^2 \theta \frac{\hat{\mathbf{n}}}{|\mathbf{x} - \mathbf{x}'(t)|^2}; \theta = \theta(\hat{\mathbf{n}}, \mathbf{a})$$

- ▶ For relativistic speeds, $|\mathbf{v}| \sim c$: Pattern depends on \mathbf{v} as well, much more complicated

Radiation from a point charge moving in media: Cerenkov Effect

Radiation from a point charge moving in media: Cerenkov Effect

- Potentials: Easy to obtain, replace $\varepsilon_0, \mu_0 \rightarrow \varepsilon, \mu$ and $c \rightarrow c/n$

$$\Phi(t, \mathbf{x}) = \frac{1}{4\pi\varepsilon} \int d^3\mathbf{x}' \frac{\rho(t - \frac{|\mathbf{x}-\mathbf{x}'|}{c/n}, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$

$$\mathbf{A}(t, \mathbf{x}) = \frac{\mu}{4\pi} \int d^3\mathbf{x}' \frac{\mathbf{j}(t - \frac{|\mathbf{x}-\mathbf{x}'|}{c/n}, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$

Here n is the refractive index of the medium

Radiation from a point charge moving in media: Cerenkov Effect

- Potentials: Easy to obtain, replace $\varepsilon_0, \mu_0 \rightarrow \varepsilon, \mu$ and $c \rightarrow c/n$

$$\Phi(t, \mathbf{x}) = \frac{1}{4\pi\varepsilon} \int d^3\mathbf{x}' \frac{\rho(t - \frac{|\mathbf{x}-\mathbf{x}'|}{c/n}, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$

$$\mathbf{A}(t, \mathbf{x}) = \frac{\mu}{4\pi} \int d^3\mathbf{x}' \frac{\mathbf{j}(t - \frac{|\mathbf{x}-\mathbf{x}'|}{c/n}, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$

Here n is the refractive index of the medium

- Radiation emitted even when there is no acceleration!. Just uniform motion with $|\mathbf{v}| > c/n$ (superluminal motion) is enough!

Radiation from a point charge moving in media: Cerenkov Effect

- Potentials: Easy to obtain, replace $\varepsilon_0, \mu_0 \rightarrow \varepsilon, \mu$ and $c \rightarrow c/n$

$$\Phi(t, \mathbf{x}) = \frac{1}{4\pi\varepsilon} \int d^3\mathbf{x}' \frac{\rho(t - \frac{|\mathbf{x}-\mathbf{x}'|}{c/n}, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$
$$\mathbf{A}(t, \mathbf{x}) = \frac{\mu}{4\pi} \int d^3\mathbf{x}' \frac{\mathbf{j}(t - \frac{|\mathbf{x}-\mathbf{x}'|}{c/n}, \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$

Here n is the refractive index of the medium

- Radiation emitted even when there is no acceleration!. Just uniform motion with $|\mathbf{v}| > c/n$ (superluminal motion) is enough!
- EM radiation fields confined inside a (Mach) cone trailing the charge. The cone angle is,

$$\sin \theta = \frac{c}{nv}$$

