

Essentials of Fourier Series and Gamma Function

Pradipto Banerjee

pradipto@iith.ac.in

IIT, Hyderabad/Bhilai

November 3, 2018

Lesson Plan

- Develop the basic theory of **Fourier** series

Lesson Plan

- Develop the basic theory of **Fourier** series
- **Bessel's** inequality and **Parseval's** identity

Lesson Plan

- Develop the basic theory of **Fourier** series
- **Bessel's** inequality and **Parseval's** identity
- Discuss (mostly without proofs) *convergence results* for Fourier series

Lesson Plan

- Develop the basic theory of **Fourier** series
- **Bessel's** inequality and **Parseval's** identity
- Discuss (mostly without proofs) *convergence results* for Fourier series
- Examples/applications of Fourier series assuming convergence

Lesson Plan

- Develop the basic theory of **Fourier** series
- **Bessel's** inequality and **Parseval's** identity
- Discuss (mostly without proofs) *convergence results* for Fourier series
- Examples/applications of Fourier series assuming convergence
- The **Gamma** and **Beta** functions with an application

Periodic functions

- A real valued function f is said to be **periodic** if there is a $L \in \mathbb{R}$ such that

$$f(x + 2L) = f(x) \quad \forall \quad x \in \mathbb{R}$$

Periodic functions

- A real valued function f is said to be **periodic** if there is a $L \in \mathbb{R}$ such that

$$f(x + 2L) = f(x) \quad \forall \quad x \in \mathbb{R}$$

- Generic examples are $\sin \omega x$ and $\cos \omega x$ having period $2\pi/\omega$

Periodic functions

- A real valued function f is said to be **periodic** if there is a $L \in \mathbb{R}$ such that

$$f(x + 2L) = f(x) \quad \forall \quad x \in \mathbb{R}$$

- Generic examples are $\sin \omega x$ and $\cos \omega x$ having period $2\pi/\omega$
- Fourier philosophy – “Every reasonably behaved periodic function can be expressed in terms of \sin and \cos ”

A useful integral identity

Lemma

Let f be a real valued $2L$ -periodic integrable function defined on \mathbb{R} . Then for any $a \in \mathbb{R}$, we have

$$\int_{-L}^L f(x) dx = \int_{-L+a}^{L+a} f(x) dx.$$

Proof of the Lemma

- We have

$$\int_{-L}^L f(x) dx = \int_{-L}^{-L+a} f(x) dx + \int_{-L+a}^L f(x) dx$$

Proof of the Lemma

- We have

$$\int_{-L}^L f(x) dx = \int_{-L}^{-L+a} f(x) dx + \int_{-L+a}^L f(x) dx$$

- Since $f(x+2L) = f(x)$, the first integral is

$$\begin{aligned} \int_{-L}^{-L+a} f(x) dx &= \int_{-L}^{-L+a} f(x+2L) dx \\ &= \int_{-L}^{-L+a} f(x+2L) d(x+2L) \\ &= \int_L^{L+a} f(x) dx \end{aligned}$$

Proof continued

- Therefore,

$$\begin{aligned}\int_{-L}^L f(x) dx &= \int_{-L}^{-L+a} f(x) dx + \int_{-L+a}^L f(x) dx \\ &= \int_L^{L+a} f(x) dx + \int_{-L+a}^L f(x) dx \\ &= \int_{-L+a}^{L+a} f(x) dx\end{aligned}$$

Proof continued

- Therefore,

$$\begin{aligned}\int_{-L}^L f(x) dx &= \int_{-L}^{-L+a} f(x) dx + \int_{-L+a}^L f(x) dx \\ &= \int_L^{L+a} f(x) dx + \int_{-L+a}^L f(x) dx \\ &= \int_{-L+a}^{L+a} f(x) dx\end{aligned}$$

- Proof is complete

Even and odd functions

- A function

$$f(x) \text{ is } \begin{cases} \text{even} & \text{if } f(-x) = f(x) \\ \text{odd} & \text{if } f(-x) = -f(x) \end{cases}$$

Even and odd functions

- A function

$$f(x) \text{ is } \begin{cases} \text{even} & \text{if } f(-x) = f(x) \\ \text{odd} & \text{if } f(-x) = -f(x) \end{cases}$$

- An easy yet useful fact about even/odd functions is that if $L > 0$ and the function f is integrable on $[-L, L]$, then

$$\int_{-L}^L f(x) dx = \begin{cases} 2 \int_0^L f(x) dx & \text{if } f \text{ is even} \\ 0 & \text{if } f \text{ is odd} \end{cases}$$

Orthogonal functions

- Two real valued functions $f(x)$ and $g(x)$ are called **orthogonal** on $[a, b]$ if

$$\int_a^b f(x)g(x)dx = 0$$

Orthogonal functions

- Two real valued functions $f(x)$ and $g(x)$ are called **orthogonal** on $[a, b]$ if

$$\int_a^b f(x)g(x)dx = 0$$

- A family $\{f_n\}$ of real valued functions is called an **orthonormal** family on $[a, b]$ if

$$\int_a^b f_m(x)f_n(x)dx = \begin{cases} 0 & \text{if } m \neq n \\ 1 & \text{if } m = n \end{cases}$$

Some orthonormal families

Proposition

Given a $L > 0$, we have

- The family $\mathcal{F}_{\sin} = \left\{ \frac{1}{\sqrt{L}} \sin \left(\frac{\pi n x}{L} \right) \right\}_{n=0}^{\infty}$ is orthonormal on $[-L, L]$
- The family $\mathcal{F}_{\cos} = \left\{ \frac{1}{\sqrt{e_n L}} \cos \left(\frac{\pi n x}{L} \right) \right\}_{n=0}^{\infty}$ is orthonormal on $[-L, L]$, where $e_0 = 2$ and $e_n = 1$ for all $n > 0$

Proof of the proposition

Really a high school level exercise!

- For nonnegative integers m and n , define

$$I_{m,n} = \frac{1}{L} \int_{-L}^L \sin\left(\frac{\pi mx}{L}\right) \sin\left(\frac{\pi nx}{L}\right) dx$$

and

$$J_{m,n} = \frac{1}{\sqrt{e_m e_n} L} \int_{-L}^L \cos\left(\frac{\pi mx}{L}\right) \cos\left(\frac{\pi nx}{L}\right) dx$$

Proof of the proposition

Really a high school level exercise!

- For nonnegative integers m and n , define

$$I_{m,n} = \frac{1}{L} \int_{-L}^L \sin\left(\frac{\pi mx}{L}\right) \sin\left(\frac{\pi nx}{L}\right) dx$$

and

$$J_{m,n} = \frac{1}{\sqrt{e_m e_n} L} \int_{-L}^L \cos\left(\frac{\pi mx}{L}\right) \cos\left(\frac{\pi nx}{L}\right) dx$$

- Then we have to show that

$$\text{both, } I_{m,n} \text{ and } J_{m,n} = \begin{cases} 0 & \text{if } m \neq n \\ 1 & \text{if } m = n \end{cases}$$

Proof of the proposition

Make use of

- Let N be any integer, then

$$\int_{-L}^L \cos(N\pi x/L) dx = \begin{cases} 2L & \text{if } N = 0 \\ 0 & \text{if } N \neq 0 \end{cases}$$

Proof of the proposition

Make use of

- Let N be any integer, then

$$\int_{-L}^L \cos(N\pi x/L) dx = \begin{cases} 2L & \text{if } N = 0 \\ 0 & \text{if } N \neq 0 \end{cases}$$

- Now use the trigonometric “product to sum” formulas

Proof of the proposition

Make use of

- Let N be any integer, then

$$\int_{-L}^L \cos(N\pi x/L) dx = \begin{cases} 2L & \text{if } N = 0 \\ 0 & \text{if } N \neq 0 \end{cases}$$

- Now use the trigonometric “product to sum” formulas
- Also, note that $\int_{-L}^L \sin(N\pi x/L) dx = 0$

A complex orthonormal family

- From $e^{(\pi i n x / L)} = \cos(\pi n x / L) + i \sin(\pi n x / L)$, we deduce that

$$\frac{1}{2L} \int_{-L}^L e^{(\pi i n x / L)} dx = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{if } n = \pm 1, \pm 2, \dots \end{cases}$$

A complex orthonormal family

- From $e^{(\pi i n x / L)} = \cos(\pi n x / L) + i \sin(\pi n x / L)$, we deduce that

$$\frac{1}{2L} \int_{-L}^L e^{(\pi i n x / L)} dx = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{if } n = \pm 1, \pm 2, \dots \end{cases}$$

- Thus, the family $\{e^{(\pi i n x / L)} / \sqrt{2L}\}_{n=-\infty}^{\infty}$ is an orthonormal family of functions.

A complex orthonormal family

- From $e^{(\pi i n x / L)} = \cos(\pi n x / L) + i \sin(\pi n x / L)$, we deduce that

$$\frac{1}{2L} \int_{-L}^L e^{(\pi i n x / L)} dx = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{if } n = \pm 1, \pm 2, \dots \end{cases}$$

- Thus, the family $\{e^{(\pi i n x / L)} / \sqrt{2L}\}_{n=-\infty}^{\infty}$ is an orthonormal family of functions.
- If the normalising factor $1/\sqrt{2L}$ is removed, then we have an orthogonal family instead

Closing remarks

- For most practical scenarios, we take $L = \pi$ so that, our last result then takes the form

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{inx} dx = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{if } n = \pm 1, \pm 2, \dots \end{cases}$$

Closing remarks

- For most practical scenarios, we take $L = \pi$ so that, our last result then takes the form

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{inx} dx = \begin{cases} 1 & \text{if } n = 0 \\ 0 & \text{if } n = \pm 1, \pm 2, \dots \end{cases}$$

- It really helps to memorise the following version of the last identity

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{imx} e^{-inx} dx = \begin{cases} 1 & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$

Trigonometric Polynomials

- Let $Q(X, Y)$ be a polynomial in two variables with real coefficients

Trigonometric Polynomials

- Let $Q(X, Y)$ be a polynomial in two variables with real coefficients
- A **Trigonometric Polynomial** (TP) $P(x)$ is a *real valued* function defined as

$$P(x) = Q(\sin x, \cos x)$$

Trigonometric Polynomials

- Let $Q(X, Y)$ be a polynomial in two variables with real coefficients
- A **Trigonometric Polynomial** (TP) $P(x)$ is a *real valued* function defined as

$$P(x) = Q(\sin x, \cos x)$$

- Thus, $P(x)$ is smooth with period 2π , i.e.,

$$\begin{aligned} P(x + 2\pi) &= Q(\sin(x + 2\pi), \cos(x + 2\pi)) \\ &= Q(\sin x, \cos x) = P(x) \end{aligned}$$

Trigonometric Polynomials

- A particular TP that we will consider here is given by

$$P_N(x) = a_0 + \sum_{n=1}^N (a_n \cos nx + b_n \sin nx),$$

where a_n, b_n are *real*, and $x \in \mathbb{R}$

Trigonometric Polynomials

- A particular TP that we will consider here is given by

$$P_N(x) = a_0 + \sum_{n=1}^N (a_n \cos nx + b_n \sin nx),$$

where a_n, b_n are *real*, and $x \in \mathbb{R}$

- Note that $\sin nx$ and $\cos nx$ can be expressed as polynomials functions of $\sin x$ and $\cos x$, deduce that P_N is indeed a TP of degree

Trigonometric Polynomials

- A particular TP that we will consider here is given by

$$P_N(x) = a_0 + \sum_{n=1}^N (a_n \cos nx + b_n \sin nx),$$

where a_n, b_n are *real*, and $x \in \mathbb{R}$

- Note that $\sin nx$ and $\cos nx$ can be expressed as polynomials functions of $\sin x$ and $\cos x$, deduce that P_N is indeed a TP of degree
- $\leq N$

Trigonometric Polynomials

Complex Version of TP

Each TP $P_N(x)$ can be expressed as

$$P_N(x) = \sum_{-N}^N c_n e^{inx}, \quad c_n \in \mathbb{C}, x \in \mathbb{R}$$

Trigonometric Polynomials

Complex Version of TP

Each TP $P_N(x)$ can be expressed as

$$P_N(x) = \sum_{-N}^N c_n e^{inx}, \quad c_n \in \mathbb{C}, x \in \mathbb{R}$$

- Write

$$\cos nx = \frac{e^{inx} + e^{-inx}}{2}, \quad \sin nx = \frac{-i(e^{inx} - e^{-inx})}{2}$$

Trigonometric Polynomials

- Then $P_N(x)$ can be expressed as

$$\begin{aligned} P_N(x) &= a_0 + \sum_{n=1}^N \left[\left(\frac{a_n - ib_n}{2} \right) e^{inx} + \left(\frac{a_n + ib_n}{2} \right) e^{-inx} \right] \\ &= \sum_{-N}^N c_n e^{inx}, \quad x \in \mathbb{R} \end{aligned}$$

Trigonometric Polynomials

- Where, one has

$$c_0 = a_0, \quad c_n = \frac{a_n - ib_n}{2}, \quad n > 0$$

Trigonometric Polynomials

- Where, one has

$$c_0 = a_0, \quad c_n = \frac{a_n - ib_n}{2}, \quad n > 0$$

- And

$$c_{-n} = \frac{a_n + ib_n}{2} = \overline{c_n}$$

Trigonometric Polynomials

- Where, one has

$$c_0 = a_0, \quad c_n = \frac{a_n - ib_n}{2}, \quad n > 0$$

- And

$$c_{-n} = \frac{a_n + ib_n}{2} = \overline{c_n}$$

- Moreover, we have that $c_n \in \mathbb{C}$ with

$$4|c_n|^2 = a_n^2 + b_n^2 \quad \text{for } n > 0$$

An Integral Identity

A Key Integral Formula

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{inx} e^{-imx} dx = \begin{cases} 1 & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$

An Integral Identity

A Key Integral Formula

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{inx} e^{-imx} dx = \begin{cases} 1 & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$

- Integrate both sides of $P_N(x)e^{-imx}$ on $[-\pi, \pi]$ to get

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} P_N(x) e^{-imx} dx = \sum_{-N}^N c_n \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(n-m)x} dx \right)$$

An Integral Identity

- Which yields

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} P_N(x) e^{-imx} dx = \begin{cases} c_m & \text{if } |m| \leq N \\ 0 & \text{if } |m| > N \end{cases}$$

An Integral Identity

- Which yields

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} P_N(x) e^{-imx} dx = \begin{cases} c_m & \text{if } |m| \leq N \\ 0 & \text{if } |m| > N \end{cases}$$

- Setting $c_m = 0$ for $|m| > N$, we thus have

$$P_N(x) = \sum_{-\infty}^{\infty} c_n e^{inx}$$

An Integral Identity

- Which yields

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} P_N(x) e^{-imx} dx = \begin{cases} c_m & \text{if } |m| \leq N \\ 0 & \text{if } |m| > N \end{cases}$$

- Setting $c_m = 0$ for $|m| > N$, we thus have

$$P_N(x) = \sum_{-\infty}^{\infty} c_n e^{inx}$$

- Where the coefficients are given by

$$c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} P_N(x) e^{-inx} dx$$

Trigonometric Series

- In the same spirit, define a **trigonometric series**

$$\sum_{-\infty}^{\infty} c_n e^{inx}, \quad c_n \in \mathbb{C}$$

Trigonometric Series

- In the same spirit, define a **trigonometric series**

$$\sum_{-\infty}^{\infty} c_n e^{inx}, \quad c_n \in \mathbb{C}$$

- Impose the condition $c_{-n} = \overline{c_n}$

Trigonometric Series

- In the same spirit, define a **trigonometric series**

$$\sum_{-\infty}^{\infty} c_n e^{inx}, \quad c_n \in \mathbb{C}$$

- Impose the condition $c_{-n} = \overline{c_n}$
- Writing $a_0 = c_0$, and $c_n = (a_n - ib_n)/2$ for $n > 0$, we get the real trigonometric series

$$a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

Fourier Series

- Given a 2π -periodic, integrable function f on $[-\pi, \pi]$, for $n \in \mathbb{Z}$, we set

$$c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx$$

Fourier Series

- Given a 2π -periodic, integrable function f on $[-\pi, \pi]$, for $n \in \mathbb{Z}$, we set

$$c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx$$

- Now, define the **Fourier series** associated with f as

$$f(x) \sim \sum_{-\infty}^{\infty} c_n e^{inx} = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

Fourier Series

- Given a 2π -periodic, integrable function f on $[-\pi, \pi]$, for $n \in \mathbb{Z}$, we set

$$c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx$$

- Now, define the **Fourier series** associated with f as

$$f(x) \sim \sum_{-\infty}^{\infty} c_n e^{inx} = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

- Numbers a_n , b_n and c_n are called the **Fourier coefficients** of $f(x)$

Remarks

- Note that for $n > 0$ the real Fourier coefficients a_n and b_n can be recovered from c_n as

$$\begin{aligned}c_n &= \frac{a_n}{2} - i\frac{b_n}{2} \\&= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx \\&= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) (\cos nx - i \sin nx) dx \\&= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \cos nxdx - \frac{i}{2\pi} \int_{-\pi}^{\pi} f(x) \sin nxdx\end{aligned}$$

Remarks

- Thus, we find that

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx, \quad a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$

Remarks

- Thus, we find that

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx, \quad a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$

- We have used a ‘ \sim ’ instead of ‘ $=$ ’ as we do not know whether

Remarks

- Thus, we find that

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx, \quad a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$

- We have used a ‘ \sim ’ instead of ‘ $=$ ’ as we do not know whether
- The series $\sum_{-\infty}^{\infty} c_n e^{inx}$ converges?

Remarks

- Thus, we find that

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx, \quad a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$

- We have used a ‘ \sim ’ instead of ‘ $=$ ’ as we do not know whether
- The series $\sum_{-\infty}^{\infty} c_n e^{inx}$ converges?
- Even if does, whether the series converges to $f(x)$?

Partial Sums of a Fourier Series

- Given $f(x) \sim \sum_{-\infty}^{\infty} c_n e^{inx}$, define the N -th partial sum $S_N(f, x) := S_N(x)$ to be the TP

$$S_N(x) = \sum_{-N}^N c_n e^{inx} = a_0 + \sum_{n=1}^N (a_n \cos nx + b_n \sin nx),$$

Partial Sums of a Fourier Series

- Given $f(x) \sim \sum_{-\infty}^{\infty} c_n e^{inx}$, define the N -th partial sum $S_N(f, x) := S_N(x)$ to be the TP

$$S_N(x) = \sum_{-N}^N c_n e^{inx} = a_0 + \sum_{n=1}^N (a_n \cos nx + b_n \sin nx),$$

- Where, for $n \leq N$, we have

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx = c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} S_N(x) e^{-inx} dx$$

Partial Sums of a Fourier Series

- Given $f(x) \sim \sum_{-\infty}^{\infty} c_n e^{inx}$, define the N -th partial sum $S_N(f, x) := S_N(x)$ to be the TP

$$S_N(x) = \sum_{-N}^N c_n e^{inx} = a_0 + \sum_{n=1}^N (a_n \cos nx + b_n \sin nx),$$

- Where, for $n \leq N$, we have

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx = c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} S_N(x) e^{-inx} dx$$

- Thus,

$$\sum_{-\infty}^{\infty} c_n e^{inx} = \lim_{N \rightarrow \infty} S_N(x)$$

Partial Sum Identities

Proposition (Integral Identities for $S_N(x)$)

Let $f(x) \sim \sum_{-\infty}^{\infty} c_n e^{inx}$, and let $S_N(x) = \sum_{-N}^N c_n e^{inx}$ be the N -th partial sum. Then

$$(i) \quad \frac{1}{2\pi} \int_{-\pi}^{\pi} |S_N(x)|^2 dx = \sum_{-N}^N |c_n|^2$$

$$(ii) \quad \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) S_N(x) dx = \sum_{-N}^N |c_n|^2.$$

Proof of the Proposition

(i) follows from the following 2 observations:

- One has

$$|S_N(x)|^2 = S_N(x) \overline{S_N(x)} = \sum_{n,m=-N}^N c_n \overline{c_m} e^{i(n-m)x}$$

Proof of the Proposition

(i) follows from the following 2 observations:

- One has

$$|S_N(x)|^2 = S_N(x) \overline{S_N(x)} = \sum_{n,m=-N}^N c_n \overline{c_m} e^{i(n-m)x}$$

- And

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} c_n \overline{c_m} e^{i(n-m)x} dx = \begin{cases} |c_n|^2 & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}$$

Proof of the Proposition

For (ii), observe that

$$\begin{aligned}\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) S_N(x) dx &= \sum_{-N}^N c_n \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{inx} dx \right) \\ &= \sum_{-N}^N c_n c_{-n} = \sum_{-N}^N c_n \overline{c_n} \\ &= \sum_{-N}^N |c_n|^2\end{aligned}$$

Bessel's Inequality & Riemann - Lebesgue Lemma

Proposition

Let f be 2π -periodic and integrable real valued function, and let $f(x) \sim \overline{\sum_{-\infty}^{\infty} c_n e^{inx}}$, then

$$(i) \quad \sum_{-\infty}^{\infty} |c_n|^2 \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx \quad (\text{Bessel's Inequality})$$

$$(ii) \quad \lim_{|n| \rightarrow \infty} c_n = 0 \quad (\text{Riemann - Lebesgue Lemma})$$

Other forms of Riemann - Lebesgue Lemma

Versions of Riemann - Lebesgue Lemma

(ii) is equivalent to

$$(1) \lim_{|n| \rightarrow \infty} \left(\int_{-\pi}^{\pi} f(x) e^{-inx} dx \right) = 0$$

$$(2) \lim_{n \rightarrow \infty} \left(\int_{-\pi}^{\pi} f(x) \cos nx dx \right) = 0$$

$$(3) \lim_{n \rightarrow \infty} \left(\int_{-\pi}^{\pi} f(x) \sin nx dx \right) = 0$$

Proof of the Proposition

For (i), observe that

$$\begin{aligned} 0 &\leq \frac{1}{2\pi} \int_{-\pi}^{\pi} |f(x) - S_N(x)|^2 dx \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} (f(x) - S_N(x)) \overline{(f(x) - S_N(x))} dx \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} (|f(x)|^2 - f(x) \overline{S_N(x)} - \overline{f(x)} S_N(x) + |S_N(x)|^2) dx \end{aligned}$$

Proof of the Proposition

At this point note that f and S_N are real valued, so that the last expression becomes

$$\begin{aligned} 0 &\leq \frac{1}{2\pi} \int_{-\pi}^{\pi} (f(x)^2 - 2f(x)S_N(x) + S_N(x)^2) dx \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx - 2 \sum_{-N}^N |c_n|^2 + \sum_{-N}^N |c_n|^2 \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx - \sum_{-N}^N |c_n|^2 \end{aligned}$$

Proof of the Proposition

- Deduce that

$$\sum_{-N}^N |c_n|^2 \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx \quad \forall \quad N$$

Proof of the Proposition

- Deduce that

$$\sum_{-N}^N |c_n|^2 \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx \quad \forall \quad N$$

- Taking $N \rightarrow \infty$ on both sides, we have (i)

Proof of the Proposition

- Deduce that

$$\sum_{-N}^N |c_n|^2 \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx \quad \forall \quad N$$

- Taking $N \rightarrow \infty$ on both sides, we have (i)
- For (ii), notice that from $c_{-n} = \overline{c_n}$, and

$$\sum_{-\infty}^{\infty} |c_n|^2 \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx < \infty$$

Proof of the Proposition

- Deduce that

$$\sum_{-N}^N |c_n|^2 \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx \quad \forall \quad N$$

- Taking $N \rightarrow \infty$ on both sides, we have (i)
- For (ii), notice that from $c_{-n} = \overline{c_n}$, and

$$\sum_{-\infty}^{\infty} |c_n|^2 \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx < \infty$$

- One has that

$$c_0^2 + 2 \sum_{n=1}^{\infty} |c_n|^2 < \infty$$

Proof of the Proposition

- That is the real numerical series $\sum_{n=1}^{\infty} |c_n|^2$ converges.

Proof of the Proposition

- That is the real numerical series $\sum_{n=1}^{\infty} |c_n|^2$ converges.
- Therefore, we necessarily have

$$\lim_{n \rightarrow \infty} |c_n|^2 = 0 \quad \implies \quad \lim_{n \rightarrow \infty} c_n = 0 \quad (\mathbf{Ex.})$$

Proof of the Proposition

- That is the real numerical series $\sum_{n=1}^{\infty} |c_n|^2$ converges.
- Therefore, we necessarily have

$$\lim_{n \rightarrow \infty} |c_n|^2 = 0 \quad \implies \quad \lim_{n \rightarrow \infty} c_n = 0 \quad (\mathbf{Ex.})$$

- Consequently, we have

$$0 = \lim_{n \rightarrow \infty} \overline{c_n} = \lim_{n \rightarrow \infty} c_{-n} = \lim_{n \rightarrow -\infty} c_n$$

Parseval's Identity

Theorem (Parseval's Theorem)

Bessel's Inequality states that if f is 2π -periodic and integrable on $[-\pi, \pi]$, then

$$\sum_{-\infty}^{\infty} |c_n|^2 \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx.$$

Parseval's result asserts something stronger, that for any f as above, we in fact have

$$\sum_{-\infty}^{\infty} |c_n|^2 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x)^2 dx.$$

Convergence Questions for FS

- We have seen that the FS associated with a 2π -periodic function f is in some sense square convergent -

$$\sum_{-\infty}^{\infty} |c_n|^2 < \infty$$

Convergence Questions for FS

- We have seen that the FS associated with a 2π -periodic function f is in some sense square convergent -

$$\sum_{-\infty}^{\infty} |c_n|^2 < \infty$$

- Naturally, we would like to know when does

$$\sum_{-\infty}^{\infty} c_n e^{inx} \quad \text{or} \quad \sum_{-\infty}^{\infty} |c_n|$$

converge?

Convergence Questions for FS

- If $\sum_{-\infty}^{\infty} c_n e^{inx} \searrow$ on $[-\pi, \pi]$, then does it converge to $f(x)$?

Convergence Questions for FS

- If $\sum_{-\infty}^{\infty} c_n e^{inx} \searrow$ on $[-\pi, \pi]$, then does it converge to $f(x)$?
- If $\sum_{-\infty}^{\infty} c_n e^{inx} = f(x)$ on $[-\pi, \pi]$, does it converge uniformly?

Convergence Questions for FS

- If $\sum_{-\infty}^{\infty} c_n e^{inx} \searrow$ on $[-\pi, \pi]$, then does it converge to $f(x)$?
- If $\sum_{-\infty}^{\infty} c_n e^{inx} = f(x)$ on $[-\pi, \pi]$, does it converge uniformly?
- For a piecewise continuous function f and a $x \in [-\pi, \pi]$, let us define

$$f(x \pm 0) = \lim_{\delta \rightarrow 0} f(x \pm \delta)$$

Convergence Questions for FS

- Define $g(x)$ on $[-\pi, \pi]$ as

$$g(x) = \frac{f(x+0) + f(x-0)}{2}$$

Convergence Questions for FS

- Define $g(x)$ on $[-\pi, \pi]$ as

$$g(x) = \frac{f(x+0) + f(x-0)}{2}$$

- Then g is well defined and $g(t) = f(t)$ if f is continuous at $x = t$

Convergence Questions for FS

- Define $g(x)$ on $[-\pi, \pi]$ as

$$g(x) = \frac{f(x+0) + f(x-0)}{2}$$

- Then g is well defined and $g(t) = f(t)$ if f is continuous at $x = t$
- If f has a discontinuity at $x = t$, then g is also discontinuous at $x = t$

Piecewise Smooth Functions

A real valued $f(x)$ is piecewise smooth on $[a, b]$ if

- f is piecewise continuous on $[a, b]$, and if f is continuous on an interval $I \subset [a, b]$, then f is also smooth on I

Piecewise Smooth Functions

A real valued $f(x)$ is piecewise smooth on $[a, b]$ if

- f is piecewise continuous on $[a, b]$, and if f is continuous on an interval $I \subset [a, b]$, then f is also smooth on I
- This, in particular, means that $f'(x)$ is piecewise continuous on $[a, b]$

Piecewise Smooth Functions

A real valued $f(x)$ is piecewise smooth on $[a, b]$ if

- f is piecewise continuous on $[a, b]$, and if f is continuous on an interval $I \subset [a, b]$, then f is also smooth on I
- This, in particular, means that $f'(x)$ is piecewise continuous on $[a, b]$
- Piecewise smooth functions are one class of functions on which Fourier theory yields rich results

Convergence Results for FS

Theorem

Let a real valued function f satisfies:

- (i) f is 2π -periodic*
- (ii) f is piecewise smooth on $[-\pi, \pi]$.*

Then the Fourier series at associated with f converges pointwise to $g(x) = \frac{f(x+0)+f(x-0)}{2}$ for all $x \in [-\pi, \pi]$.

Convergence Results for FS

Theorem

Let a real valued function f satisfies:

- (i) f is 2π -periodic*
- (ii) f is smooth on $[-\pi, \pi]$.*

Then the Fourier series associated with f converges uniformly to $f(x)$ for all $x \in [-\pi, \pi]$.

Examples and Applications

Problem 1.

Discuss the Fourier series expansion of

$$f(x) = \begin{cases} -1 & -\pi \leq x < 0 \\ 1 & 0 \leq x < \pi. \end{cases}$$

Examples and Applications

Problem 1.

Discuss the Fourier series expansion of

$$f(x) = \begin{cases} -1 & -\pi \leq x < 0 \\ 1 & 0 \leq x < \pi. \end{cases}$$

- We compute the coefficients c_n , a_n and b_n

Examples and Applications

Problem 1.

Discuss the Fourier series expansion of

$$f(x) = \begin{cases} -1 & -\pi \leq x < 0 \\ 1 & 0 \leq x < \pi. \end{cases}$$

- We compute the coefficients c_n , a_n and b_n
- We have

$$a_0 = c_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = 0$$

Solution Contd. ..

for $n > 0$, c_n is given by

$$\begin{aligned}c_n &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx \\&= \frac{1}{2\pi} \int_{-\pi}^0 f(x) e^{-inx} dx + \frac{1}{2\pi} \int_0^{\pi} f(x) e^{-inx} dx \\&= \frac{1}{2\pi} \int_{-\pi}^0 -e^{-inx} dx + \frac{1}{2\pi} \int_0^{\pi} e^{-inx} dx \\&= \frac{-i(1 - \cos n\pi)}{n\pi} = \frac{-i(1 - (-1)^n)}{n\pi}\end{aligned}$$

Solution Contd. ..

- Thus, on $[-\pi, \pi]$, we have

$$f(x) \sim \sum_{k=-\infty}^{\infty} \frac{-2i}{(2k+1)\pi} e^{((2k+1)ix)}$$

Solution Contd. ..

- Thus, on $[-\pi, \pi]$, we have

$$f(x) \sim \sum_{k=-\infty}^{\infty} \frac{-2i}{(2k+1)\pi} e^{((2k+1)ix)}$$

- And the real version can be given by

$$f(x) \sim \sum_{k=0}^{\infty} \left(\frac{4}{(2k+1)\pi} \right) \sin((2k+1)x)$$

Solution Contd. ..

- Thus, on $[-\pi, \pi]$, we have

$$f(x) \sim \sum_{k=-\infty}^{\infty} \frac{-2i}{(2k+1)\pi} e^{((2k+1)ix)}$$

- And the real version can be given by

$$f(x) \sim \sum_{k=0}^{\infty} \left(\frac{4}{(2k+1)\pi} \right) \sin((2k+1)x)$$

- Note that ' \sim ' can be replaced by '=' as long as $x \notin \{0, \pm\pi\}$

Solution Contd. ..

- From the periodicity property of the integrals of periodic functions, we find for any $a \in \mathbb{R}$ that

$$c_n = \int_{-\pi}^{\pi} f(x) dx = \int_{-\pi+a}^{\pi+a} f(x) dx$$

Solution Contd. ..

- From the periodicity property of the integrals of periodic functions, we find for any $a \in \mathbb{R}$ that

$$c_n = \int_{-\pi}^{\pi} f(x) dx = \int_{-\pi+a}^{\pi+a} f(x) dx$$

- Therefore, the series formula for $f(x)$ is valid for any $x \in \mathbb{R} / \{n\pi : n \in \mathbb{Z}\}$

Examples and Applications

Problem 2.

Find the Fourier expansion of $f(x) = x$ in $[0, 2\pi)$, and derive that

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}.$$

- Note that f is CD, hence the FS of f converges to f everywhere on $[0, 2\pi]$

Examples and Applications

Problem 2.

Find the Fourier expansion of $f(x) = x$ in $[0, 2\pi)$, and derive that

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}.$$

- Note that f is CD, hence the FS of f converges to f everywhere on $[0, 2\pi]$
- As before, we compute

$$a_0 = c_0 = \frac{1}{2\pi} \int_0^{2\pi} x dx = \pi$$

Solution Contd. ..

for $n > 0$, c_n is given by (integrating by parts)

$$\begin{aligned}c_n &= \frac{1}{2\pi} \int_0^{2\pi} x e^{-inx} dx \\&= \frac{1}{2\pi} \left(x \int e^{-inx} dx \right) \Big|_0^{2\pi} - \frac{1}{2\pi} \int_0^{2\pi} \left(\int e^{-inx} dx \right) dx \\&= \frac{1}{2\pi} \frac{x i e^{-inx}}{n} \Big|_0^{2\pi} + \frac{1}{2ni\pi} \int_0^{2\pi} e^{-inx} dx \\&= \frac{i}{n} + 0\end{aligned}$$

Solution Contd. ..

- Thus,

$$x = \pi + \sum_{\substack{-\infty \\ n \neq 0}}^{\infty} \frac{i}{n} e^{inx}$$

Solution Contd. ..

- Thus,

$$x = \pi + \sum_{\substack{-\infty \\ n \neq 0}}^{\infty} \frac{i}{n} e^{inx}$$

- Real version,

$$x = \pi + \sum_{n=1}^{\infty} \frac{-2}{n} \sin nx$$

Solution Contd. ..

- Thus,

$$x = \pi + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{i}{n} e^{inx}$$

- Real version,

$$x = \pi + \sum_{n=1}^{\infty} \frac{-2}{n} \sin nx$$

- Parseval's Identity, now yields

$$\frac{1}{2\pi} \int_0^{2\pi} x^2 dx = \pi^2 + \sum_{\substack{n=-\infty \\ n \neq 0}}^{\infty} \frac{1}{n^2}$$

Solution Contd. ..

- Therefore,

$$\frac{4\pi^2}{3} = \pi^2 + 2 \sum_{n=1}^{\infty} \frac{1}{n^2}$$

Solution Contd. ..

- Therefore,

$$\frac{4\pi^2}{3} = \pi^2 + 2 \sum_{n=1}^{\infty} \frac{1}{n^2}$$

- Upon rearranging, we have

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

Fourier Series - A worked out Example

Problem.


Consider the 2π -periodic function f given by $f(x) = 0$ on $[-\pi, 0)$; and $f(x) = x$ on $[0, \pi)$.

- (i) Sketch the graph of $f(x)$ in $[-3\pi, 3\pi]$.
- (ii) Find the Fourier Series of f .
- (iii) Pick appropriate values of x to show that

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots, \text{ and.}$$

$$\frac{\pi^2}{8} = 1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \cdots.$$

Solution - The Graph of f



Solution

For part (ii), we compute the Fourier coefficients, c_n , a_n and b_n .

$$\begin{aligned}a_0 = c_0 &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx \\&= \frac{1}{2\pi} \int_0^{\pi} x dx \\&= \frac{\pi}{4}.\end{aligned}$$

Solution

For $n \neq 0$,

$$\begin{aligned}c_n &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx = \frac{1}{2\pi} \int_0^{\pi} x e^{-inx} dx \\&= \frac{1}{2\pi} \left[x \cdot \frac{e^{-inx}}{-in} \right]_0^{\pi} - \frac{1}{2\pi} \int_0^{\pi} \frac{e^{-inx}}{-in} dx \\&= \frac{i \cos n\pi}{2n} - \frac{1 - \cos n\pi}{2\pi n^2}.\end{aligned}$$

Solution

- Thus, for $n > 0$, we have that

$$a_n = \begin{cases} 0, & n - \text{even} \\ -\frac{2}{\pi n^2}, & n - \text{odd}, \end{cases} \quad b_n = \frac{(-1)^{n+1}}{n}$$

Solution

- Thus, for $n > 0$, we have that

$$a_n = \begin{cases} 0, & n - \text{even} \\ -\frac{2}{\pi n^2}, & n - \text{odd}, \end{cases} \quad b_n = \frac{(-1)^{n+1}}{n}$$

- Therefore,

$$f(x) \sim \frac{\pi}{4} - \frac{2}{\pi} \sum_{k=0}^{\infty} \frac{\cos(2k+1)x}{(2k+1)^2} + \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin nx$$

Solution

- Since, only points of discontinuity of f are *odd multiples of π*

Solution

- Since, only points of discontinuity of f are *odd multiples of π*
- Deduce that for $x \neq (2m+1)\pi$, one has

$$\begin{aligned} f(x) = & \frac{\pi}{4} - \frac{2}{\pi} \left(\cos x + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right) \\ & + \left(\sin x - \frac{\sin 2x}{2} + \frac{\sin 3x}{3} - \frac{\sin 4x}{4} + \dots \right) \end{aligned}$$

Solution

- Since, only points of discontinuity of f are *odd multiples of π*
- Deduce that for $x \neq (2m+1)\pi$, one has

$$f(x) = \frac{\pi}{4} - \frac{2}{\pi} \left(\cos x + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right) \\ + \left(\sin x - \frac{\sin 2x}{2} + \frac{\sin 3x}{3} - \frac{\sin 4x}{4} + \dots \right)$$

- For part (iii), set $x = \pi/2$ and $x = 0$, respectively

Practice Problems

Problems.

Repeat (i) and (ii) of the last problem, and pick x suitably to find sums of interesting series.

1. $f(x) = x^2$ for $-\pi \leq x \leq \pi$

2. f is given by

$$\begin{cases} \pi - x, & 0 \leq x < \pi \\ 0, & \pi \leq x < 2\pi. \end{cases}$$

The Gamma Function - $\Gamma(x)$

Gamma Function

For $0 < x < \infty$, define the **Gamma** function $\Gamma(x)$ is defined as

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt.$$

The integral converges for all x in the indicated range.

Convergence of $\Gamma(x)$

If $0 < x \leq 1$, then

$$\begin{aligned}\Gamma(x) &= \int_0^{\infty} t^{x-1} e^{-t} dt \\ &= \int_0^1 t^{x-1} e^{-t} dt + \int_1^{\infty} t^{x-1} e^{-t} dt \\ &\leq \int_0^1 t^{x-1} dt + \int_1^{\infty} e^{-t} dt \\ &= \frac{1}{x} + \frac{1}{e} < \infty.\end{aligned}$$

Convergence of $\Gamma(x)$

If $x > 1$, then t^{x-1} is *dominated* by $e^{t/2}$ in $[0, \infty)$, meaning that there is a constant $B := B(x) < \infty$, depending on x such that

$$t^{x-1} < e^{t/2} \quad \forall \quad x \geq B$$

Convergence of $\Gamma(x)$

If $x > 1$, then t^{x-1} is *dominated* by $e^{t/2}$ in $[0, \infty)$, meaning that there is a constant $B := B(x) < \infty$, depending on x such that

$$t^{x-1} < e^{t/2} \quad \forall \quad x \geq B$$

Thus,

$$\begin{aligned}\Gamma(x) &= \int_0^\infty t^{x-1} e^{-t} dt \\ &= \int_0^B t^{x-1} e^{-t} dt + \int_B^\infty t^{x-1} e^{-t} dt \\ &\leq \text{a finite no.} + \int_B^\infty e^{-t/2} dt < \infty\end{aligned}$$

Properties of $\Gamma(x)$

Theorem

We have

- (a) $\Gamma(x+1) = x\Gamma(x) \quad \forall \quad 0 < x < \infty.$
- (b) $\Gamma(n+1) = n!$ for all $n = 1, 2, 3, \dots$
- (c) $\Gamma(x)$ is differentiable on $(0, \infty)$ (proving this is beyond the scope of this course).

Proof of the Theorem

For (a), integrate by parts, taking t^x as the first function, we have

$$\begin{aligned}\Gamma(x+1) &= \int_0^\infty t^x e^{-t} dt \\ &= \left[-\frac{t^x}{e^t} \right]_0^\infty + x \int_0^\infty t^{x-1} e^{-t} dt \\ &= x\Gamma(x).\end{aligned}$$

For part (b), use induction on n and then part (a).

The Beta function - $\beta(x, y)$

Beta Function

For $x > 0$ and $y > 0$, the *Beta* function $\beta(x, y)$ is defined as

$$\beta(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt.$$

The Beta function - $\beta(x, y)$

Beta Function

For $x > 0$ and $y > 0$, the *Beta* function $\beta(x, y)$ is defined as

$$\beta(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt.$$

Theorem

For $x > 0$ and $y > 0$, one has

$$\beta(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}.$$

Proof of the Theorem

Note that by invoking Fubini's Theorem, one has

$$\begin{aligned}\Gamma(x)\Gamma(y) &= \left(\int_0^\infty t^{x-1} e^{-t} dt \right) \left(\int_0^\infty s^{y-1} e^{-s} ds \right) \\ &= \int_0^\infty \int_0^\infty t^{x-1} s^{y-1} e^{-(s+t)} dt ds\end{aligned}$$

Proof of the Theorem

Note that by invoking Fubini's Theorem, one has

$$\begin{aligned}\Gamma(x)\Gamma(y) &= \left(\int_0^\infty t^{x-1} e^{-t} dt \right) \left(\int_0^\infty s^{y-1} e^{-s} ds \right) \\ &= \int_0^\infty \int_0^\infty t^{x-1} s^{y-1} e^{-(s+t)} dt ds\end{aligned}$$

Set $t = uv$, and $s = u(1 - v)$, so that the Jacobian $\partial(s, t)/\partial(u, v)$ is given by

$$\frac{\partial(s, t)}{\partial(u, v)} = \begin{vmatrix} \partial s / \partial u & \partial s / \partial v \\ \partial t / \partial u & \partial t / \partial v \end{vmatrix} = \begin{vmatrix} 1 - v & -u \\ v & u \end{vmatrix} = u$$

Proof of the Theorem

Thus, by invoking Fubini again, we have

$$\begin{aligned}\Gamma(x)\Gamma(y) &= \int_0^1 \int_0^\infty (uv)^{x-1} (u(1-v))^{y-1} e^{-u} u \, du \, dv \\ &= \int_0^1 \int_0^\infty v^{x-1} (1-v)^{y-1} u^{x+y-1} e^{-u} \, du \, dv \\ &= \left(\int_0^\infty u^{x+y-1} e^{-u} \, du \right) \left(\int_0^1 v^{x-1} (1-v)^{y-1} \, dv \right) \\ &= \Gamma(x+y) \beta(x, y).\end{aligned}$$

Consequences of $\Gamma - \beta$ formula

- Observe that

$$\beta(x, y) = 2 \int_0^{\pi/2} (\sin \theta)^{2x-1} (\cos \theta)^{2y-1} d\theta$$

Consequences of $\Gamma - \beta$ formula

- Observe that

$$\beta(x, y) = 2 \int_0^{\pi/2} (\sin \theta)^{2x-1} (\cos \theta)^{2y-1} d\theta$$

- To see this, substitute $t = \sin^2 \theta$, and an easy calculation yields the result

Consequences of $\Gamma - \beta$ formula

- Observe that

$$\beta(x, y) = 2 \int_0^{\pi/2} (\sin \theta)^{2x-1} (\cos \theta)^{2y-1} d\theta$$

- To see this, substitute $t = \sin^2 \theta$, and an easy calculation yields the result
- The last formula gives for $x = y = 1/2$

$$\frac{\Gamma(1/2)\Gamma(1/2)}{\Gamma(1)} = \beta(1/2, 1/2) = 2 \int_0^{\pi/2} d\theta = \pi$$

Consequences of $\Gamma - \beta$ formula

- Note that

$$\Gamma(1) = \int_0^1 e^{-t} dt = 1$$

Consequences of $\Gamma - \beta$ formula

- Note that

$$\Gamma(1) = \int_0^1 e^{-t} dt = 1$$

- Thus $\Gamma(1/2)^2 = \pi$

Consequences of $\Gamma - \beta$ formula

- Note that

$$\Gamma(1) = \int_0^1 e^{-t} dt = 1$$

- Thus $\Gamma(1/2)^2 = \pi$
- Since $\Gamma(x)$ is nonnegative, deduce that

$$\Gamma(1/2) = \sqrt{\pi}$$

Consequences of $\Gamma - \beta$ formula

- Now, substitute $t = s^2$ in the formula for $\Gamma(x)$ to get

$$\Gamma(x) = 2 \int_0^\infty s^{2x-1} e^{-s^2} ds$$

Consequences of $\Gamma - \beta$ formula

- Now, substitute $t = s^2$ in the formula for $\Gamma(x)$ to get

$$\Gamma(x) = 2 \int_0^\infty s^{2x-1} e^{-s^2} ds$$

- In the particular case $x = 1/2$, we have

$$\int_0^\infty e^{-s^2} ds = \Gamma(1/2)/2 = \sqrt{\pi}/2$$

Consequences of $\Gamma - \beta$ formula

- Now, substitute $t = s^2$ in the formula for $\Gamma(x)$ to get

$$\Gamma(x) = 2 \int_0^\infty s^{2x-1} e^{-s^2} ds$$

- In the particular case $x = 1/2$, we have

$$\int_0^\infty e^{-s^2} ds = \Gamma(1/2)/2 = \sqrt{\pi}/2$$

- Thus

$$\int_{-\infty}^\infty e^{-s^2} ds = 2 \int_0^\infty e^{-s^2} ds = \sqrt{\pi}$$

Good Luck!