

EP 1027: Maxwell's Equations and Electromagnetic Waves

Instructor: Shubho Roy¹
(Dept. of Physics)

Lecture 3

March 29, 2019

¹Office: C 313 D

Email: sroy@iith.ac.in

Office hrs. - Walk in or by Email appointment

Agenda

Agenda

- ▶ Recap of Lecture 2

Agenda

- ▶ Recap of Lecture 2
- ▶ Useful Vector identities

Agenda

- ▶ Recap of Lecture 2
- ▶ Useful Vector identities
- ▶ Potentials for curl free and divergenceless fields

Agenda

- ▶ Recap of Lecture 2
- ▶ Useful Vector identities
- ▶ Potentials for curl free and divergenceless fields
- ▶ Generic vector fields: Helmholtz theorem

Agenda

- ▶ Recap of Lecture 2
- ▶ Useful Vector identities
- ▶ Potentials for curl free and divergenceless fields
- ▶ Generic vector fields: Helmholtz theorem
- ▶ Geometric meaning of Gradient, Divergence, Curl

Agenda

- ▶ Recap of Lecture 2
- ▶ Useful Vector identities
- ▶ Potentials for curl free and divergenceless fields
- ▶ Generic vector fields: Helmholtz theorem
- ▶ Geometric meaning of Gradient, Divergence, Curl
- ▶ Integration of vector fields: Gauss' and Stokes' Theorem

Agenda

- ▶ Recap of Lecture 2
- ▶ Useful Vector identities
- ▶ Potentials for curl free and divergenceless fields
- ▶ Generic vector fields: Helmholtz theorem
- ▶ Geometric meaning of Gradient, Divergence, Curl
- ▶ Integration of vector fields: Gauss' and Stokes' Theorem
- ▶ Application of Gauss divergence theorem: Continuity equation for conservation of mass or charge

References/Readings

References/Readings

- ▶ Griffiths, D.J., **Introduction to Electrodynamics, Ch. 1**
- ▶ Spiegel M.R., **Schaum's Outline of Vector Analysis**
- ▶ Boas, M. L., **Mathematical Methods in the Physical Sciences Ch. 6**
- ▶ Arfken, G. B., **Mathematical Methods for Physicists Ch.3**

Cartesian Vectors: General treatment

- ▶ Vector: Objects whose components transform under rotation of coordinate axes as

$$a'_i = O_{ij}a_j$$

Cartesian Vectors: General treatment

- ▶ Vector: Objects whose components transform under rotation of coordinate axes as

$$a'_i = O_{ij}a_j$$

- ▶ Scalar: Objects w/o free indices, remain unchanged under rotations

$$\phi' = \phi,$$

e.g., $\phi = \mathbf{a} \cdot \mathbf{b}$ (Check), mass, time, norm/magnitude of a vector, . . .

Cartesian Vectors: General treatment

- ▶ Vector: Objects whose components transform under rotation of coordinate axes as

$$a'_i = O_{ij}a_j$$

- ▶ Scalar: Objects w/o free indices, remain unchanged under rotations

$$\phi' = \phi,$$

e.g., $\phi = \mathbf{a} \cdot \mathbf{b}$ (Check), mass, time, norm/magnitude of a vector, . . .

- ▶ Tensor: Objects with components having multiple indices, $T_{i_1 \dots i_p}$ (rank p tensor), transform with p -factors of O_{ij} ,

Cartesian Vectors: General treatment

- ▶ Vector: Objects whose components transform under rotation of coordinate axes as

$$a'_i = O_{ij} a_j$$

- ▶ Scalar: Objects w/o free indices, remain unchanged under rotations

$$\phi' = \phi,$$

e.g., $\phi = \mathbf{a} \cdot \mathbf{b}$ (Check), mass, time, norm/magnitude of a vector, . . .

- ▶ Tensor: Objects with components having multiple indices, $T_{i_1 \dots i_p}$ (rank p tensor), transform with p -factors of O_{ij} ,
- ▶ Rank 2 tensor: Under coordinate axes rotation,

$$T'_{ij} = O_{il} O_{jm} T_{lm},$$

e.g., Outer product of two vectors- $a^i b^j$; Kronecker delta- δ_{ij} ; Moment of Inertia tensor, $I_{ij} = m(\delta_{ij} x_k x_k - x_i x_j)$.

Cartesian Vectors: General treatment

Cartesian Vectors: General treatment

- ▶ Levi-Civita tensor, ϵ_{ijk} is a rank 3 tensor

$$\begin{aligned}\epsilon'_{ijk} &= O_{il}O_{jm}O_{kn}\epsilon_{lmn} \\ &= |O|\epsilon_{ijk} \\ &= \epsilon_{ijk}.\end{aligned}$$

Cartesian Vectors: General treatment

- ▶ Levi-Civita tensor, ϵ_{ijk} is a rank 3 tensor

$$\begin{aligned}\epsilon'_{ijk} &= O_{il}O_{jm}O_{kn}\epsilon_{lmn} \\ &= |O|\epsilon_{ijk} \\ &= \epsilon_{ijk}.\end{aligned}$$

- ▶ Invariant tensors: δ_{ij} , ϵ_{ijk}

Cartesian Vectors: General treatment

- ▶ Levi-Civita tensor, ϵ_{ijk} is a rank 3 tensor

$$\begin{aligned}\epsilon'_{ijk} &= O_{il}O_{jm}O_{kn}\epsilon_{lmn} \\ &= |O|\epsilon_{ijk} \\ &= \epsilon_{ijk}.\end{aligned}$$

- ▶ Invariant tensors: δ_{ij} , ϵ_{ijk}
- ▶ Check: Cross-product of two tensors give a vector:

$$(\mathbf{a} \times \mathbf{b})_k = \epsilon_{ijk}a_ib_j$$

Scalar field Differentiation: Gradient

Scalar field Differentiation: Gradient

- Components are functions of position:

$$\phi(\mathbf{x}), V_k(\mathbf{x}), T_{ij}(\mathbf{x})$$

At each point in space, \mathbf{x} , there is a scalar/vector/tensor.

Scalar field Differentiation: Gradient

- Components are functions of position:

$$\phi(\mathbf{x}), V_k(\mathbf{x}), T_{ij}(\mathbf{x})$$

At each point in space, \mathbf{x} , there is a scalar/vector/tensor.

- Examples:

$$\rho(\mathbf{x}), E_k(x), \chi_{ij}(\mathbf{x}).$$

Scalar field Differentiation: Gradient

- Components are functions of position:

$$\phi(\mathbf{x}), V_k(\mathbf{x}), T_{ij}(\mathbf{x})$$

At each point in space, \mathbf{x} , there is a scalar/vector/tensor.

- Examples:

$$\rho(\mathbf{x}), E_k(x), \chi_{ij}(\mathbf{x}).$$

- Gradient of a scalar

$$\begin{aligned}\phi(\mathbf{x} + d\mathbf{x}) - \phi(\mathbf{x}) &= dx_1 \frac{\partial \phi}{\partial x_1} + dx_2 \frac{\partial \phi}{\partial x_2} + dx_3 \frac{\partial \phi}{\partial x_3} \\ &= dx_i \frac{\partial \phi}{\partial x_i} \\ &= d\mathbf{x} \cdot (\nabla \phi),\end{aligned}$$

where we have introduced the **Gradient operator**, ∇ ,

Scalar field Differentiation: Gradient

- Components are functions of position:

$$\phi(\mathbf{x}), V_k(\mathbf{x}), T_{ij}(\mathbf{x})$$

At each point in space, \mathbf{x} , there is a scalar/vector/tensor.

- Examples:

$$\rho(\mathbf{x}), E_k(x), \chi_{ij}(\mathbf{x}).$$

- Gradient of a scalar

$$\begin{aligned}\phi(\mathbf{x} + d\mathbf{x}) - \phi(\mathbf{x}) &= dx_1 \frac{\partial \phi}{\partial x_1} + dx_2 \frac{\partial \phi}{\partial x_2} + dx_3 \frac{\partial \phi}{\partial x_3} \\ &= dx_i \frac{\partial \phi}{\partial x_i} \\ &= d\mathbf{x} \cdot (\nabla \phi),\end{aligned}$$

where we have introduced the **Gradient operator**, ∇ ,



$$\nabla \equiv \hat{\mathbf{e}}_1 \frac{\partial}{\partial x_1} + \hat{\mathbf{e}}_2 \frac{\partial}{\partial x_2} + \hat{\mathbf{e}}_3 \frac{\partial}{\partial x_3} = \hat{\mathbf{e}}_k \frac{\partial}{\partial x_k}.$$

Vector differentiation: Divergence and Curl

Vector differentiation: Divergence and Curl

- ▶ Since, ∇ acts like a vector, we can construct a scalar by taking the inner product, with a vector field, $\mathbf{A}(\mathbf{x})$,

$$\nabla \cdot \mathbf{A}(\mathbf{x}) = \partial_k A_k.$$

Vector differentiation: Divergence and Curl

- ▶ Since, ∇ acts like a vector, we can construct a scalar by taking the inner product, with a vector field, $\mathbf{A}(\mathbf{x})$,

$$\nabla \cdot \mathbf{A}(\mathbf{x}) = \partial_k A_k.$$

- ▶ Call it **Divergence of the vector field**,

$$\nabla \cdot \mathbf{A} = \frac{\partial A_1}{\partial x_1} + \frac{\partial A_2}{\partial x_2} + \frac{\partial A_3}{\partial x_3}$$

Vector differentiation: Divergence and Curl

- ▶ Since, ∇ acts like a vector, we can construct a scalar by taking the inner product, with a vector field, $\mathbf{A}(\mathbf{x})$,

$$\nabla \cdot \mathbf{A}(\mathbf{x}) = \partial_k A_k.$$

- ▶ Call it **Divergence of the vector field**,

$$\nabla \cdot \mathbf{A} = \frac{\partial A_1}{\partial x_1} + \frac{\partial A_2}{\partial x_2} + \frac{\partial A_3}{\partial x_3}$$

- ▶ We can further create a vector by taking the cross product of ∇ and $\mathbf{A}(\mathbf{x})$,

$$\nabla \times \mathbf{A} = (\nabla \times \mathbf{A})_k \hat{\mathbf{e}}_k,$$

$$\begin{aligned} (\nabla \times \mathbf{A})_k &= \epsilon_{ijk} \partial_i A_j \\ &= \epsilon_{kij} \partial_i A_j. \end{aligned}$$

Vector differentiation: Divergence and Curl

- ▶ Since, ∇ acts like a vector, we can construct a scalar by taking the inner product, with a vector field, $\mathbf{A}(\mathbf{x})$,

$$\nabla \cdot \mathbf{A}(\mathbf{x}) = \partial_k A_k.$$

- ▶ Call it **Divergence of the vector field**,

$$\nabla \cdot \mathbf{A} = \frac{\partial A_1}{\partial x_1} + \frac{\partial A_2}{\partial x_2} + \frac{\partial A_3}{\partial x_3}$$

- ▶ We can further create a vector by taking the cross product of ∇ and $\mathbf{A}(\mathbf{x})$,

$$\nabla \times \mathbf{A} = (\nabla \times \mathbf{A})_k \hat{\mathbf{e}}_k,$$

$$\begin{aligned} (\nabla \times \mathbf{A})_k &= \epsilon_{ijk} \partial_i A_j \\ &= \epsilon_{kij} \partial_i A_j. \end{aligned}$$

- ▶ call it **Curl of the vector field**

Vector differentiation: Divergence and Curl

Vector differentiation: Divergence and Curl



$$\nabla r^n = n r^{n-2} \mathbf{x}, \quad r = |\mathbf{x}|.$$

Vector differentiation: Divergence and Curl



$$\nabla r^n = n r^{n-2} \mathbf{x}, \quad r = |\mathbf{x}|.$$



$$\nabla \cdot \mathbf{x} = \frac{\partial x_1}{\partial x_1} + \frac{\partial x_2}{\partial x_2} + \frac{\partial x_3}{\partial x_3} = 3$$

Vector differentiation: Divergence and Curl



$$\nabla r^n = n r^{n-2} \mathbf{x}, \quad r = |\mathbf{x}|.$$



$$\nabla \cdot \mathbf{x} = \frac{\partial x_1}{\partial x_1} + \frac{\partial x_2}{\partial x_2} + \frac{\partial x_3}{\partial x_3} = 3$$



$$(\nabla \times \mathbf{x})_k = \epsilon_{ijk} \partial_i x_j = \epsilon_{ijk} \delta_{ij} = 0.$$

Vector differentiation: Divergence and Curl



$$\nabla r^n = n r^{n-2} \mathbf{x}, \quad r = |\mathbf{x}|.$$



$$\nabla \cdot \mathbf{x} = \frac{\partial x_1}{\partial x_1} + \frac{\partial x_2}{\partial x_2} + \frac{\partial x_3}{\partial x_3} = 3$$



$$(\nabla \times \mathbf{x})_k = \epsilon_{ijk} \partial_i x_j = \epsilon_{ijk} \delta_{ij} = 0.$$

- ▶ Can define a double derivative thru the inner product, the **Laplacian**

$$\nabla^2 \equiv \nabla \cdot \nabla = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \frac{\partial^2}{\partial x_3^2}.$$

Useful Identity I

Useful Identity I

- ▶ Product of two Levi-Civita's with one index repeated ("contraction")

$$\epsilon_{ijk}\epsilon_{lmk} = \delta_{il}\delta_{jm} - \delta_{im}\delta_{jl}. \quad (1)$$

k is the repeated index.

Useful Identity I

- ▶ Product of two Levi-Civita's with one index repeated (“contraction”)

$$\epsilon_{ijk}\epsilon_{lmk} = \delta_{il}\delta_{jm} - \delta_{im}\delta_{jl}. \quad (1)$$

k is the repeated index.

- ▶ Use it to derive $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B})$ i.e. the “BAC minus CAB” rule for vector triple products.

Useful Identity I

- ▶ Product of two Levi-Civita's with one index repeated (“contraction”)

$$\epsilon_{ijk}\epsilon_{lmk} = \delta_{il}\delta_{jm} - \delta_{im}\delta_{jl}. \quad (1)$$

k is the repeated index.

- ▶ Use it to derive $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B})$ i.e. the “BAC minus CAB” rule for vector triple products.

▶

$$\begin{aligned} [\mathbf{A} \times (\mathbf{B} \times \mathbf{C})]_k &= \epsilon_{ijk} A_i \underbrace{(\mathbf{B} \times \mathbf{C})_j} \\ &= \epsilon_{ijk} A_i (\epsilon_{lmj} B_l C_m) \\ &= \epsilon_{ijk} \epsilon_{lmj} A_i B_l C_m \\ &= \epsilon_{kij} \epsilon_{lmj} A_i B_l C_m \\ &= (\delta_{kl} \delta_{im} - \delta_{km} \delta_{il}) A_i B_l C_m \\ &= (\delta_{kl} B_l) A_i (\delta_{im} C_m) - (\delta_{km} C_m) A_i (\delta_{il} B_l) \\ &= B_k A_i C_i - C_k A_i B_i = B_k (\mathbf{A} \cdot \mathbf{C}) - C_k (\mathbf{A} \cdot \mathbf{B}) \end{aligned}$$

where in the fourth line we have used the fact that the ϵ -tensor is unchanged under cyclic permutation of its indices $\epsilon_{ijk} = \epsilon_{kij}$. In the 5th line we have used the identity (1) and in going from 7th to 8th line we have used,

$$\delta_{kl} B_l = B_k, \quad \delta_{im} C_m = C_i, \quad \delta_{km} C_m = C_k, \quad \delta_{il} B_l = B_i,$$

Useful identities II and III

Useful identities II and III

- Curl of gradient of a scalar field vanishes,

$$\nabla \times (\nabla \Phi(\mathbf{x})) = 0 \quad (2)$$

Useful identities II and III

- Curl of gradient of a scalar field vanishes,

$$\nabla \times (\nabla \Phi(\mathbf{x})) = 0 \quad (2)$$

- Proof:

$$\begin{aligned} [\nabla \times (\nabla \Phi)]_k &= \epsilon_{ijk} \partial_i (\nabla \Phi)_j \\ &= \epsilon_{ijk} \partial_i (\partial_j \Phi) \\ &= \epsilon_{ijk} \frac{\partial^2 \Phi}{\partial x_i \partial x_j} = 0, \end{aligned}$$

because ϵ_{ijk} is antisymmetric in i, j while $\frac{\partial^2 \Phi}{\partial x_i \partial x_j}$ is symmetric in i, j , as order of derivatives do not matter. The product of symmetric and antisymmetric objects in i, j vanishes.

Useful identities II and III

- Curl of gradient of a scalar field vanishes,

$$\nabla \times (\nabla \Phi(\mathbf{x})) = 0 \quad (2)$$

- Proof:

$$\begin{aligned} [\nabla \times (\nabla \Phi)]_k &= \epsilon_{ijk} \partial_i (\nabla \Phi)_j \\ &= \epsilon_{ijk} \partial_i (\partial_j \Phi) \\ &= \epsilon_{ijk} \frac{\partial^2 \Phi}{\partial x_i \partial x_j} = 0, \end{aligned}$$

because ϵ_{ijk} is antisymmetric in i, j while $\frac{\partial^2 \Phi}{\partial x_i \partial x_j}$ is symmetric in i, j , as order of derivatives do not matter. The product of symmetric and antisymmetric objects in i, j vanishes.

- Divergence of a curl of a vector field vanishes,

$$\nabla \cdot (\nabla \times \mathbf{A}) = 0 \quad (3)$$

Useful identities II and III

- ▶ Curl of gradient of a scalar field vanishes,

$$\nabla \times (\nabla \Phi(\mathbf{x})) = 0 \quad (2)$$

- ▶ Proof:

$$\begin{aligned} [\nabla \times (\nabla \Phi)]_k &= \epsilon_{ijk} \partial_i (\nabla \Phi)_j \\ &= \epsilon_{ijk} \partial_i (\partial_j \Phi) \\ &= \epsilon_{ijk} \frac{\partial^2 \Phi}{\partial x_i \partial x_j} = 0, \end{aligned}$$

because ϵ_{ijk} is antisymmetric in i, j while $\frac{\partial^2 \Phi}{\partial x_i \partial x_j}$ is symmetric in i, j , as order of derivatives do not matter. The product of symmetric and antisymmetric objects in i, j vanishes.

- ▶ Divergence of a curl of a vector field vanishes,

$$\nabla \cdot (\nabla \times \mathbf{A}) = 0 \quad (3)$$

- ▶ Proof:

$$\begin{aligned} \nabla \cdot (\nabla \times \mathbf{A}) &= \partial_k (\nabla \times \mathbf{A})_k \\ &= \partial_k (\epsilon_{ijk} \partial_i A_j) = \epsilon_{ijk} \frac{\partial^2 A_j}{\partial x_k \partial x_i} = 0, \end{aligned}$$

Irrotational and Solenoidal Vector fields: Scalar and Vector potentials

Irrotational and Solenoidal Vector fields: Scalar and Vector potentials

- ▶ If \mathbf{C} is curl-free vector field (also called irrotational or conservative), i.e., if

$$\nabla \times \mathbf{C} = 0,$$

then \mathbf{C} can be expressed as the gradient of a scalar, say, Φ

$$\mathbf{C} = -\nabla\Phi.$$

Irrotational and Solenoidal Vector fields: Scalar and Vector potentials

- ▶ If \mathbf{C} is curl-free vector field (also called irrotational or conservative), i.e., if

$$\nabla \times \mathbf{C} = 0,$$

then \mathbf{C} can be expressed as the gradient of a scalar, say, Φ

$$\mathbf{C} = -\nabla\Phi.$$

- ▶ Φ is called the **Scalar potential of the conservative vector field, \mathbf{C} .**

Irrotational and Solenoidal Vector fields: Scalar and Vector potentials

- ▶ If \mathbf{C} is curl-free vector field (also called irrotational or conservative), i.e., if

$$\nabla \times \mathbf{C} = 0,$$

then \mathbf{C} can be expressed as the gradient of a scalar, say, Φ

$$\mathbf{C} = -\nabla\Phi.$$

- ▶ Φ is called the **Scalar potential of the conservative vector field, \mathbf{C} .**
- ▶ If \mathbf{B} is divergenceless vector field (also called solenoidal), i.e., if

$$\nabla \cdot \mathbf{B} = 0,$$

then, \mathbf{B} can be expressed as a divergence of another vector, say, \mathbf{A}

$$\mathbf{B} = \nabla \times \mathbf{A}.$$

Irrotational and Solenoidal Vector fields: Scalar and Vector potentials

- ▶ If \mathbf{C} is curl-free vector field (also called irrotational or conservative), i.e., if

$$\nabla \times \mathbf{C} = 0,$$

then \mathbf{C} can be expressed as the gradient of a scalar, say, Φ

$$\mathbf{C} = -\nabla\Phi.$$

- ▶ Φ is called the **Scalar potential of the conservative vector field, \mathbf{C} .**
- ▶ If \mathbf{B} is divergenceless vector field (also called solenoidal), i.e., if

$$\nabla \cdot \mathbf{B} = 0,$$

then, \mathbf{B} can be expressed as a divergence of another vector, say, \mathbf{A}

$$\mathbf{B} = \nabla \times \mathbf{A}.$$

- ▶ \mathbf{A} is called the **Vector potential of the solenoidal vector field, \mathbf{B} .**

Irrotational and Solenoidal Vector fields: Scalar and Vector potentials

- ▶ If \mathbf{C} is curl-free vector field (also called irrotational or conservative), i.e., if

$$\nabla \times \mathbf{C} = 0,$$

then \mathbf{C} can be expressed as the gradient of a scalar, say, Φ

$$\mathbf{C} = -\nabla\Phi.$$

- ▶ Φ is called the **Scalar potential of the conservative vector field, \mathbf{C} .**
- ▶ If \mathbf{B} is divergenceless vector field (also called solenoidal), i.e., if

$$\nabla \cdot \mathbf{B} = 0,$$

then, \mathbf{B} can be expressed as a divergence of another vector, say, \mathbf{A}

$$\mathbf{B} = \nabla \times \mathbf{A}.$$

- ▶ \mathbf{A} is called the **Vector potential of the solenoidal vector field, \mathbf{B} .**
- ▶ \mathbf{A} is non-unique, $\mathbf{A}' \sim \mathbf{A} + \nabla\chi$.

General Vector fields: Helmholtz theorem

General Vector fields: Helmholtz theorem

- ▶ Generic vector field, \mathbf{C}

$$\mathbf{C} = -\nabla\Phi + \nabla \times \mathbf{A}$$

General Vector fields: Helmholtz theorem

- ▶ Generic vector field, \mathbf{C}

$$\mathbf{C} = -\nabla\Phi + \nabla \times \mathbf{A}$$



$$\Phi(\mathbf{x}) = \frac{1}{4\pi} \iiint d^3\mathbf{x}' \frac{\nabla' \cdot \mathbf{C}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$
$$\mathbf{A}(\mathbf{x}) = \frac{1}{4\pi} \iiint d^3\mathbf{x}' \frac{\nabla' \times \mathbf{C}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$

General Vector fields: Helmholtz theorem

- ▶ Generic vector field, \mathbf{C}

$$\mathbf{C} = -\nabla\Phi + \nabla \times \mathbf{A}$$



$$\Phi(\mathbf{x}) = \frac{1}{4\pi} \iiint d^3\mathbf{x}' \frac{\nabla' \cdot \mathbf{C}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$
$$\mathbf{A}(\mathbf{x}) = \frac{1}{4\pi} \iiint d^3\mathbf{x}' \frac{\nabla' \times \mathbf{C}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$

- ▶ Conditions apply: $\mathbf{C} \sim \frac{1}{|\mathbf{x}|^{2+\epsilon}}, \epsilon > 0$

General Vector fields: Helmholtz theorem

- ▶ Generic vector field, \mathbf{C}

$$\mathbf{C} = -\nabla\Phi + \nabla \times \mathbf{A}$$



$$\Phi(\mathbf{x}) = \frac{1}{4\pi} \iiint d^3\mathbf{x}' \frac{\nabla' \cdot \mathbf{C}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$
$$\mathbf{A}(\mathbf{x}) = \frac{1}{4\pi} \iiint d^3\mathbf{x}' \frac{\nabla' \times \mathbf{C}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$

- ▶ Conditions apply: $\mathbf{C} \sim \frac{1}{|\mathbf{x}|^{2+\epsilon}}, \epsilon > 0$
- ▶ Powerful result: Specifying Divergence and Curl of a vector field all over can determine the vector field itself!

General Vector fields: Helmholtz theorem

- ▶ Generic vector field, \mathbf{C}

$$\mathbf{C} = -\nabla\Phi + \nabla \times \mathbf{A}$$



$$\Phi(\mathbf{x}) = \frac{1}{4\pi} \iiint d^3\mathbf{x}' \frac{\nabla' \cdot \mathbf{C}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$
$$\mathbf{A}(\mathbf{x}) = \frac{1}{4\pi} \iiint d^3\mathbf{x}' \frac{\nabla' \times \mathbf{C}(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|}$$

- ▶ Conditions apply: $\mathbf{C} \sim \frac{1}{|\mathbf{x}|^{2+\epsilon}}, \epsilon > 0$
- ▶ Powerful result: Specifying Divergence and Curl of a vector field all over can determine the vector field itself!
- ▶ Only true for 3 dim space!

Geometric meaning of gradient, divergence, curl

Geometric meaning of gradient, divergence, curl

- **Gradient:** Consider a volume element, $\Delta V = \Delta x \Delta y \Delta z$, around point, \mathbf{x}

$$\lim_{\Delta V \rightarrow 0} \frac{\oiint dS \, \hat{\mathbf{n}} \cdot \nabla \Phi(\mathbf{x})}{\Delta V} = \nabla \Phi,$$

$\hat{\mathbf{n}}$ is the unit outward normal vector on the surface S .

Geometric meaning of gradient, divergence, curl

- **Gradient:** Consider a volume element, $\Delta V = \Delta x \Delta y \Delta z$, around point, \mathbf{x}

$$\lim_{\Delta V \rightarrow 0} \frac{\oint_S dS \hat{\mathbf{n}} \cdot \nabla \phi(\mathbf{x})}{\Delta V} = \nabla \phi,$$

$\hat{\mathbf{n}}$ is the unit outward normal vector on the surface S .

- **Divergence:** Consider a volume element, $\Delta V = \Delta x \Delta y \Delta z$, around \mathbf{x} ,

$$\lim_{\Delta V \rightarrow 0} \frac{\oint_S dS \hat{\mathbf{n}} \cdot \mathbf{A}}{\Delta V} = \nabla \cdot \mathbf{A}$$

$\hat{\mathbf{n}}$ is the unit outward normal vector on the surface S . Thus,

Divergence = flux over an infinitesimal closed surface per unit volume enclosed by the surface.

Geometric meaning of gradient, divergence, curl

- **Gradient:** Consider a volume element, $\Delta V = \Delta x \Delta y \Delta z$, around point, \mathbf{x}

$$\lim_{\Delta V \rightarrow 0} \frac{\oiint dS \hat{\mathbf{n}} \Phi(\mathbf{x})}{\Delta V} = \nabla \Phi,$$

$\hat{\mathbf{n}}$ is the unit outward normal vector on the surface S .

- **Divergence:** Consider a volume element, $\Delta V = \Delta x \Delta y \Delta z$, around \mathbf{x} ,

$$\lim_{\Delta V \rightarrow 0} \frac{\oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{A}}{\Delta V} = \nabla \cdot \mathbf{A}$$

$\hat{\mathbf{n}}$ is the unit outward normal vector on the surface S . Thus,

Divergence = flux over an infinitesimal closed surface per unit volume enclosed by the surface.

- **Curl:** Consider an area element, $\Delta S_{yz} = \Delta y \Delta z$, around \mathbf{x} ,

$$\lim_{\Delta S_{yz} \rightarrow 0} \frac{\oint \mathbf{A} \cdot d\mathbf{l}}{\Delta S_{yz}} = (\nabla \times \mathbf{A})_x,$$

$d\mathbf{l}$ is the (tangential) line element. So,

Curl = Anticlockwise circulation in an infinitesimal loop per unit normal area bounded by the loop.

Integration of vector fields

Integration of vector fields

- ▶ **Gauss Divergence theorem:** If S is a closed surface enclosing a volume, V

$$\iiint_V d^3\mathbf{x} \nabla \cdot \mathbf{A} = \oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{A},$$

$\hat{\mathbf{n}}$ is the unit outward normal vector on the surface S .

Integration of vector fields

- ▶ **Gauss Divergence theorem:** If S is a closed surface enclosing a volume, V

$$\iiint_V d^3\mathbf{x} \nabla \cdot \mathbf{A} = \oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{A},$$

$\hat{\mathbf{n}}$ is the unit outward normal vector on the surface S .

- ▶ **Stokes Curl theorem:** If S is an open surface, with a boundary, C (closed curve)

$$\iint_S dS \hat{\mathbf{n}} \cdot (\nabla \times \mathbf{A}) = \oint_C d\mathbf{l} \cdot \mathbf{A}$$

Integration of vector fields

- ▶ **Gauss Divergence theorem:** If S is a closed surface enclosing a volume, V

$$\iiint_V d^3\mathbf{x} \nabla \cdot \mathbf{A} = \oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{A},$$

$\hat{\mathbf{n}}$ is the unit outward normal vector on the surface S .

- ▶ **Stokes Curl theorem:** If S is an open surface, with a boundary, C (closed curve)

$$\iint_S dS \hat{\mathbf{n}} \cdot (\nabla \times \mathbf{A}) = \oint_C d\mathbf{l} \cdot \mathbf{A}$$

- ▶ Should be thought of as vector generalizations of *Fundamental theorem* of single variable calculus:

$$\int_a^b dx \frac{df(x)}{dx} = f(b) - f(a)$$

Gauss and Stokes theorem





Figure: Pictorial representation of Gauss and Stokes Theorems.

Application of Gauss' theorem: Continuity Equation

Application of Gauss' theorem: Continuity Equation

- Consider a closed surface, S enclosing a volume, V containing a fluid of mass density (or electric charge density), ρ . The total mass/charge inside is then,

$$\iiint_V d^3\mathbf{x} \rho$$

Application of Gauss' theorem: Continuity Equation

- ▶ Consider a closed surface, S enclosing a volume, V containing a fluid of mass density (or electric charge density), ρ . The total mass/charge inside is then,

$$\iiint_V d^3\mathbf{x} \rho$$

- ▶ The amount of mass or charge flowing out per unit area perpendicular to the flow, per unit time, is called the current density, \mathbf{j} .

Application of Gauss' theorem: Continuity Equation

- ▶ Consider a closed surface, S enclosing a volume, V containing a fluid of mass density (or electric charge density), ρ . The total mass/charge inside is then,

$$\iiint_V d^3\mathbf{x} \rho$$

- ▶ The amount of mass or charge flowing out per unit area perpendicular to the flow, per unit time, is called the current density, \mathbf{j} .
- ▶ The amount of mass/charge coming out of the volume V by crossing the surface, S per unit time is equal the outward flux per unit time thru the entire surface, S

$$\oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{j}$$

Application of Gauss' theorem: Continuity Equation

- ▶ Consider a closed surface, S enclosing a volume, V containing a fluid of mass density (or electric charge density), ρ . The total mass/charge inside is then,

$$\iiint_V d^3\mathbf{x} \rho$$

- ▶ The amount of mass or charge flowing out per unit area perpendicular to the flow, per unit time, is called the current density, \mathbf{j} .
- ▶ The amount of mass/charge coming out of the volume V by crossing the surface, S per unit time is equal the outward flux per unit time thru the entire surface, S

$$\oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{j}$$

- ▶ The outflow of the fluid across the surface, S will lead to a reduce in the mass or charge inside the volume, V . This rate of decrease of mass/charge inside V is,

$$-\frac{d}{dt} \left(\iiint_V d^3\mathbf{x} \rho \right)$$

Application of Gauss' theorem: Continuity Equation

- ▶ Consider a closed surface, S enclosing a volume, V containing a fluid of mass density (or electric charge density), ρ . The total mass/charge inside is then,

$$\iiint_V d^3\mathbf{x} \rho$$

- ▶ The amount of mass or charge flowing out per unit area perpendicular to the flow, per unit time, is called the current density, \mathbf{j} .
- ▶ The amount of mass/charge coming out of the volume V by crossing the surface, S per unit time is equal the outward flux per unit time thru the entire surface, S

$$\oint_S dS \hat{\mathbf{n}} \cdot \mathbf{j}$$

- ▶ The outflow of the fluid across the surface, S will lead to a reduce in the mass or charge inside the volume, V . This rate of decrease of mass/charge inside V is,

$$-\frac{d}{dt} \left(\iiint_V d^3\mathbf{x} \rho \right)$$

- ▶ Since there are no sinks (or sources) where the fluid can disappear to (or appear from),
Amount of mass (or charge) escaped by crossing the surface, S = Amount of mass (or charge) decreased in the volume, V

Continuity Equation

Continuity Equation

- Conservation of mass or electric charge

$$\oiint_S dS \, \hat{\mathbf{n}} \cdot \mathbf{j} = -\frac{d}{dt} \left(\iiint_V d^3\mathbf{x} \, \rho \right) \quad (4)$$

Continuity Equation

- Conservation of mass or electric charge

$$\oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{j} = -\frac{d}{dt} \left(\iiint_V d^3\mathbf{x} \rho \right) \quad (4)$$

- One can convert the surface integral on the LHS into a volume integral using Gauss' theorem,

$$\oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{j} = \iiint_V d^3\mathbf{x} \nabla \cdot \mathbf{j},$$

Continuity Equation

- Conservation of mass or electric charge

$$\oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{j} = -\frac{d}{dt} \left(\iiint_V d^3\mathbf{x} \rho \right) \quad (4)$$

- One can convert the surface integral on the LHS into a volume integral using Gauss' theorem,

$$\oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{j} = \iiint_V d^3\mathbf{x} \nabla \cdot \mathbf{j},$$

- And in the RHS one can take the time-derivative from outside the volume integral to inside the volume integral,

$$-\frac{d}{dt} \left(\iiint_V d^3\mathbf{x} \rho \right) = \iiint_V d^3\mathbf{x} \left(-\frac{\partial \rho}{\partial t} \right).$$

Continuity Equation

- Conservation of mass or electric charge

$$\oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{j} = -\frac{d}{dt} \left(\iiint_V d^3\mathbf{x} \rho \right) \quad (4)$$

- One can convert the surface integral on the LHS into a volume integral using Gauss' theorem,

$$\oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{j} = \iiint_V d^3\mathbf{x} \nabla \cdot \mathbf{j},$$

- And in the RHS one can take the time-derivative from outside the volume integral to inside the volume integral,

$$-\frac{d}{dt} \left(\iiint_V d^3\mathbf{x} \rho \right) = \iiint_V d^3\mathbf{x} \left(-\frac{\partial \rho}{\partial t} \right).$$

- Thus, the conservation equation, (4), becomes,

$$\iiint_V d^3\mathbf{x} = \iiint_V d^3\mathbf{x} \left(-\frac{\partial \rho}{\partial t} \right),$$

or,

$$\begin{aligned} \iiint_V d^3\mathbf{x} \left(\frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{j} \right) &= 0, \\ \implies \frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{j} &= 0. \end{aligned}$$