

EP 1027: Maxwell's Equations and Electromagnetic Waves

Instructor: Shubho Roy¹
(Dept. of Physics)

Lecture 11

April 20, 2019

¹Email: sroy@iith.ac.in, Office: C 313 D, Office hrs: Walk in or Email Appointment

Agenda

Agenda

- ▶ EM fields in Conductors: Skin depth & Skin Effect, Plasma frequency

Agenda

- ▶ EM fields in Conductors: Skin depth & Skin Effect, Plasma frequency
- ▶ Reflection from surface of a conductor

Agenda

- ▶ EM fields in Conductors: Skin depth & Skin Effect, Plasma frequency
- ▶ Reflection from surface of a conductor
- ▶ Dispersion: Cauchy's formula

References/Readings

References/Readings

- ▶ Griffiths, D.J., **Introduction to Electrodynamics, Ch. 9**

EM waves in Conductors

EM waves in Conductors

- ▶ Conductors have free charges, ρ_f , and Ohmic current,

$$\mathbf{j}_{free} = \sigma \mathbf{E},$$

EM waves in Conductors

- ▶ Conductors have free charges, ρ_f , and Ohmic current,

$$\mathbf{j}_{free} = \sigma \mathbf{E},$$

- ▶ Maxwell's Equations in conductors (ϵ, μ, σ)

$$\nabla \cdot \mathbf{E} = \frac{\rho_{free}}{\epsilon}, \quad \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t},$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{B} = \mu\sigma \mathbf{E} + \mu\epsilon \frac{\partial \mathbf{E}}{\partial t}$$

EM waves in Conductors

- ▶ Conductors have free charges, ρ_f , and Ohmic current,

$$\mathbf{j}_{free} = \sigma \mathbf{E},$$

- ▶ Maxwell's Equations in conductors (ϵ, μ, σ)

$$\nabla \cdot \mathbf{E} = \frac{\rho_{free}}{\epsilon}, \quad \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t},$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{B} = \mu \sigma \mathbf{E} + \mu \epsilon \frac{\partial \mathbf{E}}{\partial t}$$

- ▶ Continuity Equation: Conservation of free charge

$$\nabla \cdot \mathbf{j}_{free} + \frac{\partial \rho_{free}}{\partial t} = 0,$$

$$\implies \rho(t) = e^{-(\sigma/\epsilon)t} \rho(0).$$

Charge quickly dissipates inside!

EM waves in Conductors

- ▶ Conductors have free charges, ρ_f , and Ohmic current,

$$\mathbf{j}_{free} = \sigma \mathbf{E},$$

- ▶ Maxwell's Equations in conductors (ϵ, μ, σ)

$$\nabla \cdot \mathbf{E} = \frac{\rho_{free}}{\epsilon}, \quad \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t},$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{B} = \mu\sigma \mathbf{E} + \mu\epsilon \frac{\partial \mathbf{E}}{\partial t}$$

- ▶ Continuity Equation: Conservation of free charge

$$\nabla \cdot \mathbf{j}_{free} + \frac{\partial \rho_{free}}{\partial t} = 0,$$

$$\implies \rho(t) = e^{-(\sigma/\epsilon)t} \rho(0).$$

Charge quickly dissipates inside!

- ▶ Ultimately

$$\nabla \cdot \mathbf{E} = 0, \quad \nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t},$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{B} = \mu\sigma \mathbf{E} + \mu\epsilon \frac{\partial \mathbf{E}}{\partial t}$$

EM waves in Conductors: Skin Effect

EM waves in Conductors: Skin Effect

- ▶ Dissipative Wave equations

$$\left(\nabla^2 - \mu\epsilon \frac{\partial^2}{\partial t^2} \right) \mathbf{E}, \mathbf{B} = \mu\sigma \frac{\partial \mathbf{E}, \mathbf{B}}{\partial t}$$

EM waves in Conductors: Skin Effect

- ▶ Dissipative Wave equations

$$\left(\nabla^2 - \mu\epsilon \frac{\partial^2}{\partial t^2} \right) \mathbf{E}, \mathbf{B} = \mu\sigma \frac{\partial \mathbf{E}, \mathbf{B}}{\partial t}$$

- ▶ Plane wave solutions with amplitude declining in space

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{i(\tilde{k}z - \omega t)}, \quad \mathbf{B} = \tilde{\mathbf{B}}_0 e^{i(\tilde{k}z - \omega t)}$$
$$\tilde{k}^2 = \mu\epsilon\omega^2 + i\mu\sigma\omega.$$

EM waves in Conductors: Skin Effect

- ▶ Dissipative Wave equations

$$\left(\nabla^2 - \mu\epsilon \frac{\partial^2}{\partial t^2}\right) \mathbf{E}, \mathbf{B} = \mu\sigma \frac{\partial \mathbf{E}, \mathbf{B}}{\partial t}$$

- ▶ Plane wave solutions with amplitude declining in space

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{i(\tilde{k}z - \omega t)}, \quad \mathbf{B} = \tilde{\mathbf{B}}_0 e^{i(\tilde{k}z - \omega t)}$$
$$\tilde{k}^2 = \mu\epsilon\omega^2 + i\mu\sigma\omega.$$

- ▶ $\tilde{k} = k + i\kappa,$

$$k = \omega \sqrt{\frac{\mu\epsilon}{2}} \left(\sqrt{1 + \left(\frac{\sigma}{\epsilon\omega}\right)^2} + 1 \right)^{\frac{1}{2}} \approx \sqrt{\frac{\mu\sigma\omega}{2}},$$

$$\kappa = \omega \sqrt{\frac{\mu\epsilon}{2}} \left(\sqrt{1 + \left(\frac{\sigma}{\epsilon\omega}\right)^2} - 1 \right)^{\frac{1}{2}} \approx \sqrt{\frac{\mu\sigma\omega}{2}}$$

EM waves in Conductors: Skin Effect

- ▶ Dissipative Wave equations

$$\left(\nabla^2 - \mu\epsilon \frac{\partial^2}{\partial t^2}\right) \mathbf{E}, \mathbf{B} = \mu\sigma \frac{\partial \mathbf{E}, \mathbf{B}}{\partial t}$$

- ▶ Plane wave solutions with amplitude declining in space

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{i(\tilde{k}z - \omega t)}, \quad \mathbf{B} = \tilde{\mathbf{B}}_0 e^{i(\tilde{k}z - \omega t)}$$
$$\tilde{k}^2 = \mu\epsilon\omega^2 + i\mu\sigma\omega.$$

- ▶ $\tilde{k} = k + i\kappa,$

$$k = \omega \sqrt{\frac{\mu\epsilon}{2}} \left(\sqrt{1 + \left(\frac{\sigma}{\epsilon\omega}\right)^2} + 1 \right)^{\frac{1}{2}} \approx \sqrt{\frac{\mu\sigma\omega}{2}},$$
$$\kappa = \omega \sqrt{\frac{\mu\epsilon}{2}} \left(\sqrt{1 + \left(\frac{\sigma}{\epsilon\omega}\right)^2} - 1 \right)^{\frac{1}{2}} \approx \sqrt{\frac{\mu\sigma\omega}{2}}$$

- ▶ Amplitude falls to 1/e after a distance $d = \frac{1}{\kappa} \approx \sqrt{\frac{2}{\mu\sigma\omega}}$, (**Skin depth**)

EM waves in conductors: E and B out of phase

EM waves in conductors: \mathbf{E} and \mathbf{B} out of phase

- ▶ Gauss law and “No sinks or sources” law means: \mathbf{E} and \mathbf{B} still transverse,

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{-\kappa z} e^{i(kz - \omega t)} \hat{\mathbf{x}}$$

EM waves in conductors: \mathbf{E} and \mathbf{B} out of phase

- ▶ Gauss law and “No sinks or sources” law means: \mathbf{E} and \mathbf{B} still transverse,

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{-\kappa z} e^{i(kz - \omega t)} \hat{\mathbf{x}}$$

- ▶ \mathbf{B} has a phase difference wrt \mathbf{E}

$$\mathbf{B} = \left(\frac{\tilde{k}}{\omega} \tilde{\mathbf{E}}_0 \right) e^{-\kappa z} e^{i(kz - \omega t)} \hat{\mathbf{y}}.$$

EM waves in conductors: \mathbf{E} and \mathbf{B} out of phase

- ▶ Gauss law and “No sinks or sources” law means: \mathbf{E} and \mathbf{B} still transverse,

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{-\kappa z} e^{i(kz - \omega t)} \hat{\mathbf{x}}$$

- ▶ \mathbf{B} has a phase difference wrt \mathbf{E}

$$\mathbf{B} = \left(\frac{\tilde{k}}{\omega} \tilde{\mathbf{E}}_0 \right) e^{-\kappa z} e^{i(kz - \omega t)} \hat{\mathbf{y}}.$$

- ▶ Ratio of magnitudes

$$\frac{B_0}{E_0} = \frac{|\tilde{k}|}{\omega} = \sqrt{\epsilon\mu} \sqrt{1 + \left(\frac{\sigma}{\epsilon\omega} \right)^2} \approx \sqrt{\frac{\mu\sigma}{\omega}}$$

EM waves in conductors: \mathbf{E} and \mathbf{B} out of phase

- ▶ Gauss law and “No sinks or sources” law means: \mathbf{E} and \mathbf{B} still transverse,

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{-\kappa z} e^{i(kz - \omega t)} \hat{\mathbf{x}}$$

- ▶ \mathbf{B} has a phase difference wrt \mathbf{E}

$$\mathbf{B} = \left(\frac{\tilde{k}}{\omega} \tilde{\mathbf{E}}_0 \right) e^{-\kappa z} e^{i(kz - \omega t)} \hat{\mathbf{y}}.$$

- ▶ Ratio of magnitudes

$$\frac{B_0}{E_0} = \frac{|\tilde{k}|}{\omega} = \sqrt{\epsilon\mu} \sqrt{1 + \left(\frac{\sigma}{\epsilon\omega} \right)^2} \approx \sqrt{\frac{\mu\sigma}{\omega}}$$

- ▶ Phase lag of B wrt E

$$\varphi = \tan^{-1} \frac{\kappa}{k}.$$

EM waves in Plasma: Plasma frequency (ω_p)

EM waves in Plasma: Plasma frequency (ω_p)

- ▶ Plasma i.e. ionized gas has abundance of free charges \Rightarrow conductor.

EM waves in Plasma: Plasma frequency (ω_p)

- ▶ Plasma i.e. ionized gas has abundance of free charges \Rightarrow conductor.
- ▶ Electronic oscillations,

$$\mathbf{x}(t) = \frac{e}{m_e \omega^2} \tilde{\mathbf{E}}_0 e^{-i\omega t}$$

EM waves in Plasma: Plasma frequency (ω_p)

- ▶ Plasma i.e. ionized gas has abundance of free charges \Rightarrow conductor.
- ▶ Electronic oscillations,

$$\mathbf{x}(t) = \frac{e}{m_e \omega^2} \tilde{\mathbf{E}}_0 e^{-i\omega t}$$



$$\mathbf{j} = -Ne \dot{\mathbf{x}}, \Rightarrow \sigma = i \frac{Ne^2}{m_e \omega}$$

EM waves in Plasma: Plasma frequency (ω_p)

- ▶ Plasma i.e. ionized gas has abundance of free charges \Rightarrow conductor.
- ▶ Electronic oscillations,

$$\mathbf{x}(t) = \frac{e}{m_e \omega^2} \tilde{\mathbf{E}}_0 e^{-i\omega t}$$



$$\mathbf{j} = -Ne \dot{\mathbf{x}}, \Rightarrow \sigma = i \frac{Ne^2}{m_e \omega}$$



$$\tilde{k}^2 = \mu \epsilon \omega^2 + i \mu \sigma \omega = \mu \epsilon \omega^2 \left(1 - \frac{\omega_p^2}{\omega^2} \right), \omega_p^2 = \frac{Ne^2}{m_e \epsilon}$$

EM waves in Plasma: Plasma frequency (ω_p)

- ▶ Plasma i.e. ionized gas has abundance of free charges \Rightarrow conductor.
- ▶ Electronic oscillations,

$$\mathbf{x}(t) = \frac{e}{m_e \omega^2} \tilde{\mathbf{E}}_0 e^{-i\omega t}$$



$$\mathbf{j} = -Ne \dot{\mathbf{x}}, \Rightarrow \sigma = i \frac{Ne^2}{m_e \omega}$$



$$\tilde{k}^2 = \mu\epsilon\omega^2 + i\mu\sigma\omega = \mu\epsilon\omega^2 \left(1 - \frac{\omega_p^2}{\omega^2}\right), \omega_p^2 = \frac{Ne^2}{m_e \epsilon}$$

- ▶ For $\omega < \omega_p$, no wave propagation! Used in shortwave radio communication by bouncing off the ionosphere ($N \approx 10^{11}/m^3$, $\omega_p \sim 20$ MHz)

Reflection (& Transmission?) of waves at conductor surface

Reflection (& Transmission?) of waves at conductor surface

- Dielectric-conductor interface at $z = 0$

$$\begin{aligned}\epsilon_1 E_1^\perp - \epsilon_2 E_2^\perp &= \sigma_{free}, & \mathbf{E}_1^\parallel &= \mathbf{E}_2^\parallel, \\ B_1^\perp &= B_2^\perp, & \frac{1}{\mu_2} \mathbf{B}_1^\parallel - \frac{1}{\mu_2} \mathbf{B}_2^\parallel &= \mathbf{K}_{free} \times \hat{\mathbf{n}}\end{aligned}$$

Reflection (& Transmission?) of waves at conductor surface

- Dielectric-conductor interface at $z = 0$

$$\epsilon_1 E_1^\perp - \epsilon_2 E_2^\perp = \sigma_{free}, \quad \mathbf{E}_1^\parallel = \mathbf{E}_2^\parallel,$$
$$B_1^\perp = B_2^\perp, \quad \frac{1}{\mu_2} \mathbf{B}_1^\parallel - \frac{1}{\mu_2} \mathbf{B}_2^\parallel = \mathbf{K}_{free} \times \hat{\mathbf{n}}$$

- Say medium 1 is a dielectric and medium 2 is a conductor, then for normal incidence

$$\frac{\tilde{E}_{0R}}{\tilde{E}_{0I}} = \frac{1 - \tilde{\beta}}{1 + \tilde{\beta}}, \quad \frac{\tilde{E}_{0T}}{\tilde{E}_{0I}} = \frac{2}{1 + \tilde{\beta}}$$

where now $\tilde{\beta}$ is **complex**! To wit,

$$\tilde{\beta} = \frac{\mu_1 v_1}{\mu_2 \omega} \tilde{k}_2$$

Reflection (& Transmission?) of waves at conductor surface

- ▶ Dielectric-conductor interface at $z = 0$

$$\epsilon_1 E_1^\perp - \epsilon_2 E_2^\perp = \sigma_{free}, \quad \mathbf{E}_1^\parallel = \mathbf{E}_2^\parallel,$$
$$B_1^\perp = B_2^\perp, \quad \frac{1}{\mu_2} \mathbf{B}_1^\parallel - \frac{1}{\mu_2} \mathbf{B}_2^\parallel = \mathbf{K}_{free} \times \hat{\mathbf{n}}$$

- ▶ Say medium 1 is a dielectric and medium 2 is a conductor, then for normal incidence

$$\frac{\tilde{E}_{0R}}{\tilde{E}_{0I}} = \frac{1 - \tilde{\beta}}{1 + \tilde{\beta}}, \quad \frac{\tilde{E}_{0T}}{\tilde{E}_{0I}} = \frac{2}{1 + \tilde{\beta}}$$

where now $\tilde{\beta}$ is **complex**! To wit,

$$\tilde{\beta} = \frac{\mu_1 v_1}{\mu_2 \omega} \tilde{k}_2$$

- ▶ Perfect conductor, $\sigma \rightarrow \infty$

$$\frac{\tilde{E}_{0R}}{\tilde{E}_{0I}} = -1, \quad \frac{\tilde{E}_{0T}}{\tilde{E}_{0I}} = 0$$

Conductors (metals) are excellent reflectors!

Theory of Dispersion

Theory of Dispersion

- ▶ Dispersion of waves = Frequency components travel with different phase velocity, $v \equiv \frac{\omega}{k} = v(k)$

Theory of Dispersion

- ▶ Dispersion of waves = Frequency components travel with different phase velocity, $v \equiv \frac{\omega}{k} = v(k)$
- ▶ The envelope (amplitude) travels with the speed $v_g = \frac{d\omega}{dk}$ (Group velocity)

Theory of Dispersion

- ▶ Dispersion of waves = Frequency components travel with different phase velocity, $v \equiv \frac{\omega}{k} = v(k)$
- ▶ The envelope (amplitude) travels with the speed $v_g = \frac{d\omega}{dk}$ (Group velocity)
- ▶ What causes dispersion: Theoretical picture of dispersion is a model based forced oscillation of electrons inside molecules of a substance when EM waves (light) propagate thru dielectrics

$$\begin{aligned} m\ddot{x} &= F_{\text{restoring}} + F_{\text{damping}} + F_{\text{EM-Wave}} \\ &= -m\omega_0^2 x - m\gamma \dot{x} + q (E_0 e^{i\omega t}) \end{aligned}$$

Theory of Dispersion

- ▶ Dispersion of waves = Frequency components travel with different phase velocity, $v \equiv \frac{\omega}{k} = v(k)$
- ▶ The envelope (amplitude) travels with the speed $v_g = \frac{d\omega}{dk}$ (Group velocity)
- ▶ What causes dispersion: Theoretical picture of dispersion is a model based forced oscillation of electrons inside molecules of a substance when EM waves (light) propagate thru dielectrics

$$\begin{aligned} m\ddot{x} &= F_{\text{restoring}} + F_{\text{damping}} + F_{\text{EM-Wave}} \\ &= -m\omega_0^2 x - m\gamma \dot{x} + q (E_0 e^{i\omega t}) \end{aligned}$$

- ▶ Suffices to consider 1d motion (radial motion). Driven (radial) oscillations given by (complex) position amplitude,

$$\tilde{x}_0 = \frac{q/m}{\omega_0^2 - \omega^2 - i\gamma\omega} E_0 e^{i\omega t}$$

- ▶ Displacement of electrons from the center of charge leads to creation of dipole moment,

$$\tilde{p} = q \tilde{x}_0$$

Theory of Dispersion

Theory of Dispersion

- ▶ Total induced polarization of the medium = sum over dipole moments of all electrons in a molecule and then multiply by # of molecules per unit volume (N),

$$\tilde{P}(\omega) = N \frac{q^2}{m} \tilde{E} \sum_j \frac{n_j}{\omega_j^2 - \omega^2 - i\gamma_j\omega},$$

n_j is the number of electrons in a molecule with natural frequency ω_j , j is being summed over all the different natural frequencies of the molecule.

Theory of Dispersion

- ▶ Total induced polarization of the medium = sum over dipole moments of all electrons in a molecule and then multiply by # of molecules per unit volume (N),

$$\tilde{P}(\omega) = N \frac{q^2}{m} \tilde{E} \sum_j \frac{n_j}{\omega_j^2 - \omega^2 - i\gamma_j\omega},$$

n_j is the number of electrons in a molecule with natural frequency ω_j , j is being summed over all the different natural frequencies of the molecule.

- ▶ Permittivity is then a function of frequency, ω

$$\frac{\tilde{\epsilon}(\omega)}{\epsilon_0} \equiv 1 + \frac{\tilde{P}}{\epsilon_0 \tilde{E}} = 1 + \frac{Nq^2}{m\epsilon_0} \sum_j \frac{n_j}{\omega_j^2 - \omega^2 - i\gamma_j\omega}$$

Theory of Dispersion

- ▶ Total induced polarization of the medium = sum over dipole moments of all electrons in a molecule and then multiply by # of molecules per unit volume (N),

$$\tilde{P}(\omega) = N \frac{q^2}{m} \tilde{E} \sum_j \frac{n_j}{\omega_j^2 - \omega^2 - i\gamma_j \omega},$$

n_j is the number of electrons in a molecule with natural frequency ω_j , j is being summed over all the different natural frequencies of the molecule.

- ▶ Permittivity is then a function of frequency, ω

$$\frac{\tilde{\epsilon}(\omega)}{\epsilon_0} \equiv 1 + \frac{\tilde{P}}{\epsilon_0 \tilde{E}} = 1 + \frac{Nq^2}{m\epsilon_0} \sum_j \frac{n_j}{\omega_j^2 - \omega^2 - i\gamma_j \omega}$$

- ▶ Refractive Index as a result varies with ω

$$n = \sqrt{\operatorname{Re} \left(\frac{\tilde{\epsilon}(\omega)}{\epsilon_0} \right)} = 1 + \frac{Nq^2}{2m\epsilon_0} \sum_j \frac{f_j (\omega_j^2 - \omega^2)}{(\omega_j^2 - \omega^2)^2 + \gamma_j^2 \omega^2}.$$

Theory of Dispersion

Theory of Dispersion

- ▶ Normal dispersion (refractive index increases with frequency): For frequencies of the EM wave much less than the natural frequencies of the molecular electrons, $\omega \ll \omega_j$, n increases with ω

Theory of Dispersion

- ▶ Normal dispersion (refractive index increases with frequency): For frequencies of the EM wave much less than the natural frequencies of the molecular electrons, $\omega \ll \omega_j$, n increases with ω
- ▶ Anomalous dispersion (refractive index decreases with frequency): When the frequency of the propagating EM wave, $\omega \sim \omega_j$ for some j , index suddenly plunges

$$n \propto 1 + \omega^{-2}$$

!

Theory of Dispersion

- ▶ Normal dispersion (refractive index increases with frequency): For frequencies of the EM wave much less than the natural frequencies of the molecular electrons, $\omega \ll \omega_j$, n increases with ω
- ▶ Anomalous dispersion (refractive index decreases with frequency): When the frequency of the propagating EM wave, $\omega \sim \omega_j$ for some j , index suddenly plunges

$$n \propto 1 + \omega^{-2}$$

!

- ▶ For normal dispersion, i.e. far from resonant we can neglect damping ($\gamma_j \rightarrow 0$), as a result,

$$n = 1 + A(1 + B\omega^2) = 1 + A\left(1 + \frac{B}{\lambda^2}\right)$$

Celebrated result due to Cauchy (Cauchy dispersion formula, verified in the lab)!