

EP 1027: Maxwell's Equations and Electromagnetic waves

Instructor: Shubho Roy¹
(Dept. of Physics)

Lecture 1

March 26, 2019

¹ Office: C 313 D

Email: sroy@iith.ac.in

Office hours By email appointment or walk in

Rough Plan

Rough Plan

- ▶ Vector Calculus: Notations and algebra of Cartesian Vectors, Vector Differential operators, Integration of vectors, Curvilinear coordinates

Rough Plan

- ▶ Vector Calculus: Notations and algebra of Cartesian Vectors, Vector Differential operators, Integration of vectors, Curvilinear coordinates
- ▶ Static charges: Electric fields in vacuum and matter (conductors and dielectrics)
- ▶ Steady Currents: Magnetic fields in vacuum and matter (paramagnetics)

Rough Plan

- ▶ Vector Calculus: Notations and algebra of Cartesian Vectors, Vector Differential operators, Integration of vectors, Curvilinear coordinates
- ▶ Static charges: Electric fields in vacuum and matter (conductors and dielectrics)
- ▶ Steady Currents: Magnetic fields in vacuum and matter (paramagnetics)
- ▶ Electrodynamics (general case): Maxwell's equations

Rough Plan

- ▶ Vector Calculus: Notations and algebra of Cartesian Vectors, Vector Differential operators, Integration of vectors, Curvilinear coordinates
- ▶ Static charges: Electric fields in vacuum and matter (conductors and dielectrics)
- ▶ Steady Currents: Magnetic fields in vacuum and matter (paramagnetics)
- ▶ Electrodynamics (general case): Maxwell's equations
- ▶ Electromagnetic (EM) Waves in vacuum, in media (reflection, refraction, absorption, dispersion), Guided Waves .

References

References

- ▶ Griffiths, D.J. **Introduction to Electrodynamics.**
- ▶ Purcell, E. **Electricity and Magnetism**
- ▶ **Feynman lectures on Physics**, Vol. 2
- ▶ Reitz, Milford and Christy: **Foundations of Electromagnetic Theory**
- ▶ Schaum's Outline of Electromagnetics (Edminister, J)
- ▶ ** Advanced books: Jackson, Panofsky & Phillips, Schwartz, Schwinger

Omissions?

Omissions?

- ▶ Boundary Value problems: Solving Laplace or Poisson's equations (Method of Images, Separation of variables method, Green's function method, Computational approaches)

Omissions?

- ▶ Boundary Value problems: Solving Laplace or Poisson's equations (Method of Images, Separation of variables method, Green's function method, Computational approaches)
- ▶ Multipole Expansions

Omissions?

- ▶ Boundary Value problems: Solving Laplace or Poisson's equations (Method of Images, Separation of variables method, Green's function method, Computational approaches)
- ▶ Multipole Expansions
- ▶ DC and AC Circuits

Omissions?

- ▶ Boundary Value problems: Solving Laplace or Poisson's equations (Method of Images, Separation of variables method, Green's function method, Computational approaches)
- ▶ Multipole Expansions
- ▶ DC and AC Circuits
- ▶ Special Relativistic formulation of Electrodynamics: Spacetime tensor notation

Omissions?

- ▶ Boundary Value problems: Solving Laplace or Poisson's equations (Method of Images, Separation of variables method, Green's function method, Computational approaches)
- ▶ Multipole Expansions
- ▶ DC and AC Circuits
- ▶ Special Relativistic formulation of Electrodynamics: Spacetime tensor notation
- ▶ Radiation from accelerating charges: Cerenkov, Synchrotron, ... only if we have time

Evaluation

Evaluation

- ▶ Assessment: Final (100 points, 3 hrs.) + 3(4) Quizzes (10 points, 10 min.)

Evaluation

- ▶ Assessment: Final (100 points, 3 hrs.) + 3(4) Quizzes (10 points, 10 min.)
- ▶ Grading is absolute ($A > 80$)

Evaluation

- ▶ Assessment: Final (100 points, 3 hrs.) + 3(4) Quizzes (10 points, 10 min.)
- ▶ Grading is absolute ($A > 80$)
- ▶ Tentative final date (May 4, 10 AM - 1 PM), Review of grades (May 7)

Review of Cartesian Vectors: Notation and Terminology

Review of Cartesian Vectors: Notation and Terminology

- ▶ What is a vector?

Review of Cartesian Vectors: Notation and Terminology

- ▶ What is a vector?
- ▶ Quantity w/ both magnitude and a direction (*directed line segment*). e.g. Position vector, \mathbf{x} .

Review of Cartesian Vectors: Notation and Terminology

- ▶ What is a vector?
- ▶ Quantity w/ both magnitude and a direction (*directed line segment*). e.g. Position vector, \mathbf{x} .
- ▶ behaves like position vector under vector addition and multiplication by a scalar, **and under rotation of coordinate axes**.

Review of Cartesian Vectors: Notation and Terminology

- ▶ What is a vector?
- ▶ Quantity w/ both magnitude and a direction (*directed line segment*). e.g. Position vector, \mathbf{x} .
- ▶ behaves like position vector under vector addition and multiplication by a scalar, **and under rotation of coordinate axes**.
- ▶ Position vector: 3-component object (triplet) giving the location from a origin of Cartesian coordinate system: row

vector (x, y, z) or a column vector $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$ or $\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$.

Review of Cartesian Vectors: Notation and Terminology

- ▶ What is a vector?
- ▶ Quantity w/ both magnitude and a direction (*directed line segment*). e.g. Position vector, \mathbf{x} .
- ▶ behaves like position vector under vector addition and multiplication by a scalar, **and under rotation of coordinate axes**.
- ▶ Position vector: 3-component object (triplet) giving the location from a origin of Cartesian coordinate system: row vector (x, y, z) or a column vector $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$ or $\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$.
- ▶ Notation: Call the components of the vector: x_k , k can take values from 1 to 3. Whole vector is denoted in boldface, \mathbf{x} or, with an arrow overhead, \vec{x} .

Cartesian Vectors: Notation and Terminology

Cartesian Vectors: Notation and Terminology

- ▶ Rule for addition of 2 vectors: Add the respective components

$$\mathbf{x} + \mathbf{y} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \equiv \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{pmatrix}$$

Cartesian Vectors: Notation and Terminology

- ▶ Rule for addition of 2 vectors: Add the respective components

$$\mathbf{x} + \mathbf{y} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \equiv \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{pmatrix}$$

- ▶ Rule for multiplication with a number, λ : Multiply each component by λ

$$\lambda \mathbf{x} = \lambda \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \equiv \begin{pmatrix} \lambda x_1 \\ \lambda x_2 \\ \lambda x_3 \end{pmatrix} .$$

Cartesian Vectors: Notation and Terminology

- ▶ Rule for addition of 2 vectors: Add the respective components

$$\mathbf{x} + \mathbf{y} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} + \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix} \equiv \begin{pmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{pmatrix}$$

- ▶ Rule for multiplication with a number, λ : Multiply each component by λ

$$\lambda \mathbf{x} = \lambda \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} \equiv \begin{pmatrix} \lambda x_1 \\ \lambda x_2 \\ \lambda x_3 \end{pmatrix}.$$

- ▶ Once we choose the vector to be a column, then to denote the row vector, we will use the transpose

$$\mathbf{x}^T = (x_1, x_2, x_3)$$

Cartesian Vectors: Notation and Terminology

Cartesian Vectors: Notation and Terminology

- Notation: Basis Vectors, $\hat{\mathbf{e}}_k$

$$\begin{aligned}\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} &= x_1 \underbrace{\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}}_{\hat{\mathbf{e}}_1} + x_2 \underbrace{\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}}_{\hat{\mathbf{e}}_2} + x_3 \underbrace{\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}}_{\hat{\mathbf{e}}_3} \\ &= x_1 \hat{\mathbf{e}}_1 + x_2 \hat{\mathbf{e}}_2 + x_3 \hat{\mathbf{e}}_3 \\ &= \sum_{k=1}^3 x_k \hat{\mathbf{e}}_k.\end{aligned}$$

Cartesian Vectors: Notation and Terminology

- Notation: Basis Vectors, $\hat{\mathbf{e}}_k$

$$\begin{aligned}\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} &= x_1 \underbrace{\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}}_{\hat{\mathbf{e}}_1} + x_2 \underbrace{\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}}_{\hat{\mathbf{e}}_2} + x_3 \underbrace{\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}}_{\hat{\mathbf{e}}_3} \\ &= x_1 \hat{\mathbf{e}}_1 + x_2 \hat{\mathbf{e}}_2 + x_3 \hat{\mathbf{e}}_3 \\ &= \sum_{k=1}^3 x_k \hat{\mathbf{e}}_k.\end{aligned}$$

- “Dot product/ inner product/ scalar product” of \mathbf{x} and \mathbf{y} :

$$\mathbf{x} \cdot \mathbf{y} \equiv \mathbf{x}^T \mathbf{y} = x_1 y_1 + x_2 y_2 + x_3 y_3 = \sum_{k=1}^3 x_k y_k.$$

Cartesian Vectors: Notation and Terminology

- ▶ **Einstein summation convention:** Clean up notation by eradicating summation sign $\sum_{k=1}^3$

Cartesian Vectors: Notation and Terminology

- ▶ **Einstein summation convention:** Clean up notation by eradicating summation sign $\sum_{k=1}^3$
- ▶ Rule 1) Repeating an index \Rightarrow it is summed over all values 1, 2, 3. E.g.

$$\mathbf{x} = \sum_{i=1}^3 x_i \hat{\mathbf{e}}_i = x_i \hat{\mathbf{e}}_i, \quad \mathbf{x} \cdot \mathbf{y} = \sum_{i=1}^3 x_i y_i = x_i y_i.$$

Cartesian Vectors: Notation and Terminology

- ▶ **Einstein summation convention:** Clean up notation by eradicating summation sign $\sum_{k=1}^3$
- ▶ Rule 1) Repeating an index \Rightarrow it is summed over all values 1, 2, 3. E.g.

$$\mathbf{x} = \sum_{i=1}^3 x_i \hat{\mathbf{e}}_i = x_i \hat{\mathbf{e}}_i, \quad \mathbf{x} \cdot \mathbf{y} = \sum_{i=1}^3 x_i y_i = x_i y_i.$$

- ▶ Rule 2) Non-repeated indices are “free” to take any values, should not be summed,

Cartesian Vectors: Notation and Terminology

- ▶ **Einstein summation convention:** Clean up notation by eradicating summation sign $\sum_{k=1}^3$
- ▶ Rule 1) Repeating an index \Rightarrow it is summed over all values 1, 2, 3. E.g.

$$\mathbf{x} = \sum_{i=1}^3 x_i \hat{\mathbf{e}}_i = x_i \hat{\mathbf{e}}_i, \quad \mathbf{x} \cdot \mathbf{y} = \sum_{i=1}^3 x_i y_i = x_i y_i.$$

- ▶ Rule 2) Non-repeated indices are “free” to take any values, should not be summed,
- ▶ Subrule 3) Repeated index cannot appear more than twice, $x_k x_k y_k$ is illegal/inconsistent

Cartesian Vectors: Notation and Terminology

- ▶ **Einstein summation convention:** Clean up notation by eradicating summation sign $\sum_{k=1}^3$
- ▶ Rule 1) Repeating an index \Rightarrow it is summed over all values 1, 2, 3. E.g.

$$\mathbf{x} = \sum_{i=1}^3 x_i \hat{\mathbf{e}}_i = x_i \hat{\mathbf{e}}_i, \quad \mathbf{x} \cdot \mathbf{y} = \sum_{i=1}^3 x_i y_i = x_i y_i.$$

- ▶ Rule 2) Non-repeated indices are “free” to take any values, should not be summed,
- ▶ Subrule 3) Repeated index cannot appear more than twice, $x_k x_k y_k$ is illegal/inconsistent
- ▶ Subrule 4) A repeated index can be relabeled anytime (dummy). E.g. $\mathbf{x} \cdot \mathbf{y} = x_i y_i = x_l y_l = x_m y_m$.

Cartesian Vectors: Notation and Terminology

Cartesian Vectors: Notation and Terminology

- ▶ **Norm** (or size) of a vector: $||\mathbf{x}|| \equiv \sqrt{\mathbf{x} \cdot \mathbf{x}} = \sqrt{x_k x_k}$

Cartesian Vectors: Notation and Terminology

- ▶ **Norm** (or size) of a vector: $\|\mathbf{x}\| \equiv \sqrt{\mathbf{x} \cdot \mathbf{x}} = \sqrt{x_k x_k}$
- ▶ **Outer product**/ **Tensor product** of \mathbf{x} and \mathbf{y} ,

$$T_{lk} = x_l y_k, \quad \overleftrightarrow{T} = \mathbf{T} = \begin{pmatrix} x_1 y_1 & x_1 y_2 & x_1 y_3 \\ x_2 y_1 & \dots & \\ x_3 y_1 & \dots & \end{pmatrix}$$

Cartesian Vectors: Notation and Terminology

- ▶ **Norm** (or size) of a vector: $\|\mathbf{x}\| \equiv \sqrt{\mathbf{x} \cdot \mathbf{x}} = \sqrt{x_k x_k}$
- ▶ **Outer product/ Tensor product** of \mathbf{x} and \mathbf{y} ,

$$T_{lk} = x_l y_k, \quad \overleftrightarrow{T} = \mathbf{T} = \begin{pmatrix} x_1 y_1 & x_1 y_2 & x_1 y_3 \\ x_2 y_1 & \dots & \\ x_3 y_1 & \dots & \end{pmatrix}$$

- ▶ **Vector product/Cross product**

$$\mathbf{x} \times \mathbf{y} \equiv (\epsilon_{ijk} x_i y_j) \hat{\mathbf{e}}_k,$$

where

$$\epsilon_{123} = \epsilon_{231} = \epsilon_{312} = 1,$$

$$\epsilon_{132} = \epsilon_{213} = \epsilon_{321} = -1,$$

$$\epsilon_{112} = \epsilon_{121} = \epsilon_{223} \dots = 0.$$

ϵ_{ijk} = “Levi Civita Symbol or Completely antisymmetric symbol”

Cartesian Vectors: Notation and Terminology

Cartesian Vectors: Notation and Terminology

- ▶ e.g. Let's compute the 1st component of $\mathbf{x} \times \mathbf{y}$:

$$(\mathbf{x} \times \mathbf{y})_1 = \epsilon_{ij1} x_i y_j,$$

i and j are both repeated \implies we have to sum over all values of i and j . Sum over i first,

$$\begin{aligned} (\mathbf{x} \times \mathbf{y})_1 = \epsilon_{ij1} x_i y_j &= \underbrace{\epsilon_{1j1}}_{=0} x_1 y_j + \epsilon_{2j1} x_2 y_j + \epsilon_{3j1} x_3 y_j. \\ &= \epsilon_{2j1} x_2 y_j + \epsilon_{3j1} x_3 y_j. \end{aligned}$$

Cartesian Vectors: Notation and Terminology

- ▶ e.g. Let's compute the 1st component of $\mathbf{x} \times \mathbf{y}$:

$$(\mathbf{x} \times \mathbf{y})_1 = \epsilon_{ij1} x_i y_j,$$

i and j are both repeated \implies we have to sum over all values of i and j . Sum over i first,

$$\begin{aligned} (\mathbf{x} \times \mathbf{y})_1 = \epsilon_{ij1} x_i y_j &= \underbrace{\epsilon_{1j1}}_{=0} x_1 y_j + \epsilon_{2j1} x_2 y_j + \epsilon_{3j1} x_3 y_j. \\ &= \epsilon_{2j1} x_2 y_j + \epsilon_{3j1} x_3 y_j. \end{aligned}$$

- ▶ Next, sum over index j . The first term,

$$\epsilon_{2j1} x_2 y_j = \underbrace{\epsilon_{211}}_{=0} x_2 y_1 + \underbrace{\epsilon_{221}}_{=0} x_2 y_2 + \underbrace{\epsilon_{231}}_{=1} x_2 y_3 = x_2 y_3.$$

Cartesian Vectors: Notation and Terminology

- ▶ e.g. Let's compute the 1st component of $\mathbf{x} \times \mathbf{y}$:

$$(\mathbf{x} \times \mathbf{y})_1 = \epsilon_{ij1} x_i y_j,$$

i and j are both repeated \implies we have to sum over all values of i and j . Sum over i first,

$$\begin{aligned} (\mathbf{x} \times \mathbf{y})_1 = \epsilon_{ij1} x_i y_j &= \underbrace{\epsilon_{1j1}}_{=0} x_1 y_j + \epsilon_{2j1} x_2 y_j + \epsilon_{3j1} x_3 y_j. \\ &= \epsilon_{2j1} x_2 y_j + \epsilon_{3j1} x_3 y_j. \end{aligned}$$

- ▶ Next, sum over index j . The first term,

$$\epsilon_{2j1} x_2 y_j = \underbrace{\epsilon_{211}}_{=0} x_2 y_1 + \underbrace{\epsilon_{221}}_{=0} x_2 y_2 + \underbrace{\epsilon_{231}}_{=1} x_2 y_3 = x_2 y_3.$$

- ▶ Similarly, can show the second term: $\epsilon_{3j1} x_3 y_j = -x_3 y_2$.

Cartesian Vectors: Notation and Terminology

- ▶ e.g. Let's compute the 1st component of $\mathbf{x} \times \mathbf{y}$:

$$(\mathbf{x} \times \mathbf{y})_1 = \epsilon_{ij1} x_i y_j,$$

i and j are both repeated \implies we have to sum over all values of i and j . Sum over i first,

$$\begin{aligned} (\mathbf{x} \times \mathbf{y})_1 = \epsilon_{ij1} x_i y_j &= \underbrace{\epsilon_{1j1}}_{=0} x_1 y_j + \epsilon_{2j1} x_2 y_j + \epsilon_{3j1} x_3 y_j. \\ &= \epsilon_{2j1} x_2 y_j + \epsilon_{3j1} x_3 y_j. \end{aligned}$$

- ▶ Next, sum over index j . The first term,

$$\epsilon_{2j1} x_2 y_j = \underbrace{\epsilon_{211}}_{=0} x_2 y_1 + \underbrace{\epsilon_{221}}_{=0} x_2 y_2 + \underbrace{\epsilon_{231}}_{=1} x_2 y_3 = x_2 y_3.$$

- ▶ Similarly, can show the second term: $\epsilon_{3j1} x_3 y_j = -x_3 y_2$.
- ▶ Thus we have derived: $(\mathbf{x} \times \mathbf{y})_1 = x_2 y_3 - x_3 y_2$.

Cartesian Vectors: Notation and Terminology

Cartesian Vectors: Notation and Terminology

- ▶ Vector transformation rule under rotation of coordinate axes: Implemented through Matrix operations, \mathbf{O}

$$\mathbf{x}' = \mathbf{O}\mathbf{x},$$
$$x'_i = O_{ij} x_j = \begin{pmatrix} O_{11} & O_{12} & O_{13} \\ O_{21} & & \\ O_{22} & & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

Cartesian Vectors: Notation and Terminology

- ▶ Vector transformation rule under rotation of coordinate axes: Implemented through Matrix operations, \mathbf{O}

$$\mathbf{x}' = \mathbf{O}\mathbf{x},$$
$$x'_i = O_{ij} x_j = \begin{pmatrix} O_{11} & O_{12} & O_{13} \\ O_{21} & & \\ O_{22} & & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

- ▶ Rotation matrices must be Orthogonal matrices. Why?

Cartesian Vectors: Notation and Terminology

- ▶ Vector transformation rule under rotation of coordinate axes: Implemented through Matrix operations, \mathbf{O}

$$\mathbf{x}' = \mathbf{O}\mathbf{x},$$
$$x'_i = O_{ij} x_j = \begin{pmatrix} O_{11} & O_{12} & O_{13} \\ O_{21} & & \\ O_{22} & & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

- ▶ Rotation matrices must be Orthogonal matrices. Why?
- ▶ Under Rotation, length/ Norm of vector remains same, i.e.

$$\mathbf{x}' \cdot \mathbf{x}' = \mathbf{x} \cdot \mathbf{x},$$

Cartesian Vectors: Notation and Terminology

- ▶ Vector transformation rule under rotation of coordinate axes: Implemented through Matrix operations, \mathbf{O}

$$\mathbf{x}' = \mathbf{O}\mathbf{x},$$
$$x'_i = O_{ij} x_j = \begin{pmatrix} O_{11} & O_{12} & O_{13} \\ O_{21} & & \\ O_{22} & & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

- ▶ Rotation matrices must be Orthogonal matrices. Why?
- ▶ Under Rotation, length/ Norm of vector remains same, i.e.

$$\mathbf{x}' \cdot \mathbf{x}' = \mathbf{x} \cdot \mathbf{x},$$

- ▶ This means

$$\mathbf{x}'^T \mathbf{x}' = (\mathbf{x}^T \mathbf{O}^T) \mathbf{O} \mathbf{x} = \mathbf{x}^T \mathbf{x},$$
$$\implies \mathbf{O}^T \mathbf{O} = \mathbf{I}$$

Cartesian Vectors: Notation and Terminology

- ▶ Vector transformation rule under rotation of coordinate axes: Implemented through Matrix operations, \mathbf{O}

$$\mathbf{x}' = \mathbf{O}\mathbf{x},$$
$$x'_i = O_{ij} x_j = \begin{pmatrix} O_{11} & O_{12} & O_{13} \\ O_{21} & & \\ O_{22} & & \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

- ▶ Rotation matrices must be Orthogonal matrices. Why?
- ▶ Under Rotation, length/ Norm of vector remains same, i.e.

$$\mathbf{x}' \cdot \mathbf{x}' = \mathbf{x} \cdot \mathbf{x},$$

- ▶ This means

$$\mathbf{x}'^T \mathbf{x}' = (\mathbf{x}^T \mathbf{O}^T) \mathbf{O} \mathbf{x} = \mathbf{x}^T \mathbf{x},$$
$$\implies \mathbf{O}^T \mathbf{O} = \mathbf{I}$$

- ▶ Component notation: $O_{ij} O_{ik} = \delta_{jk}$.

Summary of Lecture 1

- ▶ Indices introduced for components of a vector.

Summary of Lecture 1

- ▶ Indices introduced for components of a vector.
- ▶ Summation convention for indexed objects to clean up notation, no more Σ signs, only dummy indices or free indices.

Summary of Lecture 1

- ▶ Indices introduced for components of a vector.
- ▶ Summation convention for indexed objects to clean up notation, no more Σ signs, only dummy indices or free indices.
- ▶ Summation convention applied to simplify dot products and cross products.

Summary of Lecture 1

- ▶ Indices introduced for components of a vector.
- ▶ Summation convention for indexed objects to clean up notation, no more Σ signs, only dummy indices or free indices.
- ▶ Summation convention applied to simplify dot products and cross products.
- ▶ Two new symbols (indexed objects) introduced, namely, the Levi Civita (antisymmetric) ϵ_{ijk} and the Kronecker (symmetric) δ_{ij} .

Summary of Lecture 1

- ▶ Indices introduced for components of a vector.
- ▶ Summation convention for indexed objects to clean up notation, no more Σ signs, only dummy indices or free indices.
- ▶ Summation convention applied to simplify dot products and cross products.
- ▶ Two new symbols (indexed objects) introduced, namely, the Levi Civita (antisymmetric) ϵ_{ijk} and the Kronecker (symmetric) δ_{ij} .
- ▶ Effect of Rotation of coordinate axis accomplished through orthogonal transformations on the vector.