

EP 1027: Maxwell's Equations and Electromagnetic Waves

Instructor: Shubho Roy¹
(Dept. of Physics)

Lecture 9 & 10

April 12, 2019

¹Email: sroy@iith.ac.in, Office: C 313 D, Office hrs: Walk in/ By Email Appointment

Agenda

Agenda

- ▶ Wave equation basics: d'Alembert's solution, Bernoulli's solution, superposition

Agenda

- ▶ Wave equation basics: d'Alembert's solution, Bernoulli's solution, superposition
- ▶ EM waves in vacuum

Agenda

- ▶ Wave equation basics: d'Alembert's solution, Bernoulli's solution, superposition
- ▶ EM waves in vacuum
- ▶ EM waves in matter

Agenda

- ▶ Wave equation basics: d'Alembert's solution, Bernoulli's solution, superposition
- ▶ EM waves in vacuum
- ▶ EM waves in matter
- ▶ Laws of reflection and refraction for waves

Agenda

- ▶ Wave equation basics: d'Alembert's solution, Bernoulli's solution, superposition
- ▶ EM waves in vacuum
- ▶ EM waves in matter
- ▶ Laws of reflection and refraction for waves
- ▶ Fresnel Equations

References/Readings

References/Readings

- ▶ Griffiths, D.J., **Introduction to Electrodynamics, Ch. 9**

Wave mechanics : Basics

Wave mechanics : Basics

- ▶ Generic Wave equation (in 1D),

$$\left(\frac{\partial^2}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) f(x, t) = 0,$$

c is the wave-velocity

Wave mechanics : Basics

- ▶ Generic Wave equation (in 1D),

$$\left(\frac{\partial^2}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) f(x, t) = 0,$$

c is the wave-velocity

- ▶ d'Alembert's solution

$$f(x, t) = g(x - ct) + h(x + ct).$$

Wave mechanics : Basics

- ▶ Generic Wave equation (in 1D),

$$\left(\frac{\partial^2}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) f(x, t) = 0,$$

c is the wave-velocity

- ▶ d'Alembert's solution

$$f(x, t) = g(x - ct) + h(x + ct).$$

- ▶ Linear Equation: So, waves superpose to give a new solution,

$$f(x, t) = \sum_i c_i f(x, t).$$

Wave mechanics : Basics

- ▶ Generic Wave equation (in 1D),

$$\left(\frac{\partial^2}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) f(x, t) = 0,$$

c is the wave-velocity

- ▶ d'Alembert's solution

$$f(x, t) = g(x - ct) + h(x + ct).$$

- ▶ Linear Equation: So, waves superpose to give a new solution,

$$f(x, t) = \sum_i c_i f(x, t).$$

- ▶ Bernoulli's solution: Sinusoidal

$$f(x, t) = A \cos(kx - \omega t + \Delta_1) + B \cos(kx + \omega t + \Delta_2), \omega = kc.$$

Wave mechanics : Basics

Wave mechanics : Basics

- Complex form,

$$\tilde{f}(x, t) = \tilde{A} \exp(i k x - i \omega t), \quad \tilde{A} = A e^{i \Delta}$$

Wave mechanics : Basics

- Complex form,

$$\tilde{f}(x, t) = \tilde{A} \exp(i k x - i \omega t), \quad \tilde{A} = A e^{i \Delta}$$

- Expansion in Sinusoidal basis

$$\tilde{f}(x, t) = \int_{-\infty}^{\infty} dk \tilde{A}(k) e^{i(kx - \omega t)}.$$

Wave mechanics : Basics

- ▶ Complex form,

$$\tilde{f}(x, t) = \tilde{A} \exp(i kx - i\omega t), \quad \tilde{A} = A e^{i\Delta}$$

- ▶ Expansion in Sinusoidal basis

$$\tilde{f}(x, t) = \int_{-\infty}^{\infty} dk \tilde{A}(k) e^{i(kx - \omega t)}.$$

- ▶ Boundary conditions: Frequency of (Incident, Reflected, transmitted) must remain same, speed and wavelength can change,

Wave mechanics : Basics

- ▶ Complex form,

$$\tilde{f}(x, t) = \tilde{A} \exp(i k x - i \omega t), \quad \tilde{A} = A e^{i \Delta}$$

- ▶ Expansion in Sinusoidal basis

$$\tilde{f}(x, t) = \int_{-\infty}^{\infty} dk \tilde{A}(k) e^{i(kx - \omega t)}.$$

- ▶ Boundary conditions: Frequency of (Incident, Reflected, transmitted) must remain same, speed and wavelength can change,

Maxwell's Equations: EM waves in vacuum

Maxwell's Equations: EM waves in vacuum

- In the absence of sources,

$$\nabla \cdot \mathbf{E} = \nabla \cdot \mathbf{B} = 0,$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}, \nabla \times \mathbf{B} = \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t}$$

Maxwell's Equations: EM waves in vacuum

- In the absence of sources,

$$\nabla \cdot \mathbf{E} = \nabla \cdot \mathbf{B} = 0,$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}, \nabla \times \mathbf{B} = \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t}$$

- 3D Wave Equations for \mathbf{E}, \mathbf{B}

$$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \mathbf{E} = 0,$$

$$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \mathbf{B} = 0.$$

Maxwell's Equations: EM waves in vacuum

- In the absence of sources,

$$\nabla \cdot \mathbf{E} = \nabla \cdot \mathbf{B} = 0,$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}, \nabla \times \mathbf{B} = \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t}$$

- 3D Wave Equations for \mathbf{E}, \mathbf{B}

$$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \mathbf{E} = 0,$$

$$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) \mathbf{B} = 0.$$

- c is the speed of light in vacuum, light is an EM wave!

EM waves in vacuum: Plane Waves

EM waves in vacuum: Plane Waves

- ▶ Wave front is planar & perpendicular to direction of propagation

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}, \mathbf{B} = \tilde{\mathbf{B}}_0 e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)},$$

\mathbf{k} is the wave vector, normal to wave fronts.

EM waves in vacuum: Plane Waves

- ▶ Wave front is planar & perpendicular to direction of propagation

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}, \mathbf{B} = \tilde{\mathbf{B}}_0 e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)},$$

\mathbf{k} is the wave vector, normal to wave fronts.

- ▶ Gauss law ($\nabla \cdot \mathbf{E} = \nabla \cdot \mathbf{B} = 0$) implies,

$$\mathbf{k} \cdot \mathbf{E} = \mathbf{k} \cdot \mathbf{B} = 0,$$

EM waves are transverse!

EM waves in vacuum: Plane Waves

- ▶ Wave front is planar & perpendicular to direction of propagation

$$\mathbf{E} = \tilde{\mathbf{E}}_0 e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}, \mathbf{B} = \tilde{\mathbf{B}}_0 e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)},$$

\mathbf{k} is the wave vector, normal to wave fronts.

- ▶ Gauss law ($\nabla \cdot \mathbf{E} = \nabla \cdot \mathbf{B} = 0$) implies,

$$\mathbf{k} \cdot \mathbf{E} = \mathbf{k} \cdot \mathbf{B} = 0,$$

EM waves are transverse!

- ▶ Faraday law ($\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$) implies,

$$\mathbf{B} = \hat{\mathbf{k}} \times \frac{\mathbf{E}}{c}.$$

EM waves in vacuum: Energy and Momentum

EM waves in vacuum: Energy and Momentum

► Using, $\mathbf{B} = \hat{\mathbf{k}} \times \frac{\mathbf{E}}{c}$, can show,

$$\frac{1}{2\mu_0} \mathbf{B}^2 = \frac{\epsilon_0}{2} \mathbf{E}^2, u_{\mathbf{E}} = u_{\mathbf{B}},$$

$$u = \epsilon_0 \mathbf{E}^2.$$

EM waves in vacuum: Energy and Momentum

- ▶ Using, $\mathbf{B} = \hat{\mathbf{k}} \times \frac{\mathbf{E}}{c}$, can show,

$$\frac{1}{2\mu_0} \mathbf{B}^2 = \frac{\epsilon_0}{2} \mathbf{E}^2, u_{\mathbf{E}} = u_{\mathbf{B}},$$

$$u = \epsilon_0 \mathbf{E}^2.$$

- ▶ Energy flux density (per unit cross-sectional area, per unit time), also called intensity

$$\mathbf{S} = \frac{\mathbf{E}^2}{\mu_0 c} \hat{\mathbf{k}} = \epsilon_0 c \mathbf{E}^2 \hat{\mathbf{k}}.$$

EM waves in vacuum: Energy and Momentum

- ▶ Using, $\mathbf{B} = \hat{\mathbf{k}} \times \frac{\mathbf{E}}{c}$, can show,

$$\frac{1}{2\mu_0} \mathbf{B}^2 = \frac{\epsilon_0}{2} \mathbf{E}^2, u_{\mathbf{E}} = u_{\mathbf{B}},$$

$$u = \epsilon_0 \mathbf{E}^2.$$

- ▶ Energy flux density (per unit cross-sectional area, per unit time), also called intensity

$$\mathbf{S} = \frac{\mathbf{E}^2}{\mu_0 c} \hat{\mathbf{k}} = \epsilon_0 c \mathbf{E}^2 \hat{\mathbf{k}}.$$

- ▶ Momentum density, π_{EM}

$$\pi_{EM} = \frac{1}{c^2} \mathbf{S}, \pi_{EM} = \frac{u}{c}.$$

EM waves in vacuum: Pressure of radiation

²Note that $\langle \mathbf{E}^2 \rangle = \frac{1}{2} \mathbf{E}_0^2$, since $\langle \sin^2 \theta \rangle = \frac{1}{2}$.

EM waves in vacuum: Pressure of radiation

- Absorption of EM waves: Momentum transferred to the absorber,

$$P = \frac{\text{Normal Force}}{\text{Area}} = \frac{\text{Force}}{\text{Cross-sectional area}} = \frac{\text{Momentum Transferred}}{\text{time} \times \text{Cross-sectional area}}$$

$$P = \pi_{EM} c, \langle P \rangle = \langle \pi_{EM} \rangle c = \frac{1}{2} \epsilon_0 \mathbf{E}_0^2$$

2

²Note that $\langle \mathbf{E}^2 \rangle = \frac{1}{2} \mathbf{E}_0^2$, since $\langle \sin^2 \theta \rangle = \frac{1}{2}$.

EM waves in vacuum: Pressure of radiation

- Absorption of EM waves: Momentum transferred to the absorber,

$$P = \frac{\text{Normal Force}}{\text{Area}} = \frac{\text{Force}}{\text{Cross-sectional area}} = \frac{\text{Momentum Transferred}}{\text{time} \times \text{Cross-sectional area}}$$

$$P = \pi_{EM} c, \langle P \rangle = \langle \pi_{EM} \rangle c = \frac{1}{2} \epsilon_0 \mathbf{E}_0^2$$

2

- Perfect reflector, twice as much momentum transfer

$$P = 2\pi_{EM} c = \epsilon_0 \mathbf{E}_0^2.$$

²Note that $\langle \mathbf{E}^2 \rangle = \frac{1}{2} \mathbf{E}_0^2$, since $\langle \sin^2 \theta \rangle = \frac{1}{2}$.

EM waves in vacuum: Pressure of radiation

- ▶ Absorption of EM waves: Momentum transferred to the absorber,

$$P = \frac{\text{Normal Force}}{\text{Area}} = \frac{\text{Force}}{\text{Cross-sectional area}} = \frac{\text{Momentum Transferred}}{\text{time} \times \text{Cross-sectional area}}$$

$$P = \pi_{EM} c, \langle P \rangle = \langle \pi_{EM} \rangle c = \frac{1}{2} \epsilon_0 \mathbf{E}_0^2$$

2

- ▶ Perfect reflector, twice as much momentum transfer

$$P = 2\pi_{EM} c = \epsilon_0 \mathbf{E}_0^2.$$

- ▶ Microscopic level, \mathbf{E} and \mathbf{B} push/pull on charges in the wall.

²Note that $\langle \mathbf{E}^2 \rangle = \frac{1}{2} \mathbf{E}_0^2$, since $\langle \sin^2 \theta \rangle = \frac{1}{2}$.

EM waves in matter

EM waves in matter

- ▶ Maxwell Equations in absence of sources in matter,

$$\nabla \cdot \mathbf{D} = 0, \quad \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0,$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{H} - \frac{\partial \mathbf{D}}{\partial t} = 0$$

EM waves in matter

- ▶ Maxwell Equations in absence of sources in matter,

$$\nabla \cdot \mathbf{D} = 0, \quad \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0,$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{H} - \frac{\partial \mathbf{D}}{\partial t} = 0$$

- ▶ Linear homogenous media: Constitutive relations,

$$\mathbf{D} = \epsilon \mathbf{E}, \quad \mathbf{B} = \mu \mathbf{H}.$$

EM waves in matter

- ▶ Maxwell Equations in absence of sources in matter,

$$\nabla \cdot \mathbf{D} = 0, \quad \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0,$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{H} - \frac{\partial \mathbf{D}}{\partial t} = 0$$

- ▶ Linear homogenous media: Constitutive relations,

$$\mathbf{D} = \epsilon \mathbf{E}, \quad \mathbf{B} = \mu \mathbf{H}.$$

- ▶ Maxwell's equations for linear homogenous media

$$\nabla \cdot \mathbf{E} = 0, \quad \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0,$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{B} - \epsilon\mu \frac{\partial \mathbf{E}}{\partial t} = 0.$$

EM waves in matter

- ▶ Maxwell Equations in absence of sources in matter,

$$\nabla \cdot \mathbf{D} = 0, \quad \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0,$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{H} - \frac{\partial \mathbf{D}}{\partial t} = 0$$

- ▶ Linear homogenous media: Constitutive relations,

$$\mathbf{D} = \epsilon \mathbf{E}, \quad \mathbf{B} = \mu \mathbf{H}.$$

- ▶ Maxwell's equations for linear homogenous media

$$\nabla \cdot \mathbf{E} = 0, \quad \nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0,$$

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{B} - \epsilon \mu \frac{\partial \mathbf{E}}{\partial t} = 0.$$

- ▶ Speed of EM waves,

$$v = \frac{1}{\sqrt{\mu\epsilon}} = \frac{c}{n},$$

n is called the refractive index.

EM waves in matter

EM waves in matter

- ▶ Energy density in media,

$$\begin{aligned} u &= \frac{1}{2} \mathbf{D} \cdot \mathbf{H} + \frac{1}{2} \mathbf{B} \cdot \mathbf{H} \\ &= \frac{\epsilon}{2} \mathbf{E}^2 + \frac{1}{2\mu} \mathbf{B}^2. \end{aligned}$$

EM waves in matter

- ▶ Energy density in media,

$$\begin{aligned}u &= \frac{1}{2} \mathbf{D} \cdot \mathbf{H} + \frac{1}{2} \mathbf{B} \cdot \mathbf{H} \\&= \frac{\epsilon}{2} \mathbf{E}^2 + \frac{1}{2\mu} \mathbf{B}^2.\end{aligned}$$

- ▶ Energy flux rate: Poynting vector,

$$\mathbf{S} = \frac{1}{\mu} (\mathbf{E} \times \mathbf{B})$$

EM waves in matter

- ▶ Energy density in media,

$$\begin{aligned}u &= \frac{1}{2} \mathbf{D} \cdot \mathbf{H} + \frac{1}{2} \mathbf{B} \cdot \mathbf{H} \\&= \frac{\epsilon}{2} \mathbf{E}^2 + \frac{1}{2\mu} \mathbf{B}^2.\end{aligned}$$

- ▶ Energy flux rate: Poynting vector,

$$\mathbf{S} = \frac{1}{\mu} (\mathbf{E} \times \mathbf{B})$$



$$\mathbf{B} = \hat{\mathbf{k}} \times \frac{\mathbf{E}}{v}.$$

EM waves in matter

- ▶ Energy density in media,

$$\begin{aligned}u &= \frac{1}{2} \mathbf{D} \cdot \mathbf{H} + \frac{1}{2} \mathbf{B} \cdot \mathbf{H} \\&= \frac{\epsilon}{2} \mathbf{E}^2 + \frac{1}{2\mu} \mathbf{B}^2.\end{aligned}$$

- ▶ Energy flux rate: Poynting vector,

$$\mathbf{S} = \frac{1}{\mu} (\mathbf{E} \times \mathbf{B})$$



$$\mathbf{B} = \hat{\mathbf{k}} \times \frac{\mathbf{E}}{v}.$$

- ▶ Boundary Conditions at the interface of media:

$$\epsilon^1 E_{\perp}^1 = \epsilon^2 E_{\perp}^2, \quad \mathbf{E}_{\parallel}^1 = \mathbf{E}_{\parallel}^2,$$

$$B_{\perp}^1 = B_{\perp}^2, \quad \frac{1}{\mu_1} \mathbf{B}_{\parallel}^1 = \frac{1}{\mu_2} \mathbf{B}_{\parallel}^2.$$

Reflection and Transmission of EM waves

Reflection and Transmission of EM waves

- ▶ Monochromatic waves incident on the interface from left (fig. 9.14 of Griffiths' text)

$$\mathbf{E}_I = \tilde{\mathbf{E}}_{0I} e^{i(\mathbf{k}_I \cdot \mathbf{x} - \omega t)}, \quad \mathbf{B}_I = \hat{\mathbf{k}}_I \times \frac{\mathbf{E}_I}{v_1}.$$

Reflection and Transmission of EM waves

- ▶ Monochromatic waves incident on the interface from left (fig. 9.14 of Griffiths' text)

$$\mathbf{E}_I = \tilde{\mathbf{E}}_{0I} e^{i(\mathbf{k}_I \cdot \mathbf{x} - \omega t)}, \quad \mathbf{B}_I = \hat{\mathbf{k}}_I \times \frac{\mathbf{E}_I}{v_1}.$$

- ▶ Reflected wave

$$\mathbf{E}_R = \tilde{\mathbf{E}}_{0R} e^{i(\mathbf{k}_R \cdot \mathbf{x} - \omega t)}, \quad \mathbf{B}_R = \hat{\mathbf{k}}_R \times \frac{\mathbf{E}_R}{v_1}.$$

Reflection and Transmission of EM waves

- ▶ Monochromatic waves incident on the interface from left (fig. 9.14 of Griffiths' text)

$$\mathbf{E}_I = \tilde{\mathbf{E}}_{0I} e^{i(\mathbf{k}_I \cdot \mathbf{x} - \omega t)}, \quad \mathbf{B}_I = \hat{\mathbf{k}}_I \times \frac{\mathbf{E}_I}{v_1}.$$

- ▶ Reflected wave

$$\mathbf{E}_R = \tilde{\mathbf{E}}_{0R} e^{i(\mathbf{k}_R \cdot \mathbf{x} - \omega t)}, \quad \mathbf{B}_R = \hat{\mathbf{k}}_R \times \frac{\mathbf{E}_R}{v_1}.$$

- ▶ Transmitted wave

$$\mathbf{E}_T = \tilde{\mathbf{E}}_{0T} e^{i(\mathbf{k}_T \cdot \mathbf{x} - \omega t)}, \quad \mathbf{B}_T = \hat{\mathbf{k}}_T \times \frac{\mathbf{E}_T}{v_2}.$$

Reflection and Transmission of EM waves

- ▶ Monochromatic waves incident on the interface from left (fig. 9.14 of Griffiths' text)

$$\mathbf{E}_I = \tilde{\mathbf{E}}_{0I} e^{i(\mathbf{k}_I \cdot \mathbf{x} - \omega t)}, \quad \mathbf{B}_I = \hat{\mathbf{k}}_I \times \frac{\mathbf{E}_I}{v_1}.$$

- ▶ Reflected wave

$$\mathbf{E}_R = \tilde{\mathbf{E}}_{0R} e^{i(\mathbf{k}_R \cdot \mathbf{x} - \omega t)}, \quad \mathbf{B}_R = \hat{\mathbf{k}}_R \times \frac{\mathbf{E}_R}{v_1}.$$

- ▶ Transmitted wave

$$\mathbf{E}_T = \tilde{\mathbf{E}}_{0T} e^{i(\mathbf{k}_T \cdot \mathbf{x} - \omega t)}, \quad \mathbf{B}_T = \hat{\mathbf{k}}_T \times \frac{\mathbf{E}_T}{v_2}.$$

- ▶ Frequency same,

$$\omega = |\mathbf{k}|v,$$

$$\Rightarrow |\mathbf{k}_I| = |\mathbf{k}_R| = \frac{\omega}{v_1}, \quad |\mathbf{k}_T| = \frac{\omega}{v_2} = \frac{v_1}{v_2} |\mathbf{k}_I| = \frac{n_2}{n_1} |\mathbf{k}_I|.$$

Reflection and Transmission of waves at an interface

Reflection and Transmission of waves at an interface

- We take the interface of the media to be the xy -plane i.e. $z = 0$.
Then,

$$\mathbf{k} \cdot \mathbf{x}|_{z=0} = (k_x x + k_y y + k_z z)|_{z=0} = k_x x + k_y y = \mathbf{k}^{\parallel} \cdot \mathbf{x}$$

where we have introduced \mathbf{k}^{\parallel} as the part of the wave-vector parallel to the interface

$$\mathbf{k}^{\parallel} = \mathbf{k} - k_z \hat{\mathbf{z}} = k_x \hat{\mathbf{x}} + k_y \hat{\mathbf{y}}.$$

Reflection and Transmission of waves at an interface

- We take the interface of the media to be the xy -plane i.e. $z = 0$. Then,

$$\mathbf{k} \cdot \mathbf{x}|_{z=0} = (k_x x + k_y y + k_z z)|_{z=0} = k_x x + k_y y = \mathbf{k}^{\parallel} \cdot \mathbf{x}$$

where we have introduced \mathbf{k}^{\parallel} as the part of the wave-vector parallel to the interface

$$\mathbf{k}^{\parallel} = \mathbf{k} - k_z \hat{\mathbf{z}} = k_x \hat{\mathbf{x}} + k_y \hat{\mathbf{y}}.$$

- Boundary conditions for Maxwell's equations at the interface i.e. $z = 0$ generically look like

$$(\dots) e^{i(\mathbf{k}_I \cdot \mathbf{x} - \omega t)} + (\dots) e^{i(\mathbf{k}_R \cdot \mathbf{x} - \omega t)} \Big|_{z=0} = (\dots) e^{i(\mathbf{k}_T \cdot \mathbf{x} - \omega t)} \Big|_{z=0} \quad \forall x, y, t,$$

Reflection and Transmission of waves at an interface

- ▶ We take the interface of the media to be the xy -plane i.e. $z = 0$. Then,

$$\mathbf{k} \cdot \mathbf{x}|_{z=0} = (k_x x + k_y y + k_z z)|_{z=0} = k_x x + k_y y = \mathbf{k}^{\parallel} \cdot \mathbf{x}$$

where we have introduced \mathbf{k}^{\parallel} as the part of the wave-vector parallel to the interface

$$\mathbf{k}^{\parallel} = \mathbf{k} - k_z \hat{\mathbf{z}} = k_x \hat{\mathbf{x}} + k_y \hat{\mathbf{y}}.$$

- ▶ Boundary conditions for Maxwell's equations at the interface i.e. $z = 0$ generically look like

$$(\dots) e^{i(\mathbf{k}_I \cdot \mathbf{x} - \omega t)} + (\dots) e^{i(\mathbf{k}_R \cdot \mathbf{x} - \omega t)} \Big|_{z=0} = (\dots) e^{i(\mathbf{k}_T \cdot \mathbf{x} - \omega t)} \Big|_{z=0} \quad \forall x, y, t,$$

- ▶ or equivalently

$$(\dots) e^{i(\mathbf{k}_I^{\parallel} \cdot \mathbf{x} - \omega t)} + (\dots) e^{i(\mathbf{k}_R^{\parallel} \cdot \mathbf{x} - \omega t)} = (\dots) e^{i(\mathbf{k}_T^{\parallel} \cdot \mathbf{x} - \omega t)} \quad \forall x, y, t.$$

Reflection and Transmission of waves at an interface

- ▶ We take the interface of the media to be the xy -plane i.e. $z = 0$. Then,

$$\mathbf{k} \cdot \mathbf{x}|_{z=0} = (k_x x + k_y y + k_z z)|_{z=0} = k_x x + k_y y = \mathbf{k}^{\parallel} \cdot \mathbf{x}$$

where we have introduced \mathbf{k}^{\parallel} as the part of the wave-vector parallel to the interface

$$\mathbf{k}^{\parallel} = \mathbf{k} - k_z \hat{\mathbf{z}} = k_x \hat{\mathbf{x}} + k_y \hat{\mathbf{y}}.$$

- ▶ Boundary conditions for Maxwell's equations at the interface i.e. $z = 0$ generically look like

$$(\dots) e^{i(\mathbf{k}_I \cdot \mathbf{x} - \omega t)} + (\dots) e^{i(\mathbf{k}_R \cdot \mathbf{x} - \omega t)} \Big|_{z=0} = (\dots) e^{i(\mathbf{k}_T \cdot \mathbf{x} - \omega t)} \Big|_{z=0} \quad \forall x, y, t,$$

- ▶ or equivalently

$$(\dots) e^{i(\mathbf{k}_I^{\parallel} \cdot \mathbf{x} - \omega t)} + (\dots) e^{i(\mathbf{k}_R^{\parallel} \cdot \mathbf{x} - \omega t)} = (\dots) e^{i(\mathbf{k}_T^{\parallel} \cdot \mathbf{x} - \omega t)} \quad \forall x, y, t.$$

- ▶ Since this is a complex equation the phases on both sides must be equal,

$$\Rightarrow \mathbf{k}_I^{\parallel} \cdot \mathbf{x} = \mathbf{k}_R^{\parallel} \cdot \mathbf{x} = \mathbf{k}_T^{\parallel} \cdot \mathbf{x}, \quad \forall x, y$$

Reflection and Transmission of waves at an interface

- ▶ We take the interface of the media to be the xy -plane i.e. $z = 0$. Then,

$$\mathbf{k} \cdot \mathbf{x}|_{z=0} = (k_x x + k_y y + k_z z)|_{z=0} = k_x x + k_y y = \mathbf{k}^{\parallel} \cdot \mathbf{x}$$

where we have introduced \mathbf{k}^{\parallel} as the part of the wave-vector parallel to the interface

$$\mathbf{k}^{\parallel} = \mathbf{k} - k_z \hat{\mathbf{z}} = k_x \hat{\mathbf{x}} + k_y \hat{\mathbf{y}}.$$

- ▶ Boundary conditions for Maxwell's equations at the interface i.e. $z = 0$ generically look like

$$(\dots) e^{i(\mathbf{k}_I \cdot \mathbf{x} - \omega t)} + (\dots) e^{i(\mathbf{k}_R \cdot \mathbf{x} - \omega t)} \Big|_{z=0} = (\dots) e^{i(\mathbf{k}_T \cdot \mathbf{x} - \omega t)} \Big|_{z=0} \quad \forall x, y, t,$$

- ▶ or equivalently

$$(\dots) e^{i(\mathbf{k}_I^{\parallel} \cdot \mathbf{x} - \omega t)} + (\dots) e^{i(\mathbf{k}_R^{\parallel} \cdot \mathbf{x} - \omega t)} = (\dots) e^{i(\mathbf{k}_T^{\parallel} \cdot \mathbf{x} - \omega t)} \quad \forall x, y, t.$$

- ▶ Since this is a complex equation the phases on both sides must be equal,

$$\Rightarrow \mathbf{k}_I^{\parallel} \cdot \mathbf{x} = \mathbf{k}_R^{\parallel} \cdot \mathbf{x} = \mathbf{k}_T^{\parallel} \cdot \mathbf{x}, \quad \forall x, y$$

Reflection and Transmission of waves at an interface

Reflection and Transmission of waves at an interface

► From last slide

$$\Rightarrow \mathbf{k}_I^{\parallel} = \mathbf{k}_R^{\parallel} = \mathbf{k}_T^{\parallel}$$

Reflection and Transmission of waves at an interface

- From last slide

$$\Rightarrow \mathbf{k}_I^{\parallel} = \mathbf{k}_R^{\parallel} = \mathbf{k}_T^{\parallel}$$

- First law: All three wave vectors lie in a plane (plane of incidence containing \mathbf{k}_I and $\hat{\mathbf{z}}$).

$$|\mathbf{k}_I| \sin \theta_I = |\mathbf{k}_R| \sin \theta_R = |\mathbf{k}_T| \sin \theta_T.$$

Reflection and Transmission of waves at an interface

- From last slide

$$\Rightarrow \mathbf{k}_I^{\parallel} = \mathbf{k}_R^{\parallel} = \mathbf{k}_T^{\parallel}$$

- First law: All three wave vectors lie in a plane (plane of incidence containing \mathbf{k}_I and $\hat{\mathbf{z}}$).

$$|\mathbf{k}_I| \sin \theta_I = |\mathbf{k}_R| \sin \theta_R = |\mathbf{k}_T| \sin \theta_T.$$

- Second law: Angle of reflection = Angle of incidence,

$$\theta_I = \theta_R.$$

Reflection and Transmission of waves at an interface

- ▶ From last slide

$$\Rightarrow \mathbf{k}_I^{\parallel} = \mathbf{k}_R^{\parallel} = \mathbf{k}_T^{\parallel}$$

- ▶ First law: All three wave vectors lie in a plane (plane of incidence containing \mathbf{k}_I and $\hat{\mathbf{z}}$).

$$|\mathbf{k}_I| \sin \theta_I = |\mathbf{k}_R| \sin \theta_R = |\mathbf{k}_T| \sin \theta_T.$$

- ▶ Second law: Angle of reflection = Angle of incidence,

$$\theta_I = \theta_R.$$

- ▶ Third law: Snell's law of Refraction

$$\frac{\sin \theta_I}{\sin \theta_T} = \frac{|\mathbf{k}_T|}{|\mathbf{k}_I|} = \frac{v_1}{v_2} = \frac{n_2}{n_1}$$

Satisfying boundary conditions

Satisfying boundary conditions

- Boundary conditions relates amplitudes

$$\epsilon^1 \left(\tilde{E}_{0I}^3 + \tilde{E}_{0R}^3 \right) = \epsilon^2 \tilde{E}_{0T}^3,$$

$$\tilde{B}_{0I}^3 + \tilde{B}_{0R}^3 = \tilde{B}_{0T}^3,$$

$$\tilde{E}_{0I}^{1,2} + \tilde{E}_{0R}^{1,2} = \tilde{E}_{0T}^{1,2},$$

$$\frac{1}{\mu_1} \left(\tilde{B}_{0I}^{1,2} + \tilde{B}_{0R}^{1,2} \right) = \frac{1}{\mu_2} \tilde{B}_{0T}^{1,2}.$$

Satisfying boundary conditions

- Boundary conditions relates amplitudes

$$\epsilon^1 \left(\tilde{E}_{0I}^3 + \tilde{E}_{0R}^3 \right) = \epsilon^2 \tilde{E}_{0T}^3,$$

$$\tilde{B}_{0I}^3 + \tilde{B}_{0R}^3 = \tilde{B}_{0T}^3,$$

$$\tilde{E}_{0I}^{1,2} + \tilde{E}_{0R}^{1,2} = \tilde{E}_{0T}^{1,2},$$

$$\frac{1}{\mu_1} \left(\tilde{B}_{0I}^{1,2} + \tilde{B}_{0R}^{1,2} \right) = \frac{1}{\mu_2} \tilde{B}_{0T}^{1,2}.$$

- Case I: “In plane” polarization,

$$\epsilon^1 \left(- \left| \tilde{\mathbf{E}}_{0I} \right| \sin \theta_I + \left| \tilde{\mathbf{E}}_{0R} \right| \sin \theta_R \right) = -\epsilon^2 \left| \tilde{\mathbf{E}}_{0T} \right| \sin \theta_T$$

$$\left| \tilde{\mathbf{E}}_{0I} \right| \cos \theta_I + \left| \tilde{\mathbf{E}}_{0R} \right| \cos \theta_R = \left| \tilde{\mathbf{E}}_{0T} \right| \cos \theta_T$$

$$\frac{1}{\mu_1 v_1} \left(\left| \tilde{\mathbf{E}}_{0I} \right| - \left| \tilde{\mathbf{E}}_{0R} \right| \right) = \frac{1}{\mu_2 v_2} \left| \tilde{\mathbf{E}}_{0T} \right|$$

Fresnel Equations

Fresnel Equations

► We have,

$$\left| \tilde{\mathbf{E}}_{0I} \right| - \left| \tilde{\mathbf{E}}_{0R} \right| = \beta \left| \tilde{\mathbf{E}}_{0T} \right|, \beta = \frac{\mu_1 v_1}{\mu_2 v_2} = \frac{\mu_1 n_2}{\mu_2 n_1}$$

$$\left| \tilde{\mathbf{E}}_{0I} \right| + \left| \tilde{\mathbf{E}}_{0R} \right| = \alpha \left| \tilde{\mathbf{E}}_{0T} \right|, \alpha = \frac{\cos \theta_T}{\cos \theta_I}$$

Fresnel Equations

- We have,

$$\left| \tilde{\mathbf{E}}_{0I} \right| - \left| \tilde{\mathbf{E}}_{0R} \right| = \beta \left| \tilde{\mathbf{E}}_{0T} \right|, \beta = \frac{\mu_1 v_1}{\mu_2 v_2} = \frac{\mu_1 n_2}{\mu_2 n_1}$$

$$\left| \tilde{\mathbf{E}}_{0I} \right| + \left| \tilde{\mathbf{E}}_{0R} \right| = \alpha \left| \tilde{\mathbf{E}}_{0T} \right|, \alpha = \frac{\cos \theta_T}{\cos \theta_I}$$

- Fresnel's Equations for “in plane” component of polarization

$$\frac{\left| \tilde{\mathbf{E}}_{0R} \right|}{\left| \tilde{\mathbf{E}}_{0I} \right|} = \frac{\alpha - \beta}{\alpha + \beta},$$

$$\frac{\left| \tilde{\mathbf{E}}_{0T} \right|}{\left| \tilde{\mathbf{E}}_{0I} \right|} = \frac{2}{\alpha + \beta}$$

Fresnel Equations

- We have,

$$\left| \tilde{\mathbf{E}}_{0I} \right| - \left| \tilde{\mathbf{E}}_{0R} \right| = \beta \left| \tilde{\mathbf{E}}_{0T} \right|, \beta = \frac{\mu_1 v_1}{\mu_2 v_2} = \frac{\mu_1 n_2}{\mu_2 n_1}$$

$$\left| \tilde{\mathbf{E}}_{0I} \right| + \left| \tilde{\mathbf{E}}_{0R} \right| = \alpha \left| \tilde{\mathbf{E}}_{0T} \right|, \alpha = \frac{\cos \theta_T}{\cos \theta_I}$$

- Fresnel's Equations for “in plane” component of polarization

$$\frac{\left| \tilde{\mathbf{E}}_{0R} \right|}{\left| \tilde{\mathbf{E}}_{0I} \right|} = \frac{\alpha - \beta}{\alpha + \beta},$$

$$\frac{\left| \tilde{\mathbf{E}}_{0T} \right|}{\left| \tilde{\mathbf{E}}_{0I} \right|} = \frac{2}{\alpha + \beta}$$

- Brewster Angle: Full transmission

$$\alpha = \beta,$$

Fresnel Equations

- We have,

$$\left| \tilde{\mathbf{E}}_{0I} \right| - \left| \tilde{\mathbf{E}}_{0R} \right| = \beta \left| \tilde{\mathbf{E}}_{0T} \right|, \beta = \frac{\mu_1 v_1}{\mu_2 v_2} = \frac{\mu_1 n_2}{\mu_2 n_1}$$

$$\left| \tilde{\mathbf{E}}_{0I} \right| + \left| \tilde{\mathbf{E}}_{0R} \right| = \alpha \left| \tilde{\mathbf{E}}_{0T} \right|, \alpha = \frac{\cos \theta_T}{\cos \theta_I}$$

- Fresnel's Equations for “in plane” component of polarization

$$\frac{\left| \tilde{\mathbf{E}}_{0R} \right|}{\left| \tilde{\mathbf{E}}_{0I} \right|} = \frac{\alpha - \beta}{\alpha + \beta},$$

$$\frac{\left| \tilde{\mathbf{E}}_{0T} \right|}{\left| \tilde{\mathbf{E}}_{0I} \right|} = \frac{2}{\alpha + \beta}$$

- Brewster Angle: Full transmission

$$\alpha = \beta,$$

- Next class: EM waves incident on conductors

Quiz

1. Point out where rationalized units are not used

$$(a) \mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{|\mathbf{x}|^3} \mathbf{x}, \quad (b) \mathbf{B} = \frac{\mu_0}{4\pi} \oint \frac{I d\mathbf{x}' \times (\mathbf{x} - \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|^3}, \quad (c) \nabla \cdot \mathbf{E} = 4\pi\rho, \quad (d) \nabla \times \mathbf{B} = \frac{4\pi}{c} \mathbf{j}$$

2. Direction of \mathbf{E} of point charge at origin in spherical polar coordinates is along

$$(a) \hat{\mathbf{r}}, \quad (b) \hat{\boldsymbol{\theta}}, \quad (c) \hat{\boldsymbol{\phi}}, \quad (d) \hat{\boldsymbol{\rho}}$$

3. Direction of \mathbf{B} of infinitely long current carrying wire along z-axis in cylindrical coordinates is along

$$(a) \hat{\boldsymbol{\rho}}, \quad (b) \hat{\boldsymbol{\phi}}, \quad (c) \hat{\mathbf{z}}, \quad (d) \rho \hat{\boldsymbol{\rho}} + z \hat{\mathbf{z}}$$

4. Magnetic field $\mathbf{B}(\mathbf{x})$ due to a point charge, q at \mathbf{x}' moving with vel, \mathbf{v}

$$(a) \text{absent}, \quad (b) \text{approx} \propto q \frac{\mathbf{v}}{c} \times \frac{\mathbf{x}}{|\mathbf{x} - \mathbf{v}t|^3}, \quad (c) \text{approx} \propto q \frac{\mathbf{v}}{c} \times \frac{\mathbf{x} - \mathbf{v}t}{|\mathbf{x} - \mathbf{v}t|^3}, \quad (d) \text{approx} \propto q \frac{\mathbf{v}}{c} \times \mathbf{E}$$

5. Inside a conductor, in general

$$(a) \mathbf{E} = 0, \quad (b) \rho = 0, \quad (c) \mathbf{j} = \sigma \mathbf{E} \quad (d) \mathbf{E} = \sigma \mathbf{j}$$

Quiz

1. Point out where rationalized units are not used

$$(a) \mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{|\mathbf{x}|^3} \mathbf{x}, \quad (b) \mathbf{B} = \frac{\mu_0}{4\pi} \oint \frac{I d\mathbf{x}' \times (\mathbf{x} - \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|^3}, \quad (c) \nabla \cdot \mathbf{E} = 4\pi\rho, \quad (d) \nabla \times \mathbf{B} = \frac{4\pi}{c} \mathbf{j}$$

2. Direction of \mathbf{E} of point charge at origin in spherical polar coordinates is along

$$(a) \hat{\mathbf{r}}, \quad (b) \hat{\boldsymbol{\theta}}, \quad (c) \hat{\boldsymbol{\phi}}, \quad (d) \hat{\boldsymbol{\rho}}$$

3. Direction of \mathbf{B} of infinitely long current carrying wire along z-axis in cylindrical coordinates is along

$$(a) \hat{\boldsymbol{\rho}}, \quad (b) \hat{\boldsymbol{\phi}}, \quad (c) \hat{\mathbf{z}}, \quad (d) \rho \hat{\boldsymbol{\rho}} + z \hat{\mathbf{z}}$$

4. Magnetic field $\mathbf{B}(\mathbf{x})$ due to a point charge, q at \mathbf{x}' moving with vel, \mathbf{v}

$$(a) \text{absent}, \quad (b) \text{approx} \propto q \frac{\mathbf{v}}{c} \times \frac{\mathbf{x}}{|\mathbf{x} - \mathbf{v}t|^3}, \quad (c) \text{approx} \propto q \frac{\mathbf{v}}{c} \times \frac{\mathbf{x} - \mathbf{v}t}{|\mathbf{x} - \mathbf{v}t|^3}, \quad (d) \text{approx} \propto q \frac{\mathbf{v}}{c} \times \mathbf{E}$$

5. Inside a conductor, in general

$$(a) \mathbf{E} = 0, \quad (b) \rho = 0, \quad (c) \mathbf{j} = \sigma \mathbf{E} \quad (d) \mathbf{E} = \sigma \mathbf{j}$$

Quiz

6. At the boundary of a pair of dielectrics cum paramagnetics which of the following is not continuous

$$(a) \mathbf{E}_{\parallel}, \quad (b) \mathbf{H}_{\parallel} \quad (c) D_{\perp} \quad (d) B_{\perp}$$

7. If \mathbf{E} , \mathbf{B} and ϕ , \mathbf{A} contain same amount of information, why are they different in number of components?

Quiz

6. At the boundary of a pair of dielectrics cum paramagnetics which of the following is not continuous

$$(a) \mathbf{E}_{\parallel}, \quad (b) \mathbf{H}_{\parallel} \quad (c) D_{\perp} \quad (d) B_{\perp}$$

7. If \mathbf{E} , \mathbf{B} and ϕ , \mathbf{A} contain same amount of information, why are they different in number of components?