

# EP 1027: Maxwell's Equations and Electromagnetic Waves

Instructor: Shubho Roy<sup>1</sup>  
(Dept. of Physics)

Lecture 2

March 27, 2019

---

<sup>1</sup> Office: C 313 D

Email: [sroy@iith.ac.in](mailto:sroy@iith.ac.in)

Office hrs. - Email appointment or walk in

# Agenda: Algebra and Calculus of vectors and tensors

# Agenda: Algebra and Calculus of vectors and tensors

- ▶ Short Recap of Lecture 1

# Agenda: Algebra and Calculus of vectors and tensors

- ▶ Short Recap of Lecture 1
- ▶ Define vectors, scalars, and tensors *using their transformation law under rotation of coordinate axes*

# Agenda: Algebra and Calculus of vectors and tensors

- ▶ Short Recap of Lecture 1
- ▶ Define vectors, scalars, and tensors *using their transformation law under rotation of coordinate axes*
- ▶ Define vector *field*, scalar *field* and general tensor *field*

# Agenda: Algebra and Calculus of vectors and tensors

- ▶ Short Recap of Lecture 1
- ▶ Define vectors, scalars, and tensors *using their transformation law under rotation of coordinate axes*
- ▶ Define vector *field*, scalar *field* and general tensor *field*
- ▶ Differentiation of vector fields : Gradient, Divergence, Curl, Laplacian operators

# Agenda: Algebra and Calculus of vectors and tensors

- ▶ Short Recap of Lecture 1
- ▶ Define vectors, scalars, and tensors *using their transformation law under rotation of coordinate axes*
- ▶ Define vector *field*, scalar *field* and general tensor *field*
- ▶ Differentiation of vector fields : Gradient, Divergence, Curl, Laplacian operators
- ▶ Integration of vector fields: Gauss' and Stokes' Theorem

# References/Readings



# References/Readings

- ▶ Griffiths, D.J., **Introduction to Electrodynamics, Ch. 1**
- ▶ Boas, M. L., **Mathematical Methods... Ch. 6**
- ▶ Spiegel M.R., **Schaum's Outline of Vector Analysis**
- ▶ Schey, H.M., **Div, Grad, Curl and All that - An Informal Text on Vector Calculus**

# Recap of Cartesian Vectors

# Recap of Cartesian Vectors

- ▶ Vectors:
  - a) Quantity w/ both magnitude and a direction (directed line segment).
  - b) behaves like position vector under addition and multiplication by a scalar, **and transforms like it under rotations.**

# Recap of Cartesian Vectors

- ▶ Vectors:
  - a) Quantity w/ both magnitude and a direction (directed line segment).
  - b) behaves like position vector under addition and multiplication by a scalar, **and transforms like it under rotations**.
- ▶ Position vector: 3 component object (triplet) giving the location from a origin of Cartesian coordinate system: row vector  $(x, y, z)$  or a column vector  $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$  or  $\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$ .

# Recap of Cartesian Vectors

- ▶ Vectors:
  - a) Quantity w/ both magnitude and a direction (directed line segment).
  - b) behaves like position vector under addition and multiplication by a scalar, **and transforms like it under rotations**.
- ▶ Position vector: 3 component object (triplet) giving the location from a origin of Cartesian coordinate system: row vector  $(x, y, z)$  or a column vector  $\begin{pmatrix} x \\ y \\ z \end{pmatrix}$  or  $\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$ .
- ▶ Notation: General vector -  $\begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}$ . Call the components of a general vector:  $a_k$ ,  $k$  can take values from 1 to 3, denoted by  $\mathbf{a}$  or  $\vec{a}$ .

# Recap of Cartesian Vectors

# Recap of Cartesian Vectors

- ▶ Rule for addition of 2 vectors: Add the respective components

$$\mathbf{a} + \mathbf{b} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} \equiv \begin{pmatrix} a_1 + b_1 \\ a_2 + b_2 \\ a_3 + c_3 \end{pmatrix}$$

# Recap of Cartesian Vectors

- ▶ Rule for addition of 2 vectors: Add the respective components

$$\mathbf{a} + \mathbf{b} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} \equiv \begin{pmatrix} a_1 + b_1 \\ a_2 + b_2 \\ a_3 + c_3 \end{pmatrix}$$

- ▶ Rule for multiplication with a number (scalar),  $\lambda$  :  
Multiply each component by  $\lambda$

$$\lambda \mathbf{a} = \lambda \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \equiv \begin{pmatrix} \lambda a_1 \\ \lambda a_2 \\ \lambda a_3 \end{pmatrix}.$$



# Recap of Cartesian Vectors

- ▶ Rule for addition of 2 vectors: Add the respective components

$$\mathbf{a} + \mathbf{b} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} + \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} \equiv \begin{pmatrix} a_1 + b_1 \\ a_2 + b_2 \\ a_3 + b_3 \end{pmatrix}$$

- ▶ Rule for multiplication with a number (scalar),  $\lambda$  :  
Multiply each component by  $\lambda$

$$\lambda \mathbf{a} = \lambda \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \equiv \begin{pmatrix} \lambda a_1 \\ \lambda a_2 \\ \lambda a_3 \end{pmatrix}.$$

- ▶ Once we choose the vector to be a column, then to denote the row vector, we will use the transpose

$$\mathbf{a}^T = (a_1, a_2, a_3)$$

# Recap of Cartesian Vectors

# Recap of Cartesian Vectors

- Notation: In terms of basis (Unit) Vectors,  $\hat{\mathbf{e}}_k$

$$\begin{aligned}\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} &= a_1 \underbrace{\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}}_{\hat{\mathbf{e}}_1} + a_2 \underbrace{\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}}_{\hat{\mathbf{e}}_2} + a_3 \underbrace{\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}}_{\hat{\mathbf{e}}_3} \\ &= a_1 \hat{\mathbf{e}}_1 + a_2 \hat{\mathbf{e}}_2 + a_3 \hat{\mathbf{e}}_3 \\ &= \sum_{k=1}^3 a_k \hat{\mathbf{e}}_k. \\ &= a_k \hat{\mathbf{e}}_k\end{aligned}$$

# Recap of Cartesian Vectors

- Notation: In terms of basis (Unit) Vectors,  $\hat{\mathbf{e}}_k$

$$\begin{aligned}\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} &= a_1 \underbrace{\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}}_{\hat{\mathbf{e}}_1} + a_2 \underbrace{\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}}_{\hat{\mathbf{e}}_2} + a_3 \underbrace{\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}}_{\hat{\mathbf{e}}_3} \\ &= a_1 \hat{\mathbf{e}}_1 + a_2 \hat{\mathbf{e}}_2 + a_3 \hat{\mathbf{e}}_3 \\ &= \sum_{k=1}^3 a_k \hat{\mathbf{e}}_k. \\ &= a_k \hat{\mathbf{e}}_k\end{aligned}$$

- “Dot product/ inner product/ scalar product” of  $\mathbf{a}$  and  $\mathbf{b}$ :

$$\mathbf{a} \cdot \mathbf{b} \equiv \mathbf{a}^T \mathbf{b} = a_1 b_1 + a_2 b_2 + a_3 b_3 = \sum_{k=1}^3 a_k b_k = a_k b_k = a_l b_m \delta_{lm}.$$

# Recap of Cartesian Vectors

# Recap of Cartesian Vectors

- ▶ **Norm** (or size or magnitude) of a vector:

$$||\mathbf{a}|| \equiv \sqrt{\mathbf{a} \cdot \mathbf{a}} = \sqrt{a_k a_k}$$

# Recap of Cartesian Vectors

- ▶ **Norm** (or size or magnitude) of a vector:

$$||\mathbf{a}|| \equiv \sqrt{\mathbf{a} \cdot \mathbf{a}} = \sqrt{a_k a_k}$$

- ▶ Vector product/Cross product

$$\mathbf{a} \times \mathbf{b} \equiv (\epsilon_{ijk} a_i b_j) \hat{\mathbf{e}}_k,$$

i.e.

$$(\mathbf{a} \times \mathbf{b})_k = \epsilon_{ijk} a_i b_j,$$

where

$$\epsilon_{123} = \epsilon_{231} = \epsilon_{312} = 1,$$

$$\epsilon_{132} = \epsilon_{213} = \epsilon_{321} = -1,$$

$$\epsilon_{112} = \epsilon_{121} = \epsilon_{223} \dots = 0.$$

$\epsilon_{ijk}$  = “Levi Civita Symbol” or “Completely antisymmetric symbol”,

# Recap of Cartesian Vectors

- ▶ **Norm** (or size or magnitude) of a vector:

$$||\mathbf{a}|| \equiv \sqrt{\mathbf{a} \cdot \mathbf{a}} = \sqrt{a_k a_k}$$

- ▶ Vector product/Cross product

$$\mathbf{a} \times \mathbf{b} \equiv (\epsilon_{ijk} a_i b_j) \hat{\mathbf{e}}_k,$$

i.e.

$$(\mathbf{a} \times \mathbf{b})_k = \epsilon_{ijk} a_i b_j,$$

where

$$\epsilon_{123} = \epsilon_{231} = \epsilon_{312} = 1,$$

$$\epsilon_{132} = \epsilon_{213} = \epsilon_{321} = -1,$$

$$\epsilon_{112} = \epsilon_{121} = \epsilon_{223} \dots = 0.$$

$\epsilon_{ijk}$  = “Levi Civita Symbol” or “Completely antisymmetric symbol”,

- ▶ we derived:  $(\mathbf{a} \times \mathbf{b})_1 = a_2 b_3 - a_3 b_2$ .



# Recap of Cartesian Vectors

# Recap of Cartesian Vectors

- ▶ Vector transformation rule under rotation of coordinate axes: Implemented through Matrix operations,  $\mathbf{O}$

$$\mathbf{x}' = \mathbf{O}\mathbf{x},$$
$$\begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} O_{11} & O_{12} & O_{13} \\ O_{21} & O_{22} & \dots \\ O_{31} & O_{32} & \dots \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix},$$

# Recap of Cartesian Vectors

- ▶ Vector transformation rule under rotation of coordinate axes: Implemented through Matrix operations,  $\mathbf{O}$

$$\mathbf{x}' = \mathbf{O}\mathbf{x},$$

$$\begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} O_{11} & O_{12} & O_{13} \\ O_{21} & O_{22} & \dots \\ O_{31} & O_{32} & \dots \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix},$$



$$x'_i = O_{ij}x_j$$

# Recap of Cartesian Vectors

- ▶ Vector transformation rule under rotation of coordinate axes: Implemented through Matrix operations,  $\mathbf{O}$

$$\mathbf{x}' = \mathbf{O}\mathbf{x},$$

$$\begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} O_{11} & O_{12} & O_{13} \\ O_{21} & O_{22} & \dots \\ O_{31} & O_{32} & \dots \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix},$$



$$x'_i = O_{ij}x_j$$

- ▶ Rotation matrices = Orthogonal:  $\underbrace{\mathbf{O}^T \mathbf{O}}_{O_{ji} O_{jk}} = \underbrace{\mathbf{I}}_{\delta_{ik}}$ , to preserve lengths

# Recap of Cartesian Vectors

- ▶ Vector transformation rule under rotation of coordinate axes: Implemented through Matrix operations,  $\mathbf{O}$

$$\mathbf{x}' = \mathbf{O}\mathbf{x},$$

$$\begin{pmatrix} x'_1 \\ x'_2 \\ x'_3 \end{pmatrix} = \begin{pmatrix} O_{11} & O_{12} & O_{13} \\ O_{21} & O_{22} & \dots \\ O_{31} & O_{32} & \dots \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix},$$



$$x'_i = O_{ij}x_j$$

- ▶ Rotation matrices = Orthogonal:  $\underbrace{\mathbf{O}^T \mathbf{O}}_{O_{ji} O_{jk}} = \underbrace{\mathbf{I}}_{\delta_{ik}}$ , to preserve lengths

- ▶ (anticlockwise) Rotation around z-axis/3-axis by  $\theta$ ,

$$\mathbf{O} = \begin{pmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

# Cartesian Vectors (and tensors): General treatment

- ▶ Vector: Objects whose components transform under rotation of coordinate axes as

$$a'_i = O_{ij}a_j$$

# Cartesian Vectors (and tensors): General treatment

- ▶ Vector: Objects whose components transform under rotation of coordinate axes as

$$a'_i = O_{ij} a_j$$

- ▶ Scalar: Objects w/o free indices, remain unchanged under rotations

$$\phi' = \phi,$$

e.g.,  $\phi = \mathbf{a} \cdot \mathbf{b}$  (Check), mass, time, norm/magnitude of a vector, . . .

# Cartesian Vectors (and tensors): General treatment

- ▶ Vector: Objects whose components transform under rotation of coordinate axes as

$$a'_i = O_{ij} a_j$$

- ▶ Scalar: Objects w/o free indices, remain unchanged under rotations

$$\phi' = \phi,$$

e.g.,  $\phi = \mathbf{a} \cdot \mathbf{b}$  (Check), mass, time, norm/magnitude of a vector, . . .

- ▶ Tensor: Objects with components having multiple indices,  $T_{i_1 \dots i_p}$  (rank  $p$  tensor)



# Cartesian Vectors (and tensors): General treatment

- ▶ Vector: Objects whose components transform under rotation of coordinate axes as

$$a'_i = O_{ij} a_j$$

- ▶ Scalar: Objects w/o free indices, remain unchanged under rotations

$$\phi' = \phi,$$

e.g.,  $\phi = \mathbf{a} \cdot \mathbf{b}$  (Check), mass, time, norm/magnitude of a vector, . . .

- ▶ Tensor: Objects with components having multiple indices,  $T_{i_1 \dots i_p}$  (rank  $p$  tensor)
- ▶ Rank 2 tensor: Under coordinate axes rotation,

$$T'_{ij} = O_{il} O_{jm} T_{lm},$$

e.g., Outer product,  $a_i b_j$ ; Kronecker delta,  $\delta_{ij}$ ; Moment of Inertia,  $I_{ij} = m (\delta_{ij} x_k x_k - x_i x_j)$ .

# Invariant Cartesian tensors: Kronecker delta

- ▶ Invariant tensors: Components unchanged under rotations; how can they be tensors and not scalars?

# Invariant Cartesian tensors: Kronecker delta

- ▶ Invariant tensors: Components unchanged under rotations; how can they be tensors and not scalars?
- ▶ Kronecker delta ( $\delta_{ij}$ )

$$\delta'_{ij} = \delta_{ij},$$

# Invariant Cartesian tensors: Kronecker delta

- ▶ Invariant tensors: Components unchanged under rotations; how can they be tensors and not scalars?
- ▶ Kronecker delta ( $\delta_{ij}$ )

$$\delta'_{ij} = \delta_{ij},$$

- ▶ Proof:

$$\begin{aligned}\delta'_{ij} &= O_{il} O_{jm} \delta_{lm} \\ &= O_{il} O_{jl} \\ &= O_{il} (O^T)_{lj} \\ &= (OO^T)_{ij} \\ &= \delta_{ij}.\end{aligned}$$

# Invariant Cartesian Tensors: Levi-Civita

# Invariant Cartesian Tensors: Levi-Civita

- First Recall: Definition of determinant,

$$|M| = M_{1l} M_{2m} M_{3n} \epsilon_{lmn},$$

$$\epsilon_{ijk} |M| = M_{il} M_{jm} M_{kn} \epsilon_{lmn},$$

# Invariant Cartesian Tensors: Levi-Civita

- First Recall: Definition of determinant,

$$|M| = M_{1l} M_{2m} M_{3n} \epsilon_{lmn},$$

$$\epsilon_{ijk} |M| = M_{il} M_{jm} M_{kn} \epsilon_{lmn},$$

- Levi-Civita tensor,  $\epsilon_{ijk}$ ,

$$\epsilon'_{ijk} = O_{il} O_{jm} O_{kn} \epsilon_{lmn}$$

$$= |O| \epsilon_{ijk}$$

$$= \epsilon_{ijk},$$

iff  $|O| = +1$ . Rotation matrix is indeed has unit determinant, thus Levi Civita is a tensor under rotation of coordinate axes.

# Invariant Cartesian Tensors: Levi-Civita

- ▶ First Recall: Definition of determinant,

$$|M| = M_{1l} M_{2m} M_{3n} \epsilon_{lmn},$$

$$\epsilon_{ijk} |M| = M_{il} M_{jm} M_{kn} \epsilon_{lmn},$$

- ▶ Levi-Civita tensor,  $\epsilon_{ijk}$ ,

$$\epsilon'_{ijk} = O_{il} O_{jm} O_{kn} \epsilon_{lmn}$$

$$= |O| \epsilon_{ijk}$$

$$= \epsilon_{ijk},$$

iff  $|O| = +1$ . Rotation matrix is indeed has unit determinant, thus Levi Civita is a tensor under rotation of coordinate axes.

- ▶ Caveat: Under parity (inversion of coordinate axes)  $|O| = -1$ , and Levi Civita is then not a tensor.



# Tensor fields: Differentiation

# Tensor fields: Differentiation

- Components are functions of position,

$$\phi(\mathbf{x}), V_k(\mathbf{x}), T_{ij}(\mathbf{x})$$

# Tensor fields: Differentiation

- Components are functions of position,

$$\phi(\mathbf{x}), V_k(\mathbf{x}), T_{ij}(\mathbf{x})$$

- Examples:

$$\rho(\mathbf{x}), E_k(\mathbf{x}), \chi_{ij}(\mathbf{x}).$$

# Tensor fields: Differentiation

- Components are functions of position,

$$\phi(\mathbf{x}), V_k(\mathbf{x}), T_{ij}(\mathbf{x})$$

- Examples:

$$\rho(\mathbf{x}), E_k(\mathbf{x}), \chi_{ij}(\mathbf{x}).$$

- Gradient of a scalar

$$\begin{aligned}\phi(\mathbf{x} + d\mathbf{x}) - \phi(\mathbf{x}) &= dx_1 \frac{\partial \phi}{\partial x_1} + dx_2 \frac{\partial \phi}{\partial x_2} + dx_3 \frac{\partial \phi}{\partial x_3} \\ &= dx_i \frac{\partial \phi}{\partial x^i} \\ &= d\mathbf{x} \cdot (\nabla \phi),\end{aligned}$$

# Tensor fields: Differentiation

- Components are functions of position,

$$\phi(\mathbf{x}), V_k(\mathbf{x}), T_{ij}(\mathbf{x})$$

- Examples:

$$\rho(\mathbf{x}), E_k(\mathbf{x}), \chi_{ij}(\mathbf{x}).$$

- Gradient of a scalar

$$\begin{aligned}\phi(\mathbf{x} + d\mathbf{x}) - \phi(\mathbf{x}) &= dx_1 \frac{\partial \phi}{\partial x_1} + dx_2 \frac{\partial \phi}{\partial x_2} + dx_3 \frac{\partial \phi}{\partial x_3} \\ &= dx_i \frac{\partial \phi}{\partial x^i} \\ &= d\mathbf{x} \cdot (\nabla \phi),\end{aligned}$$

- Gradient operator,  $\nabla$ ,

$$\begin{aligned}\nabla &\equiv \hat{\mathbf{e}}_k \partial_k \\ &= \hat{\mathbf{e}}_1 \frac{\partial}{\partial x_1} + \hat{\mathbf{e}}_2 \frac{\partial}{\partial x_2} + \hat{\mathbf{e}}_3 \frac{\partial}{\partial x_3}.\end{aligned}$$

# Tensor fields: Divergence and Curl

# Tensor fields: Divergence and Curl

- ▶ Since,  $\nabla$  acts like a vector, we can construct a scalar by taking the inner product, with a vector field,  $\mathbf{A}(\mathbf{x})$ ,

$$\nabla \cdot \mathbf{A}(\mathbf{x}) = \partial_k A_k(x).$$

# Tensor fields: Divergence and Curl

- ▶ Since,  $\nabla$  acts like a vector, we can construct a scalar by taking the inner product, with a vector field,  $\mathbf{A}(\mathbf{x})$ ,

$$\nabla \cdot \mathbf{A}(\mathbf{x}) = \partial_k A_k(x).$$

- ▶ Call it **Divergence of the vector field**,

$$\nabla \cdot \mathbf{A} = \frac{\partial A_1}{\partial x_1} + \frac{\partial A_2}{\partial x_2} + \frac{\partial A_3}{\partial x_3}$$



# Tensor fields: Divergence and Curl

- ▶ Since,  $\nabla$  acts like a vector, we can construct a scalar by taking the inner product, with a vector field,  $\mathbf{A}(\mathbf{x})$ ,

$$\nabla \cdot \mathbf{A}(\mathbf{x}) = \partial_k A_k(x).$$

- ▶ Call it **Divergence of the vector field**,

$$\nabla \cdot \mathbf{A} = \frac{\partial A_1}{\partial x_1} + \frac{\partial A_2}{\partial x_2} + \frac{\partial A_3}{\partial x_3}$$

- ▶ We can further create a vector by taking the cross product of  $\nabla$  and  $\mathbf{A}(\mathbf{x})$ ,

$$\nabla \times \mathbf{A} = (\nabla \times \mathbf{A})_k \hat{\mathbf{e}}_k,$$

$$(\nabla \times \mathbf{A})_k = \epsilon_{ijk} \partial_i A_j$$

# Tensor fields: Divergence and Curl

- ▶ Since,  $\nabla$  acts like a vector, we can construct a scalar by taking the inner product, with a vector field,  $\mathbf{A}(\mathbf{x})$ ,

$$\nabla \cdot \mathbf{A}(\mathbf{x}) = \partial_k A_k(x).$$

- ▶ Call it **Divergence of the vector field**,

$$\nabla \cdot \mathbf{A} = \frac{\partial A_1}{\partial x_1} + \frac{\partial A_2}{\partial x_2} + \frac{\partial A_3}{\partial x_3}$$

- ▶ We can further create a vector by taking the cross product of  $\nabla$  and  $\mathbf{A}(\mathbf{x})$ ,

$$\nabla \times \mathbf{A} = (\nabla \times \mathbf{A})_k \hat{\mathbf{e}}_k,$$

$$(\nabla \times \mathbf{A})_k = \epsilon_{ijk} \partial_i A_j$$

- ▶ call it **Curl of the vector field**

# Tensor fields: Divergence and Curl

# Tensor fields: Divergence and Curl



$$\nabla r^n = n r^{n-2} \mathbf{x}.$$

# Tensor fields: Divergence and Curl



$$\nabla r^n = n r^{n-2} \mathbf{x}.$$



$$\nabla \cdot \mathbf{x} = \frac{\partial x_1}{\partial x_1} + \frac{\partial x_2}{\partial x_2} + \frac{\partial x_3}{\partial x_3} = 3$$

# Tensor fields: Divergence and Curl



$$\nabla r^n = n r^{n-2} \mathbf{x}.$$



$$\nabla \cdot \mathbf{x} = \frac{\partial x_1}{\partial x_1} + \frac{\partial x_2}{\partial x_2} + \frac{\partial x_3}{\partial x_3} = 3$$



$$(\nabla \times \mathbf{x})_k = \epsilon_{ijk} \partial_i x_j = \epsilon_{ijk} \delta_{ij} = 0.$$

# Tensor fields: Divergence and Curl



$$\nabla r^n = n r^{n-2} \mathbf{x}.$$



$$\nabla \cdot \mathbf{x} = \frac{\partial x_1}{\partial x_1} + \frac{\partial x_2}{\partial x_2} + \frac{\partial x_3}{\partial x_3} = 3$$



$$(\nabla \times \mathbf{x})_k = \epsilon_{ijk} \partial_i x_j = \epsilon_{ijk} \delta_{ij} = 0.$$

- ▶ Can define a double derivative thru the inner product, the **Laplacian**

$$\nabla \cdot \nabla = \frac{\partial^2}{(\partial x_1)^2} + \frac{\partial^2}{(\partial x_2)^2} + \frac{\partial^2}{(\partial x_3)^2}.$$

# Tensor fields: Integration



# Tensor fields: Integration

- **Gauss Divergence theorem:** If  $S$  is a closed surface enclosing a volume,  $V$

$$\iiint_V d^3\mathbf{x} \nabla \cdot \mathbf{A} = \oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{A},$$

$\hat{\mathbf{n}}$  is the unit outward normal vector on the surface  $S$ .

# Tensor fields: Integration

- ▶ **Gauss Divergence theorem:** If  $S$  is a closed surface enclosing a volume,  $V$

$$\iiint_V d^3\mathbf{x} \nabla \cdot \mathbf{A} = \oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{A},$$

$\hat{\mathbf{n}}$  is the unit outward normal vector on the surface  $S$ .

- ▶ **Stokes Curl theorem:** If  $S$  is an open surface, with a boundary,  $C$  (closed curve)

$$\iint_S dS \hat{\mathbf{n}} \cdot (\nabla \times \mathbf{A}) = \oint_C d\mathbf{l} \cdot \mathbf{A}$$

# Tensor fields: Integration

- ▶ **Gauss Divergence theorem:** If  $S$  is a closed surface enclosing a volume,  $V$

$$\iiint_V d^3\mathbf{x} \nabla \cdot \mathbf{A} = \oiint_S dS \hat{\mathbf{n}} \cdot \mathbf{A},$$

$\hat{\mathbf{n}}$  is the unit outward normal vector on the surface  $S$ .

- ▶ **Stokes Curl theorem:** If  $S$  is an open surface, with a boundary,  $C$  (closed curve)

$$\iint_S dS \hat{\mathbf{n}} \cdot (\nabla \times \mathbf{A}) = \oint_C d\mathbf{l} \cdot \mathbf{A}$$

- ▶ Should be thought of as vector generalizations of *Fundamental theorem* of single variable calculus:

$$\int_a^b dx \frac{df(x)}{dx} = f(b) - f(a)$$

# Tensor fields: Integration





Figure: Pictorial representation of Gauss and Stokes Theorems.