RAJALAKSHMI ENGINEERING COLLEGE

Mahesh G K 250701406-2025-CSE

M2

| | |
|---|---|
| **Started on** | Wednesday, 16 September 2026, 9:19 AM |
| **State** | Finished |
| **Completed on** | Wednesday, 16 September 2026, 9:23 AM |
| **Time taken** | 3 mins 10 secs |
| **Marks** | 1.00/1.00 |
| **Grade** | **10.00** out of 10.00 (**100**%) |

**Question 1** | Correct Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that A[j] - A[i] = k, i != j.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as 5 - 1 = 4

So Return 1.

**For example:**

| Input | Result |
|---|---|
| 3<br>1 3 5<br>4 | 1 |

**Answer:** (penalty regime: 0 %)

```
#include <stdio.h>

int main() {
    int n, k;
    scanf("%d", &n);

    int a[n];

    for (int i = 0; i < n; i++) {
        scanf("%d", &a[i]);
    }

    scanf("%d", &k);

    for (int i = 0; i < n; i++) {
        for (int j = i + 1; j < n; j++) {

            if (a[j] - a[i] == k) {
                printf("1");
                return 0;
            }

            if (a[j] - a[i] > k) {
                break;
            }
        }
    }

    printf("0");

    return 0;
}
```

| | Input | Expected | Got | |
|---|---|---|---|---|
| ✔ | 3<br>1 3 5<br>4 | 1 | 1 | ✔ |
| ✔ | 10<br>1 4 6 8 12 14 15 20 21 25<br>1 | 1 | 1 | ✔ |
| ✔ | 10<br>1 2 3 5 11 14 16 24 28 29<br>0 | 0 | 0 | ✔ |
| ✔ | 10<br>0 2 3 7 13 14 15 20 24 25<br>10 | 1 | 1 | ✔ |

Passed all tests! ✔

Correct

Marks for this submission: 1.00/1.00.

Back to Course