MATHHUB_PRO // NCERT_SOLUTIONS

Answers to Some Questions in Exercises

UNIT 1

1.4 16.23 M

1.5 0.617 m, 0.01 and 0.99, 0.67

1.6 157.8 mL

1.7 33.5%

1.8 17.95 m and 9.10 M

1.9 1.5×10^{-3}%, 1.25×10^{-4} m

1.15 40.907 g mol^{-1}

1.16 73.58 *k*Pa

1.17 12.08 *k*Pa

1.18 10 g

1.19 23 g mol^{-1}, 3.53 kPa

1.20 269.07 *K*

1.21 A = 25.58 u and B = 42.64 u

1.22 0.061 M

1.24 KCl, CH_3OH, CH_3CN, Cyclohexane

1.25 Toluene, chloroform; Phenol, Pentanol; Formic acid, ethylelne glycol

1.26 5 m

1.27 $2.45\text{x}10^{-8}$ M

1.28 1.424%

1.29 3.2 g of water

1.30 4.575 g

1.32 0.65^0

1.33 i = 1.0753, $K_a = 3.07\times10^{-3}$

1.34 17.44 mm Hg

1.35 178×10^{-5}

1.36 280.7 torr, 32 torr

1.38 0.6 and 0.4

1.39 $x\,(O_2)$ $4.6\text{x}10^{-5}$, $x\,(N_2)$ 9.22×10^{-5}

1.40 0.03 mol of $CaCl_2$

1.41 $5.27\text{x}10^{-3}$ atm.

UNIT 2

2.4 (i) E^o = 0.34V, $\Delta_r G^o$ = – 196.86 kJ mol^{-1}, $K = 3.124 \times 10^{34}$
(ii) E^o = 0.03V, $\Delta_r G^o$ = – 2.895 kJ mol^{-1}, K = 3.2

2.5 (i) 2.68 V, (ii) 0.53 V, (iii) 0.08 V, (iv) –1.298 V

2.6 1.56 V

2.8 124.0 S cm^2 mol^{-1}

2.9 0.219 cm^{-1}

2.11 1.85×10^{-5}

2.12 3F, 2F, 5F

2.13 1F, 4.44F

2.14 2F, 1F

2.15 1.8258g

2.16 14.40 min, Copper 0.427g, Zinc 0.437 g

UNIT 3

3.2 (i) 8.0×10^{-9} mol L^{-1} s^{-1}; 3.89×10^{-9} mol L^{-1} s^{-1}

3.4 $bar^{-1/2}s^{-1}$

3.6 (i) 4 times (ii) ¼ times

3.8 (i) 4.67×10^{-3} mol $L^{-1}s^{-1}$ (ii) 1.98×10^{-2} s^{-1}

3.9 (i) rate = $k[A][B]^2$ (ii) 9 times

© NCERT not to be republished

3.10 Orders with respect to A is 1.5 and order with respect to B is zero.

3.11 rate law = $k[A][B]^2$; rate constant = 6.0 $M^{-2}min^{-1}$

3.13 (i) 3.47 x 10^{-3} seconds (ii) 0.35 minutes (iii) 0.173 years

3.14 1845 years

3.16 4.6×10^{-2} s

3.17 0.7814 μg and 0.227 μg.

3.19 77.7 minutes

3.20 2.20×10^{-3} s^{-1}

3.21 2.23×10^{-3} s^{-1}, 7.8×10^{-4} atm s^{-1}

3.23 3.9×10^{12} s^{-1}

3.24 0.135 M

3.25 0.158 M

3.26 232.79 kJ mol^{-1}

3.27 239.339 kJ mol^{-1}

3.28 24°C

3.29 E_a = 76.750 kJ mol^{-1}, $k = 0.9965 \times 10^{-2}$ s^{-1}

3.30 52.8 kJ mol^{-1}

UNIT 4

4.2 It is because Mn^{2+} has $3d^5$ configuration which has extra stability.

4.5 Stable oxidation states.

$3d^3$ (Vanadium): (+2), +3, +4, and +5

$3d^5$ (Chromium): +3, +4, +6

$3d^5$ (Manganese): +2, +4, +6, +7

$3d^8$ (Nickel): +2, +3 (in complexes)

$3d^4$ There is no d^4 configuration in the ground state.

4.6 Vanadate VO_3^-, chromate CrO_4^{2-}, permanganate MnO_4^-

4.10 +3 is the common oxidation state of the lanthanoids

In addition to +3, oxidation states +2 and +4 are also exhibited by some of the lanthanoids.

4.13 In transition elements the oxidation states vary from +1 to any highest oxidation state by one

For example, for manganese it may vary as +2, +3, +4, +5, +6, +7. In the nontransition elements the variation is selective, always differing by 2, e.g. +2, +4, or +3, +5 or +4, +6 etc.

4.18 Except Sc^{3+}, all others will be coloured in aqueous solution because of incompletely filled 3*d*-orbitals, will give rise to *d*-*d* transitions.

4.21 (i) Cr^{2+} is reducing as it involves change from d^4 to d^3, the latter is more stable configuration (t_{2g}^3) Mn(III) to Mn(II) is from $3d^4$ to $3d^5$ again $3d^5$ is an extra stable configuration.

(ii) Due to CFSE, which more than compensates the 3rd IE.

(iii) The hydration or lattice energy more than compensates the ionisation enthalpy involved in removing electron from d^1.

4.23 Copper, because with +1 oxidation state an extra stable configuration, $3d^{10}$ results.

4.24 Unpaired electrons Mn^{3+} = 4, Cr^{3+} = 3, V^{3+} = 2, Ti^{3+} = 1. Most stable Cr^{3+}

4.28 Second part 59, 95, 102.

4.30 Lawrencium, 103, +3

4.36 Ti^{2+} = 2, V^{2+} = 3, Cr^{3+} = 3, Mn^{2+} = 5, Fe^{2+} = 6, Fe^{3+} = 5, CO^{2+} = 7, Ni^{2+} = 8, Cu^{2+} = 9

4.38 $M\sqrt{n(n+2)}$ = 2.2, n ≈ 1, d^2 sp^3, CN^- strong ligand

= 5.3, n ≈ 4, sp^3, d^2, H_2O weak ligand

= 5.9, n ≈ 5, sp^3, Cl^- weak ligand.

© NCERT not to be republished

Reprint 2025-26

UNIT 5

5.5 (i) + 3 (ii) +3 (iii) +2 (iv) +3 (v) +3

5.6 (i) $[Zn(OH)_4]^{2-}$ (ii) $K_2[PdCl_4]$ (iii) $[Pt(NH_3)_2Cl_2]$ (iv) $K_2[Ni(CN)_4]$
(v) $[Co(NH_3)_5(ONO)]^{2+}$ (vi) $[Co(NH_3)_6]_2(SO_4)_3$ (vii) $K_3[Cr(C_2O_4)_3]$ (viii) $[Pt(NH_3)_6]^{4+}$
(ix) $[CuBr_4]^{2-}$ (x) $[Co(NH_3)_5(NO_2)]^{2+}$

5.9 (i) $[Cr(C_2O_4)_3]^{3-}$ – Nil
(ii) $[Co(NH_3)_3Cl_3]$ – Two (*fac*- and *mer*-)

5.12 Three (two *cis* and one *trans*)

5.13 Aqueous $CuSO_4$ solution exists as $[Cu(H_2O)_4]SO_4$ which has blue colour due to $[Cu(H_2O)_4]^{2+}$ ions.
(i) When KF is added, the weak H_2O ligands are replaced by F^- ligands, forming $[CuF_4]^{2-}$ ions which is a green precipitate.
$[Cu(H_2O)_4]^{2+} + 4F^- \rightarrow [CuF_4]^{2-} + 4H_2O$
(ii) When KCl is added, Cl^- ligands replace the weak H_2O ligands forming $[CuCl_4]^{2-}$ ions which has bright green colour.
$[Cu(H_2O)_4]^{2+} + 4Cl^- \rightarrow [CuCl_4]^{2-} + 4H_2O$

5.14 $[Cu(H_2O)_4]^{2+} + 4\ CN^- \rightarrow [Cu(CN)_4]^{2-} + 4H_2O$
As CN^- is a strong ligand, it forms a highly stable complex with Cu^{2+} ion. On passing H_2S, free Cu^{2+} ions are not available to form the precipitate of CuS.

5.23 (i) OS = +3, CN = 6, d-orbital occupation is $t_{2g}^6\ e_g^0$,
(ii) OS = +3, CN = 6, d^3 (t_{2g}^3),
(iii) OS = +2, CN = 4, d^7 ($t_{2g}^5\ e_g^2$),
(iv) OS = +2, CN = 6, d^5 ($t_{2g}^3\ e_g^2$).

5.28 (iii)

5.29 (ii)

5.30 (iii)

5.31 (iii)

5.32 (i) The order of the ligand in the spectrochemical series :
$H_2O < NH_3 < NO_2^-$
Hence the energy of the observed light will be in the order :
$[Ni(H_2O)_6]^{2+} < [Ni(NH_3)_6]^{2+} < [Ni(NO_2)_6]^{4-}$
Thus, wavelengths absorbed ($E = hc/\lambda$) will be in the opposite order.

© NCERT not to be republished

Reprint 2025-26

NOTES

© NCERT
be republished

NOTES

© NCERT
be republished

Answers to Some Questions in Exercises

UNIT 7

7.1 (i) 2,2,4-Trimethylpentan –3-ol (ii) 5-Ethylheptane –2, 4-diol
(iii) Butane –2,3-diol (iv) Propane –1,2,3,-triol
(v) 2- Methylphenol (vi) 4-Methylphenol
(vii) 2,5 – Dimethylphenol (viii) 2,6-Dimethylphenol
(ix) 1-Methoxy-2-methylpropane (x) Ethoxybenzene
(xi) 1-phenoxyheptane (xii) 2 –Ethoxybutane

7.2 (i) 

(ii) 

(iii) 

(iv) 

(v) $C_2H_5 - O - CH_2 - CH_2 - CH_3$

(vi)

(vii) 

(viii) 

(ix) 

(x) 

7.3 (i) (a) $CH_3CH_2CH_2CH_2CH_2OH$, Pentan-1-ol;

(b)  , 2-Methylbutan-1-ol;

(c)  , 2,2-Dimethylpropan-1-ol

(d)  , Pentan-3-ol

(e)  , Pentan-3-ol

(f)  , 3-Methylbutan-2-ol

(g)  , 2-Methylbutan-2-ol

(h)  ,
3-Methylbutan-1-ol

7.4 Hydrogen bonding in propanol.

© NCERT not to be republished

7.5 Hydrogen bonding between alcohol and water molecules.

7.8 o-Nitrophenol is steam volatile because of intramolecular hydrogen bonding.

7.12 **Hint:** Carryout sulphonation followed by nucleophilic substitution.

7.13 (i)



(ii)



(iii) $CH_3(CH_2)_4Cl + NaOH \longrightarrow CH_3(CH_2)_4OH + NaCl$

7.14 Reaction with (i) sodium and (ii) sodium hydroxide

7.15 Due to electron withdrawing effect of nitro group and electron releasing effect of methoxy group.

7.20 (i) Hydration of Propene.

(ii) By nucleophilic substitution of –Cl in benzyl chloride using dilute NaOH.

(iii) $C_2H_5MgBr + HCHO \rightarrow C_2H_5CH_2OMgBr \xrightarrow{H_2O} C_2H_5CH_2OH$

(iv)



7.23 (i) 1-Ethoxy-2-methylpropane.

(ii) 2-Chloro-1-methoxyethane.

(iii) 4-Nitroanisole.

(iv) 1-Methoxypropane.

(v) 1-Ethoxy-4,4-dimethylcyclohexane.

(vi) Ethoxybenzene.

UNIT 8

8.2 (i) 4-Methylpentanal

(ii) 6-Chloro-4-ethylhexan-3-one

(iii) But-2-enal

(iv) Pentane-2,4-dione

(v) 3,3,5-Trimethylhexan-2-one

(vi) 3,3-Dimethylbutanoic acid

(vii) Benzene –1,4-dicarbaldehyde

8.3 (i)



(ii)



(iii)



(iv)



(v)



(vi)



Reprint 2025-26

(vii) 

(viii) 

8.4 (i) Heptan-2-one (ii) 4-Bromo-2-methylhexanal (iii) Heptanal
(iv) 3-Phenylprop-2-enal (v) Cyclopentanecarbaldehyde (vi) Diphenylmethanone

8.5 (i) 

(ii) 

(iii) 

(iv) 

(v) 

(vi) 

8.6 (i) 

(ii) 

(iii) 

(iv) 

(v) 

8.7 (ii), (v), (vi), (vii): Aldol condensation. (i), (iii), (ix) Cannizaro reaction. (iv), (viii) Neither.

8.10 2-Ethylbenzaldehyde (draw the structure yourself).

8.11 (A) $CH_3CH_2CH_2COOCH_2CH_2CH_2CH_3$, butyl butanoate.
(B) $CH_3CH_2CH_2$ COOH (C) $CH_3CH_2CH_2CH_2OH$. Write equation yourself.

8.12 (i) Di-tert-butyl ketone < Methyl tert-butyl ketone < Acetone < Acetaldehyde
(ii) $(CH_3)_2CHCOOH$ < $CH_3CH_2CH_2COOH$ < $CH_3CH(Br)CH_2COOH$ < $CH_3CH_2CH(Br)COOH$
(iii) 4-Methoxybenzoic acid < Benzoic acid < 4-Nitrobenzoic acid < 3,4-Dinitrobenzoic acid.

8.17 (i) 

(ii) 

(iii) 

(iv) $C_6H_5COCl–AlCl_3$

(v) 

(vi) 

(vii)  + other products

(viii) $CH_3CH(OH)CH_2COOC_2H_5$

(ix) 

(x) 1. BH_3; 2. $H_2O_2/\bar{O}H$; 3. PCC

(xi) 

8.19 The compound is methyl ketone and its structure would be: $CH_3COCH_2CH_2CH_3$

© NCERT not republished

Reprint 2025-26

UNIT 9

9.1 (i) 1-methylethylamine or propan-2-amine (ii) Propan-1-amine
(iii) N-methyl-2-methylethylamine or N-methylpropan-2-amine (iv) 2-methylpropan-2-amine
(v) N-methylbenzenamine or N-methylaniline (vi) N-Ethyl-N-methylethanamine
(vii) 3-Bromoaniline or 3-Bromobenzenamine

9.4 (i) $C_6H_5NH_2 < C_6H_5NHCH_3 < C_2H_5NH_2 < (C_2H_5)_2NH$
(ii) $C_6H_5NH_2 < C_6H_5N(CH_3)_2 < CH_3NH_2 < (C_2H_5)_2NH$
(iii) (a) p-nitroaniline < aniline < p-toluidine
(b) $C_6H_5NH_2 < C_6H_5NHCH_3 < C_6H_5CH_2NH_2$
(iv) $(C_2H_5)_3N > (C_2H_5)_2NH > C_2H_5NH_2 > NH_3$ (v) $(CH_3)_2NH < C_2H_5NH_2 < C_2H_5OH$
(vi) $C_6H_5NH_2 < (C_2H_5)_2NH < C_2H_5NH_2$

© NCERT
be republished

Reprint 2025-26

Notes

© NCERT
be republished

Notes

© NCERT
be republished