

Reflexivity in Options Markets: A Stochastic-Volatility Model with Dealer-Gamma Feedback, Hopf Bifurcation Calculus, and a Pre-Registered Evaluation Framework

Mahimn Patel
Independent Researcher
`mahimn.patel.k@gmail.com`

April 2026

Abstract

We study a continuous-time Heston-like stochastic-volatility model in which the underlying spot price is coupled to its own option market through a dealer-gamma feedback channel, derived from the open-interest grid via a Gârleanu–Pedersen–Potesman (Gârleanu et al., 2009) demand-pressure mapping. The model state is three-dimensional—spot S_t , instantaneous variance v_t , and a low-pass-filtered log-spot memory variable z_t —and reduces to standard time-dependent Heston (Heston, 1993) at zero coupling. Our contribution is the synthesis of three previously separate ingredients into a single analytical and computational object: (i) a continuous-time SV backbone with explicit dealer-gamma drift, (ii) a formal Hopf bifurcation calculus on the deterministic skeleton with first Lyapunov coefficient ℓ_1 and Khasminskii-style stochastic shift Λ , and (iii) a pre-registered evaluation protocol for RL-trained hedging agents trained inside the simulator. For a representative dimensionless calibration we obtain a critical coupling $\kappa^* \approx 0.8964$, angular frequency $\omega^* \approx 0.5724$ rad/yr, and first Lyapunov coefficient $\ell_1 \approx -0.025 < 0$, so the bifurcation is supercritical: an attracting limit cycle in volatility is born for $\kappa > \kappa^*$. The 2D reduction (without z_t) has an upper-triangular Jacobian and admits only a saddle-node onset, recovering the discrete-time recursion of Dai (2025); the 3D extension is what unlocks the Hopf. The closest model-structure precedent (Halperin and Itkin, 2025) does not perform the bifurcation analysis we provide. For the natural parametric specification with log-normal open-interest in moneyness, we obtain a closed-form Hopf threshold and a closed-form ℓ_1 , exposing a parametric phase boundary in (σ_q, γ) space.

1 Introduction

Real options markets exhibit endogenous volatility cycles that pure stochastic-volatility (SV) models cannot reproduce. The empirical anchor is Hardiman et al. (2013), who fit Hawkes-process branching ratios to E-mini mid-price changes over 1998–2011 and find $n \approx 1$ throughout—the precise critical edge between sub- and super-criticality of endogenous events. Soros (2009) and Bouchaud (2011) are the conceptual antecedents; Filimonov and Sornette (2012) formalised the reflexivity-quantification programme that Hardiman et al. (2013) executed. The question we take up here is what continuous-time mechanism produces this critical reflexivity, and—given a candidate mechanism—how the analytical structure that quantifies it should be evaluated against historical surface dynamics.

We contribute the following.

- A 3D reflexive SV simulator (S_t, v_t, z_t) with explicit dealer-gamma feedback derived from the option open-interest grid via the Gârleanu et al. (2009) demand-pressure mapping. The variance backbone is Heston (Heston, 1993); the memory equation $\dot{z} = -\alpha z + \beta \log(S/S_0)$ closes the leverage feedback channel that the 2D reduction lacks.
- A Hopf bifurcation theorem (Theorem 1) for the deterministic skeleton at κ^* given by Liu’s criterion (Routh–Hurwitz with one zero), with closed-form first Lyapunov coefficient ℓ_1 and closed-form Hopf threshold for log-normal open-interest in moneyness (Eqs. 14–15).
- A numerical phase diagram in (ξ, ρ, σ_v) together with the Khasminskii-style stochastic-Hopf shift Λ and a parametric phase boundary in (σ_q, γ) space.
- A pre-registration of the empirical evaluation pipeline (sliced-Wasserstein-2 over 21-day surface windows, κ -sensitivity slope, Hopf-frequency spectral test) at git commit 268c061 with an OpenTimestamps Bitcoin-anchored proof, in the spirit of Camerer et al. (2018) adapted for computational finance.

Closest precedents. Halperin and Itkin (2025) (Marketrn) share the model-structural axis—continuous-time SDE, memory channel, options-aware—but do not perform the bifurcation analysis. Dai (2025) share the dealer-gamma axis but produce a saddle-node onset (the 2D reduction’s instability), not a Hopf. He et al. (2025) (NeurIPS 2025) share the RL-with-environmental-sweep idea but train separate agents per grid point rather than reporting the slope at an anchor. The continuous-time HAM-Hopf precedents (He et al., 2009; Chiarella et al., 2006) use delay-differential heterogeneous-belief models without an options-grid feedback channel; the discrete-time Brock et al. (2009) “more hedging instruments may destabilize markets” result is the methodological template for derivative-driven Neimark–Sacker (Hopf-for-maps) bifurcations. The hedger-flow-feedback foundations are Frey and Stremme (1997); Sircar and Papanicolaou (1998); Platen and Schweizer (1998). Sections 3, 5 and 6 make these contrasts precise.

Roadmap. Section 2 states the model. Section 3 proves Theorem 1 and gives the closed-form ℓ_1 for log-normal OI. Section 4 reports the numerical phase diagram. Section 5 describes the pre-registered evaluation framework. Section 6 reports the mechanism decomposition against Marketrn. Section 7 closes with the empirical-calibration roadmap. Code at <https://github.com/mahimn01/reflexive-options>; pre-registration anchored to commit 268c061.

2 The reflexive model

Let S_t be the underlying spot price, v_t the instantaneous variance, and z_t a memory variable encoding low-pass-filtered price history. The reflexive system is

$$\frac{dS_t}{S_t} = (\mu + \kappa G(S_t, z_t, v_t)) dt + \sigma(S_t, v_t) dW_t^S, \quad (1)$$

$$dv_t = (\kappa_v(\theta_v - v_t) + \gamma z_t) dt + \xi \sqrt{v_t} dW_t^v, \quad (2)$$

$$dz_t = (-\alpha z_t + \beta \log(S_t/S_0)) dt, \quad (3)$$

$$d\langle W^S, W^v \rangle_t = \rho dt, \quad (4)$$

with $\kappa \geq 0$ the reflexive coupling, $\gamma \geq 0$ the leverage-feedback strength, $\alpha > 0$ the memory decay rate, β dimensionless, and $G(\cdot)$ the aggregate dealer-gamma exposure derived from the option

open-interest grid:

$$G(S, t) = \sum_{K, T} q_{K, T} \Gamma_{K, T}(S, t) \text{sign}(K, T), \quad (5)$$

where $q_{K, T}$ is open interest at strike K and maturity T , $\Gamma_{K, T}$ is per-contract Black–Scholes gamma, and the sign convention follows the SqueezeMetrics dealer-net convention (SqueezeMetrics, 2017). When $\kappa = \gamma = 0$, the system collapses to standard Heston (Heston, 1993); this collapse is verified numerically in `tests/test_simulator.py`.

For numerical stability of the variance scheme we use the full-truncation Euler of Lord et al. (2010); the empirical reference for the dealer-gamma magnitudes in G is Barbon and Buraschi (2021); Dim et al. (2024) (calibration to SPX 0DTE / dealer-gamma data, which we will use at Phase 4 of the empirical pipeline).

Why three states. A reduced 2D specification—dropping z_t and the γz term—has Jacobian in deviation variables (y, u) around an equilibrium of the form

$$J_{2D}(\kappa) = \begin{pmatrix} a(\kappa) & b(\kappa) \\ 0 & -\kappa_v \end{pmatrix},$$

which is upper triangular: eigenvalues $\{a(\kappa), -\kappa_v\}$ are always real, and only saddle-node / transcritical bifurcations are admissible as $a(\kappa)$ crosses zero. This recovers the static onset of Dai (2025). To Hopf, the feedback must flow in both directions: price $\rightarrow$ variance and variance $\rightarrow$ price. The minimal modification is the leverage channel γz together with the auxiliary memory equation (3)—structurally analogous to the memory variable in the Marketron quasi-particle SDE (Halperin and Itkin, 2025) though with very different mechanistic content.

3 Hopf bifurcation theorem and closed-form ℓ_1

3.1 Equilibria and Jacobian

Drop the Brownian terms and work in log-deviation variables $(y, u, z) = (\log(S/S^*), v - \theta_v, z - z^*)$ around the unique equilibrium (S^*, θ_v, z^*) defined by

$$\mu - \frac{1}{2}\sigma^2(S^*, \theta_v) + \kappa G(S^*, z^*, \theta_v) = 0, \quad z^* = \frac{\beta}{\alpha} \log(S^*/S_0). \quad (6)$$

To linear order, the deterministic skeleton (1–3) becomes $\dot{\mathbf{x}} = J(\kappa) \mathbf{x}$ with $\mathbf{x} = (y, u, z)^\top$ and

$$J(\kappa) = \begin{pmatrix} a(\kappa) & b(\kappa) & \kappa G_z \\ 0 & -\kappa_v & \gamma \\ \beta & 0 & -\alpha \end{pmatrix}, \quad (7)$$

where $a(\kappa) = \kappa G_x - \frac{1}{2}\partial_x \sigma^2$, $b(\kappa) = \kappa G_v - \frac{1}{2}\partial_v \sigma^2$, and G_x, G_v, G_z denote partial derivatives of G at the equilibrium.

3.2 Routh–Hurwitz / Liu’s criterion

The characteristic polynomial of (7) is $P(\lambda; \kappa) = \lambda^3 + c_2(\kappa)\lambda^2 + c_1(\kappa)\lambda + c_0(\kappa)$, with

$$c_2 = -a + \kappa_v + \alpha, \quad c_1 = -a\kappa_v - a\alpha + \kappa_v\alpha - \kappa G_z \beta, \quad c_0 = -a\kappa_v\alpha - \beta(\kappa G_z \kappa_v + b\gamma).$$

A 3D system Hopfs iff (Liu’s criterion, equivalent to Routh–Hurwitz with one zero):

$$c_2 > 0, \quad c_0 > 0, \quad c_1 c_2 - c_0 = 0, \quad \frac{d}{d\kappa} [c_1 c_2 - c_0] \neq 0. \quad (8)$$

Define $H(\kappa) := c_1(\kappa) c_2(\kappa) - c_0(\kappa)$. The Hopf threshold is

$$\boxed{\kappa^* = \{\kappa > 0 : H(\kappa) = 0, H'(\kappa) \neq 0, c_2(\kappa) > 0, c_0(\kappa) > 0\}.} \quad (9)$$

At $\kappa = \kappa^*$ the Jacobian has a pure imaginary eigenvalue pair $\lambda_{\pm} = \pm i\omega^*$ with $\omega^* = \sqrt{c_1(\kappa^*)}$, plus a real eigenvalue $\lambda_3 = -c_2(\kappa^*) < 0$.

3.3 Theorem 1 and proof sketch

Theorem 1 (Hopf bifurcation in the gamma-coupled SV skeleton). *Let the deterministic system (1–3) (with Brownian terms removed) satisfy:*

- (A1) $G \in C^3$ and $\sigma^2 \in C^3$ in their arguments, with σ^2 bounded below by a positive constant.
- (A2) There exists a locally unique equilibrium (S^*, θ_v, z^*) given by (6) for every κ in some interval $[0, \kappa_{\max}]$, in some open neighborhood of which the linearization (7) is non-degenerate.
- (A3) The map $\kappa \mapsto (a(\kappa), G_z, G_v)$ is C^3 .
- (A4) There exists $\kappa^* \in (0, \kappa_{\max})$ satisfying (9) with $\omega^* := \sqrt{c_1(\kappa^*)} > 0$.
- (A5) The first Lyapunov coefficient $\ell_1(\kappa^*) \neq 0$ (Kuznetsov, 2004, eq. 3.20).

Then the equilibrium undergoes a Hopf bifurcation at $\kappa = \kappa^*$. Specifically:

1. For κ in a one-sided neighbourhood of κ^* (sub- or super-critical depending on $\text{sgn } \ell_1$), there exists a unique smooth family of periodic orbits Γ_{κ} of period $T_{\kappa} = 2\pi/\omega^* + O(\kappa - \kappa^*)$ and amplitude proportional to $\sqrt{|\kappa - \kappa^*|}$.
2. The orbits are stable iff $\ell_1(\kappa^*) < 0$ (supercritical).
3. On Γ_{κ} , $u_t = v_t - \theta_v$ oscillates with amplitude proportional to $\sqrt{|\kappa - \kappa^*|}$, providing endogenous volatility cycles.

Proof sketch. The eigenvalue pair $\lambda_{\pm}(\kappa)$ is smooth in κ near κ^* by the implicit function theorem applied to $P(\lambda; \kappa) = 0$. Routh–Hurwitz gives $\text{Re } \lambda_{\pm}(\kappa^*) = 0$, and the transversality $H'(\kappa^*) \neq 0$ translates to $\frac{d}{d\kappa} \text{Re } \lambda_{\pm} \neq 0$. The centre manifold theorem (Kuznetsov, 2004, Thm. 5.4) (see also Carr, 1981) gives a smooth 2D invariant manifold tangent to the eigenspace of $\pm i\omega^*$. Conjugating the restricted system to its Poincaré normal form and applying Kuznetsov (2004, Thm. 3.3) yields claims (1)–(3). $\square$

3.4 Numerical anchor

The default `BifurcationConfig` in our experiments code does not exhibit a Hopf within the literature-prior κ range when the dealer-gamma magnitudes are tuned to empirical scale—this is consistent with the empirical observation that real options markets sit *near* but not *across* κ^* , in line with the $n \approx 1$ regime of Hardiman et al. (2013). For a worked numerical example we use a representative dimensionless regime with $\{G_x, G_v, G_z\} = \{0.5, -0.5, -0.5\}$, $\alpha = 0.5/\text{yr}$, $\beta = 1$,

Quantity	Value
κ^*	0.8964
ω^*	0.5724 rad/yr
Third real eigenvalue λ_3	-2.052
First Lyapunov coefficient ℓ_1	-2.53×10^{-2}
Bifurcation type	supercritical ($\ell_1 < 0$)

Table 1: Canonical Hopf-exhibiting regime for the deterministic skeleton.

$\gamma = 0.5$, $\kappa_v = 2/\text{yr}$, and constant-volatility surrogate ($\partial_v \sigma^2 = 0$). The deterministic skeleton then satisfies:

Supercriticality means an attracting limit cycle is born for $\kappa > \kappa^*$ with amplitude proportional to $\sqrt{\kappa - \kappa^*}$ —i.e. endogenous volatility cycles are stable observables of the model rather than transients. For the stochastic lift at $(\xi, \rho) = (0.3, -0.7)$ (calibration-representative for SPX), the Khasminskii sphere-process estimator (Khasminskii, 1980; Arnold, 1998; Baxendale, 1994) gives $\Lambda(\kappa^*) \approx +1.85 \times 10^{-2}$. The positive sign means noise destabilises the equilibrium; the stochastic Hopf threshold $\kappa_{\text{stoch}}^*(\varepsilon) \approx \kappa^* - \varepsilon^2 \Lambda / |\alpha'(\kappa^*)|$ lies below the deterministic threshold, consistent with the shear-induced corrections of Engel et al. (2024).

3.5 Closed-form κ^* and ℓ_1 for log-normal OI in moneyness

For the natural parametric specification of dealer gamma—a log-normal open-interest density in log-strike—the entire computation admits a closed form, eliminating the numerical-noise vulnerability of the FD-tensor pipeline and exposing the parametric phase boundary explicitly.

Replace the discrete OI grid with the continuous density $q(\log K) = \frac{1}{\sigma_q \sqrt{2\pi}} \exp(-(\log K - \mu_q)^2 / (2\sigma_q^2))$. Take a single representative maturity T_{eff} , sign convention $s \equiv +1$, and write $a := \log S$. The aggregator becomes

$$G(a, v) = \kappa_u \int_{-\infty}^{+\infty} q(\log K) \Gamma_{\text{BS}}(S, K, T_{\text{eff}}, \sqrt{v}) d(\log K). \quad (10)$$

Since Black–Scholes gamma at fixed $T = T_{\text{eff}}$, $\sigma = \sqrt{v}$ is itself Gaussian in $\log K$, the standard Gaussian-product identity collapses (10) to

$$G(a, v) = \mathcal{C}(v) \cdot e^{-a} \cdot \exp\left(-\frac{(a - m(v))^2}{2\tau^2(v)}\right), \quad (11)$$

where

$$\mathcal{C}(v) = \frac{\kappa_u e^{-q_{\text{div}} T_{\text{eff}}}}{\sqrt{2\pi \tau^2(v)}}, \quad \tau^2(v) = \sigma_q^2 + v T_{\text{eff}}, \quad m(v) = \mu_q - (r - q_{\text{div}} + \tfrac{1}{2}v) T_{\text{eff}}. \quad (12)$$

G depends on a and v but not on z ; the leverage channel γz in $\dot{u}$ keeps the $(2, 3)$ Jacobian entry alive, so cross-coupling persists through the variance row even with $G_z = 0$.

With $G_z = 0$ and $\sigma^2 = v$ (so $\partial_y \sigma^2 = 0$, $\partial_v \sigma^2 = 1$), the Routh–Hurwitz polynomial $H(\kappa) = c_1 c_2 - c_0$ degree-collapses from cubic to quadratic in κ :

$$H(\kappa) = G_y^2 (\alpha + \kappa_v) \kappa^2 + (G_v \beta \gamma - G_y (\alpha + \kappa_v)^2) \kappa + \alpha \kappa_v (\alpha + \kappa_v) - \tfrac{1}{2} \beta \gamma. \quad (13)$$

Writing $A := \alpha + \kappa_v$, $M := \alpha\kappa_v$, $L := \beta\gamma$, the Hopf threshold is the smallest positive root of the quadratic:

$$\kappa^* = \frac{G_y A^2 - G_v L - \sqrt{(G_v L - G_y A^2)^2 - 4 G_y^2 A (MA - L/2)}}{2 G_y^2 A}. \quad (14)$$

The discriminant in (14) provides the first parametric phase boundary: when $D := (G_v L - G_y A^2)^2 - 4 G_y^2 A (MA - L/2) < 0$, the system has no Hopf at any κ and the deterministic skeleton is unconditionally stable.

The third-order partials of (11) at the equilibrium are explicit rational functions of $(\delta := a^* - m(v^*), \tau^2, T_{\text{eff}})$, so the bilinear and trilinear tensors B, C at κ^* admit a closed form. Substituting into the Kuznetsov formula

$$\ell_1(\kappa^*) = \frac{1}{2\omega^*} \text{Re} \left[\langle p, C(q, q, \bar{q}) \rangle - 2 \langle p, B(q, J^{-1} B(q, \bar{q})) \rangle + \langle p, B(\bar{q}, (2i\omega^* I - J)^{-1} B(q, q)) \rangle \right], \quad (15)$$

with $Jq = i\omega^* q$, $J^\top p = -i\omega^* p$, $\langle p, q \rangle = 1$, yields a ~ 30 -term rational in the G -partials $(G_y, G_v, G_{aa}, G_{av}, G_{vv}, G_{aaa}, G_{aav}, G_{avv}, G_{vvv})$ and $(\kappa_v, \alpha, \beta, \gamma)$. The implementation evaluates this in $\sim 50 \mu\text{s}$ per parameter tuple, with results matching the FD-tensor pipeline to better than 0.6% relative on every parameter set tested.

At the canonical regime ($\sigma_q = 0.10, T_{\text{eff}} = 0.25, \kappa_v = 2, \alpha = 0.05, \beta = 1, \gamma = 1, \mu_q = \log 100, v^* = 0.04$), (14)–(15) give $\kappa^* = 17.81$, $\omega^* = 1.18 \text{ rad/yr}$, $\ell_1 = -4.81 \times 10^{-1}$ (supercritical). The closed form is essentially noise-free; with it in hand, the sign of ℓ_1 near the boundary contour is structurally certain rather than statistically inferred.

4 Numerical phase diagram

The full $(\kappa, \xi, \rho, \sigma_v)$ phase diagram is computed by sweeping $\kappa \in [0, 2]$ on a 401-point grid for each of $31 \times 21 \times 4 = 2,604$ cells over (ξ, ρ, σ_v) at the section 3.4 Hopf-exhibiting regime. The pair (ξ, ρ) enters the deterministic Jacobian via the leading shear-induced correction $a_{\text{eff}}(\kappa) = a(\kappa) + \frac{1}{2} \xi^2 \rho G_v$ —an Engel et al. (2024)-style projection of the small-noise stochastic Hopf onto the eigenvalue envelope. Wall-clock: $\sim 16 \text{ s}$ on a single M-series core. Figure 1 renders the (ξ, ρ) heatmap and four line cuts of $\kappa^*(\sigma_v)$ at SPX-relevant (ξ, ρ) probes.

The no-Hopf wedge in the upper-left (high ξ , strong negative ρ) is consistent with the shear-induced destabilisation prediction. The four line cuts substantiate the section 3.4 claim that real options markets sit *near but not across* κ^* in the SPX-relevant $(\xi, \rho) \approx (0.30, -0.70)$ corner—which is the structural mechanism we conjecture for the empirical $n \approx 1$ branching ratio of Hardiman et al. (2013), in the same heterogeneous-agent / endogenous-feedback tradition as Brock and Hommes (1998); Filimonov and Sornette (2012); Bouchaud (2011) (Conjecture: $\kappa \leftrightarrow n$; a formal reduction would itself be a separate paper).

Figure 2 renders the (σ_q, γ) phase boundary from the closed-form ℓ_1 of section 3.5. The supercritical region ($\ell_1 < 0$) is a narrow band bounded by two $\ell_1 = 0$ contours; for very small σ_q ($\lesssim 0.05$) the band collapses entirely. The two $\ell_1 = 0$ contours are the central parametric prediction: their joint geometry constrains which (OI distribution, leverage) pairs admit stable endogenous limit cycles.




Figure 1: **Hopf bifurcation threshold κ^* across the (ξ, ρ, σ_v) slice of the reflexive parameter space.** *Left (Panel A):* heatmap of $\kappa^*(\xi, \rho)$ at the median variance-of-variance slice $\sigma_v = 0.50$, computed at the canonical Hopf-exhibiting regime of section 3.4 ($G_x = 0.5$, $G_v = -0.5$, $G_z = -0.5$, $\alpha = 0.5$, $\beta = 1$, $\gamma = 0.5$, $\kappa_v = 2$). White cells are configurations with no Hopf in the scan range $\kappa \in [0, 2]$ —concentrated in the upper-left wedge of high vol-of-vol ξ and strong negative ρ , where the Engel et al. (2024) shear correction destabilises the eigenvalue envelope. Contour lines are isoquants of κ^* at increments of $(\kappa_{\max}^* - \kappa_{\min}^*)/6$. *Right (Panel B):* four line cuts of $\kappa^*(\sigma_v)$ at representative (ξ, ρ) probes: $(0.10, -0.70)$, $(0.30, -0.70)$, $(0.70, -0.70)$, and $(0.30, +0.70)$. The first three trace how moving radially outward in vol-of-vol at fixed SPX-like $\rho < 0$ pulls the Hopf threshold down; the last confirms that flipping ρ to $+0.70$ at moderate ξ keeps κ^* in the 0.85–0.95 band.

5 Pre-registered evaluation framework

The evaluation pipeline is committed at git commit 268c061 with an OpenTimestamps Bitcoin-anchored proof of the document hash, prior to any contact with empirical SPX data. This addresses the documented computational-reproducibility crisis in finance (Pérignon et al., 2024) and adapts the Camerer et al. (2018) pre-registration template to a computational-finance / RL-for-derivatives context. We do not claim “first pre-registration in finance”—the Pacific-Basin Finance Journal pathway (Brailsford et al., 2022) and the AEA RCT Registry (American Economic Association, 2018) predate this work. The novelty is the four-way intersection of [pre-registration + git commit-hash anchoring + block-bootstrap / BH-FDR statistical pre-spec + applied to RL-for-finance].

Two evaluation primitives. (i) Sliced-Wasserstein-2 (Bonneel et al., 2015) over arbitrage-free 21-day surface windows—the realism metric. (ii) κ -sensitivity slope at the calibrated coupling κ_0 —the reflexivity-importance scalar.

H1 (primary realism). A model-free RL agent π_{κ_0} trained inside the reflexive simulator at the calibrated coupling produces, in evaluation, an IV-surface distribution closer in sliced-W2 over arbitrage-free 21-trading-day rolling windows to the empirical SPX surface than any of four non-reflexive baselines (time-dependent Heston, LSV, 3/2 SV, gamma-aware-non-reflexive). Three event windows: Volmageddon (2018-02-05), COVID (2020-03-16), Yen carry unwind (2024-08-05). Decision rule: 12 pairwise dominance checks (4 baselines $\times$ 3 windows) with non-overlapping 95%




Figure 2: **Closed-form ℓ_1 phase boundary in (σ_q, γ) space at the canonical specification.** Supercritical ($\ell_1 < 0$, stable limit cycle), sub-critical ($\ell_1 > 0$, hysteresis and abrupt jumps), and no-Hopf regions delineated by the closed form of section 3.5. Computed in $\sim 50 \mu\text{s}$ per cell.

block-bootstrap CIs (Politis and Romano, 1994). The arbitrage filter follows Roper (2010); Gatheral and Jacquier (2014) on butterfly / calendar-spread / Lee-bound (Lee, 2004) criteria; sliced- W_2 sample-complexity follows Manole et al. (2022).

H2–H4 (secondary). H2 is the κ -sensitivity slope $\tilde{\rho} = \partial \hat{m} / \partial \kappa|_{\kappa_0}$ —positive and statistically distinguishable from zero for the reflexive-trained agent, statistically indistinguishable for baselines (TOST equivalence, applied to the dimensionless elasticity). H3 is the post-FOMC RR25 and ATM-term-structure shifts. H4 is the Hopf-frequency spectral peak in $|r_t|$ at $\omega^* \pm 20\%$ via Welch’s method, with an IAAFT (Schreiber and Schmitz, 1996) surrogate null. Figure 3 reports H4 detector power on the synthetic positive-control regime.

Differentiation from precedents. Ning et al. (2024) use W_1 on single-day surface marginals; we use sliced- W_2 on 21-day windows with a hard arbitrage pre-filter, block-bootstrap CIs, and per-regime reporting. He et al. (2025) (NeurIPS 2025) sweep a Wasserstein-ball perturbation budget with a separate adversarially-trained agent at each grid point; we deploy a single agent across the κ grid and report the slope at anchor with a GP-posterior CI (Hou et al., 2025) for the elliptic-uncertainty-set analogue. Subbaswamy et al. (2022) (NeurIPS 2022) provide the closest theoretical precursor for slope-at-anchor robustness summaries (parametric robustness sets, local 2nd-order approximation), which we operationalise empirically and apply to RL-for-finance (Bai et al., 2024; Huang et al., 2024). Conditional Wasserstein for state-conditional generator comparison follows Hosseini et al. (2024). Building blocks: Vuletić and Cont (2025); Choudhary et al. (2024); Issa et al. (2023); Jin and Agarwal (2025); Chevallier et al. (2025); Stadnytska et al. (2025); Buehler et al. (2019); Cao et al. (2021); Packer et al. (2018); Murray et al. (2022).




Figure 3: **H4 spectral-peak detector power on synthetic ground truth.** Power curves at canonical settings (Welch window adaptive to ω^* ; IAAFT surrogate null per pre-registration amendment A5). Positive control: limit-cycle Stuart–Landau and supercritical reflexive simulator. Negative controls: white noise, Heston.

6 Mechanism decomposition vs Marketron

The reflexive simulator and the Marketron quasi-particle SDE (Halperin and Itkin, 2025) are not the same mechanism, and a 1:1 reproduction of Marketron’s Tables 7–8 from our simulator is mathematically impossible: in our model the memory carrier is the low-pass-filtered log-price z entering only the variance drift via γz ; in Marketron it is the unobservable y entering both drifts non-linearly through a potential $V_M(x)$. Even at zero coupling our model collapses to standard Heston while Marketron does not collapse to anything standard.

We instead report mechanism decomposition: a 5D coarse-grid tuning over $(\kappa, \gamma, T_{\text{eff}}, \mu_q, \sigma_q)$, then per-cell classification into `shape_target` / `level_artifact` / `calibration_artifact`. The headline gate is the shape-match rate on `shape_target` cells.

Aggregate finding	Value
Shape-feature cells matched (out-of-sample, 10k paths)	8 / 24 (33.3%)
$\geq 30\%$ gate (one-sided binomial $p \approx 0.27$)	passed
Long-horizon (≥ 0.5 y) excess-kurt sign agreement	5 / 6 cells
Long-horizon skew sign agreement (risk-neutral drift)	predictable disagreement

Table 2: Mechanism-decomposition headline. Long-horizon excess-kurt sign agrees with Marketron Table 8 on 5/6 horizons; long-horizon skew sign disagrees predictably under risk-neutral drift (Marketron’s positive long-horizon skew comes from Bessembinder (2018) compounding under calibrated drift; ours comes from dealer-gamma + leverage feedback under $\mu = 0$, see also Farago and Hjalmarsson, 2023).

The skew-sign disagreement is a falsifiable empirical prediction: in the Phase 4 empirical calibration with reproduced drift, long-horizon skew should flip positive. The two mechanisms are independent in parameterisation but agree on the long-horizon excess-kurt sign—the most robust agreement.

7 Conclusions and future work

We have introduced a 3D reflexive SV simulator (S_t, v_t, z_t) with explicit dealer-gamma drift, proved a Hopf bifurcation theorem with a closed-form first Lyapunov coefficient ℓ_1 and Hopf threshold κ^* for log-normal open-interest in moneyness, computed a numerical (ξ, ρ, σ_v) phase diagram and a (σ_q, γ) closed-form phase boundary, and pre-registered the empirical evaluation pipeline (sliced- W_2 , κ -sensitivity slope, Hopf-frequency spectral test) at a verifiable git commit hash with an OpenTimestamps Bitcoin-anchored proof.

The empirical calibration to real SPX surfaces is the natural Phase 4 follow-up, gated on data acquisition (UoT WRDS / OptionMetrics IvyDB US, or commercial fallback). The pre-registration locks the analysis pipeline before the calibration touches the data; results—confirming or refuting H1–H4—are pre-committed to publication regardless of direction, in the spirit of the documented reproducibility response (Pérignon et al., 2024). Open theoretical items: closed-form $\Lambda(\kappa)$ via Engel–Lamb–Rasmussen-style asymptotics adapted to Heston multiplicative noise (currently numerical only); a formal Hawkes- n to SV-Jacobian-eigenvalue reduction relating κ^* to the empirical $n \approx 1$ (Hardiman et al., 2013; Bondi et al., 2024; Bacry et al., 2015).

The Fokker–Planck stationary marginal density of the reflexive simulator deviates from the closed-form Heston comparator (Feller, 1951) on tail index and skew sign in a way that tracks $\text{sgn}(G_x)$, while the bimodality test of Hartigan and Hartigan (1985) is uninformative under the $\gamma = 0$ closure used for that scan. Detector validation uses the Benettin et al. (1980) Lyapunov-renormalisation algorithm and the Schreiber and Schmitz (1996) IAAFT surrogate; the wider Hawkes-and-rough-volatility line is Bondi et al. (2024); Bacry et al. (2015); Livieri et al. (2018); arbitrage-free neural SDE market models follow Cohen et al. (2023).

All experiments are reproducible via `bash scripts/verify.sh` from commit 268c061; pre-registration anchored via OpenTimestamps. Code: github.com/mahimn01/reflexive-options.

References

- American Economic Association. AEA RCT registry. <https://www.aeaweb.org/journals/policies/rct-registry>, 2018.
- Ludwig Arnold. *Random Dynamical Systems*. Springer, 1998.
- Emmanuel Bacry, Iacopo Mastromatteo, and Jean-François Muzy. Hawkes processes in finance. *Market Microstructure and Liquidity*, 1(1), 2015.
- Junkun Bai, Xinyu Gao, Cheng Wan, Yuan Zhang, and Le Song. A review of reinforcement learning in financial applications. *arXiv preprint arXiv:2411.12746*, 2024. URL <https://arxiv.org/abs/2411.12746>.
- Andrea Barbon and Andrea Buraschi. Gamma fragility. *SSRN Working Paper*, 2021.
- Peter H. Baxendale. A stochastic Hopf bifurcation. *Probability Theory and Related Fields*, 99: 581–616, 1994.
- Giancarlo Benettin, Luigi Galgani, Antonio Giorgilli, and Jean-Marie Strelcyn. Lyapunov characteristic exponents for smooth dynamical systems and for Hamiltonian systems. *Meccanica*, 15:9–30, 1980.
- Hendrik Bessembinder. Do stocks outperform treasury bills? *Journal of Financial Economics*, 129(3):440–457, 2018. doi: 10.1016/j.jfineco.2018.06.004.

- Alessandro Bondi, Sergio Pulido, and Simone Scotti. The rough Hawkes Heston stochastic volatility model. *Mathematical Finance*, 34(4), 2024.
- Nicolas Bonneel, Julien Rabin, Gabriel Peyré, and Hanspeter Pfister. Sliced and Radon Wasserstein barycenters of measures. *Journal of Mathematical Imaging and Vision*, 51(1):22–45, 2015. doi: 10.1007/s10851-014-0506-3.
- Jean-Philippe Bouchaud. The endogenous dynamics of markets: Price impact, feedback loops and instabilities. 2011.
- Tim Brailsford et al. Editorial: Pre-registration in finance. *Pacific-Basin Finance Journal*, 2022. doi: 10.1016/j.pacfin.2022.101859.
- William A. Brock and Cars H. Hommes. Heterogeneous beliefs and routes to chaos in a simple asset pricing model. *Journal of Economic Dynamics and Control*, 22(8–9):1235–1274, 1998. doi: 10.1016/S0165-1889(98)00011-6.
- William A. Brock, Cars H. Hommes, and Florian O. O. Wagener. More hedging instruments may destabilize markets. *Journal of Economic Dynamics and Control*, 33(11):1912–1928, 2009. doi: 10.1016/j.jedc.2009.05.004.
- Hans Buehler, Lukas Gonon, Josef Teichmann, and Ben Wood. Deep hedging. *Quantitative Finance*, 19(8):1271–1291, 2019. doi: 10.1080/14697688.2019.1571683.
- Colin F. Camerer, Anna Dreber, Felix Holzmeister, Teck-Hua Ho, Jürgen Huber, Magnus Johannesson, Michael Kirchler, Gideon Nave, Brian A. Nosek, Thomas Pfeiffer, et al. Evaluating the replicability of social science experiments in Nature and Science between 2010 and 2015. *Nature Human Behaviour*, 2:637–644, 2018. doi: 10.1038/s41562-018-0399-z.
- Jay Cao, Jacky Chen, John Hull, and Zissis Poulos. Deep hedging of derivatives using reinforcement learning. *arXiv preprint arXiv:2103.16409*, 2021. URL <https://arxiv.org/abs/2103.16409>.
- Jack Carr. *Applications of Centre Manifold Theory*. Springer, 1981.
- Julien Chevallier, Stefano De Marco, and Antoine Lévy-dit Vehel. Optimal transport-based approach to arbitrage correction of implied volatility surfaces. *arXiv preprint arXiv:2501.12195*, 2025. URL <https://arxiv.org/abs/2501.12195>.
- Carl Chiarella, Xue-Zhong He, and Cars H. Hommes. The dynamics of moving average rules in a continuous-time financial market model. Technical Report RP 268, UTS Quantitative Finance Research Centre, 2006. URL <https://www.uts.edu.au/globalassets/sites/default/files/qfr-archive-03/QFR-rp268.pdf>.
- Vedant Choudhary, Sebastian Jaimungal, and Maxime Bergeron. FuNVol: Multi-asset implied volatility market simulator using functional principal components and neural SDEs. *Quantitative Finance*, 2024. doi: 10.1080/14697688.2024.2396977.
- Samuel N. Cohen, Christoph Reisinger, and Sheng Wang. Arbitrage-free neural-SDE market models. *Applied Mathematical Finance*, 2023.
- Haoying Dai. Beta-dependent gamma feedback and endogenous volatility amplification in option markets. *arXiv preprint arXiv:2511.22766*, 2025. URL <https://arxiv.org/abs/2511.22766>.

- Chukwuma Dim, Bjørn Eraker, and Grigory Vilkov. 0DTEs: Trading, gamma risk, and volatility propagation. *SSRN Working Paper*, 2024.
- Maximilian Engel, Jeroen S. W. Lamb, and Martin Rasmussen. Lyapunov exponents and shear-induced chaos for a Hopf bifurcation with additive noise. *Probability Theory and Related Fields*, 188, 2024. doi: 10.1007/s00440-024-01301-4.
- Adam Farago and Erik Hjalmarsson. Compound returns. *Journal of Financial Economics*, 2023.
- William Feller. Two singular diffusion problems. *Annals of Mathematics*, 54(1):173–182, 1951.
- Vladimir Filimonov and Didier Sornette. Quantifying reflexivity in financial markets: Toward a prediction of flash crashes. *Physical Review E*, 85:056108, 2012. doi: 10.1103/PhysRevE.85.056108.
- Rüdiger Frey and Alexander Stremme. Market volatility and feedback effects from dynamic hedging. *Mathematical Finance*, 7(4):351–374, 1997. doi: 10.1111/1467-9965.00036.
- Nicolae Gârleanu, Lasse Heje Pedersen, and Allen M. Poteshman. Demand-based option pricing. *Review of Financial Studies*, 22(10):4259–4299, 2009. doi: 10.1093/rfs/hhp005.
- Jim Gatheral and Antoine Jacquier. Arbitrage-free SVI volatility surfaces. *Quantitative Finance*, 14(1):59–71, 2014. URL <https://arxiv.org/abs/1204.0646>.
- Igor Halperin and Andrey Itkin. Marketron through the looking glass: From equity dynamics to option pricing in incomplete markets. *arXiv preprint arXiv:2508.09863*, 2025. URL <https://arxiv.org/abs/2508.09863>.
- Stephen J. Hardiman, Nicolas Bercot, and Jean-Philippe Bouchaud. Critical reflexivity in financial markets: A Hawkes process analysis. *European Physical Journal B*, 86:442, 2013. URL <https://arxiv.org/abs/1302.1405>.
- John A. Hartigan and P. M. Hartigan. The dip test of unimodality. *Annals of Statistics*, 13(1):70–84, 1985. doi: 10.1214/aos/1176346577.
- Guangyi He, Tobias Sutter, and Lukas Gonon. Distributional adversarial attacks and training in deep hedging. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2025. URL <https://arxiv.org/abs/2508.14757>.
- Xue-Zhong He, Kai Li, and Min Zheng. Market stability switches in a continuous-time financial market with heterogeneous beliefs. *Economic Modelling*, 26:1432–1442, 2009. doi: 10.1016/j.econmod.2009.07.011.
- Steven L. Heston. A closed-form solution for options with stochastic volatility with applications to bond and currency options. *Review of Financial Studies*, 6(2):327–343, 1993. doi: 10.1093/rfs/6.2.327.
- Bamdad Hosseini, Charlotte Bunne, and Marco Cuturi. Conditional optimal transport on function spaces. *arXiv preprint arXiv:2403.18705*, 2024. URL <https://arxiv.org/abs/2403.18705>.
- Yifan Hou et al. Robust reinforcement learning in finance: Modeling market impact with elliptic uncertainty sets. *arXiv preprint arXiv:2510.19950*, 2025. URL <https://arxiv.org/abs/2510.19950>.

- Shengyi Huang et al. Open RL benchmark: Comprehensive tracked experiments for reinforcement learning. *arXiv preprint arXiv:2402.03046*, 2024. URL <https://arxiv.org/abs/2402.03046>.
- Zacharia Issa, Blanka Horvath, Maud Lemerrier, and Cristopher Salvi. Non-adversarial training of neural SDEs with signature kernel scores. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2023. URL <https://arxiv.org/abs/2305.16274>.
- Yufeng Jin and Ankur Agarwal. Forecasting implied volatility surface with generative diffusion models. *arXiv preprint arXiv:2511.07571*, 2025. URL <https://arxiv.org/abs/2511.07571>.
- Rafail Z. Khasminskii. *Stochastic Stability of Differential Equations*. Sijthoff and Noordhoff, 1980.
- Yuri A. Kuznetsov. *Elements of Applied Bifurcation Theory*. Springer, 3rd edition, 2004.
- Roger W. Lee. The moment formula for implied volatility at extreme strikes. *Mathematical Finance*, 14(3):469–480, 2004. doi: 10.1111/j.0960-1627.2004.00200.x.
- Giulia Livieri, Saad Mouti, Andrea Pallavicini, and Mathieu Rosenbaum. Rough volatility: Evidence from option prices. *IIE Transactions*, 50(9):767–776, 2018.
- Roger Lord, Remmert Koekkoek, and Dick van Dijk. A comparison of biased simulation schemes for stochastic volatility models. *Quantitative Finance*, 10(2):177–194, 2010. doi: 10.1080/14697680802392496.
- Tudor Manole, Sivaraman Balakrishnan, and Larry Wasserman. Minimax confidence intervals for the sliced Wasserstein distance. *Electronic Journal of Statistics*, 2022. URL <https://arxiv.org/abs/1909.07862>.
- Phillip Murray, Ben Wood, Hans Buehler, Magnus Wiese, and Mikko Pakkanen. Deep hedging: Continuous reinforcement learning across multiple risk aversions. *arXiv preprint arXiv:2207.07467*, 2022. URL <https://arxiv.org/abs/2207.07467>.
- Brian Ning, Sebastian Jaimungal, Xiaorong Zhang, and Maxime Bergeron. Arbitrage-free implied volatility surface generation with variational autoencoders. *SIAM Journal on Financial Mathematics*, 15(1), 2024. doi: 10.1137/21M1443546.
- Charles Packer, Katelyn Gao, Jernej Kos, Philipp Krähenbühl, Vladlen Koltun, and Dawn Song. Assessing generalization in deep reinforcement learning. *arXiv preprint arXiv:1810.12282*, 2018. URL <https://arxiv.org/abs/1810.12282>.
- Christophe Pérignon, Olivier Akmansoy, Christophe Hurlin, Anna Dreber, Felix Holzmeister, Jürgen Huber, Magnus Johannesson, Michael Kirchler, Albert J. Menkveld, Michael Razen, and Utz Weitzel. Computational reproducibility in finance: Evidence from 1,000 tests. *Review of Financial Studies*, 37(11):3558, 2024. doi: 10.1093/rfs/hhae029.
- Eckhard Platen and Martin Schweizer. On feedback effects from hedging derivatives. *Mathematical Finance*, 8(1):67–84, 1998. doi: 10.1111/1467-9965.00038.
- Dimitris N. Politis and Joseph P. Romano. The stationary bootstrap. *Journal of the American Statistical Association*, 89(428):1303–1313, 1994. doi: 10.1080/01621459.1994.10476870.
- Michael Roper. Arbitrage-free implied volatility surfaces. Technical report, University of Sydney, 2010.

- Thomas Schreiber and Andreas Schmitz. Improved surrogate data for nonlinearity tests. *Physical Review Letters*, 77(4):635–638, 1996. doi: 10.1103/PhysRevLett.77.635.
- K. Ronnie Sircar and George Papanicolaou. General Black–Scholes models accounting for increased market volatility from hedging strategies. *Applied Mathematical Finance*, 5(1):45–82, 1998.
- George Soros. The general theory of reflexivity. Open Society Foundations Working Paper, 2009.
- SqueezeMetrics. The implied order book and gamma exposure. Whitepaper, squeezemetrics.com, 2017.
- Tetyana Stadnytska et al. Automated regime classification via sliced-Wasserstein k -means. *Data Science in Finance and Economics (AIMS)*, 2025. doi: 10.3934/DSFE.2025016.
- Adarsh Subbaswamy, Roy Adams, and Suchi Saria. Evaluating robustness to dataset shift via parametric robustness sets. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2022. URL https://proceedings.neurips.cc/paper_files/paper/2022/file/6b7f9d9c1217a748391800871ff7d17d-Paper-Conference.pdf.
- Milena Vuletić and Rama Cont. VolGAN: A generative model for arbitrage-free implied volatility surfaces. *Applied Mathematical Finance*, 2025. doi: 10.1080/1350486X.2025.2471317.