

Naïve Gauss Elimination Partial Pivoting

Author: Autar Kaw et. al.

Textbook: TEXTBOOK: NUMERICAL METHODS WITH
APPLICATIONS

Naïve Gauss Elimination Pitfalls

Pitfall#1. Division by zero

$$10x_2 - 7x_3 = 3$$

$$6x_1 + 2x_2 + 3x_3 = 11$$

$$5x_1 - x_2 + 5x_3 = 9$$

$$\begin{bmatrix} 0 & 10 & -7 \\ 6 & 2 & 3 \\ 5 & -1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 11 \\ 9 \end{bmatrix}$$

Is division by zero an issue here?

$$12x_1 + 10x_2 - 7x_3 = 15$$

$$6x_1 + 5x_2 + 3x_3 = 14$$

$$5x_1 - x_2 + 5x_3 = 9$$

$$\begin{bmatrix} 12 & 10 & -7 \\ 6 & 5 & 3 \\ 5 & -1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 15 \\ 14 \\ 9 \end{bmatrix}$$

Is division by zero an issue here?

YES

$$12x_1 + 10x_2 - 7x_3 = 15$$

$$6x_1 + 5x_2 + 3x_3 = 14$$

$$24x_1 - x_2 + 5x_3 = 28$$

$$\begin{bmatrix} 12 & 10 & -7 \\ 6 & 5 & 3 \\ 24 & -1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 15 \\ 14 \\ 28 \end{bmatrix} \longrightarrow \begin{bmatrix} 12 & 10 & -7 \\ 0 & 0 & 6.5 \\ 12 & -21 & 19 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 15 \\ 6.5 \\ -2 \end{bmatrix}$$

Division by zero is a possibility at any step
of forward elimination

Pitfall#2. Large Round-off Errors

$$\begin{bmatrix} 20 & 15 & 10 \\ -3 & -2.249 & 7 \\ 5 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 45 \\ 1.751 \\ 9 \end{bmatrix}$$

Exact Solution

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Pitfall#2. Large Round-off Errors

$$\begin{bmatrix} 20 & 15 & 10 \\ -3 & -2.249 & 7 \\ 5 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 45 \\ 1.751 \\ 9 \end{bmatrix}$$

Solve it on a computer using **6** significant digits with chopping

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0.9625 \\ 1.05 \\ 0.999995 \end{bmatrix}$$

Pitfall#2. Large Round-off Errors

$$\begin{bmatrix} 20 & 15 & 10 \\ -3 & -2.249 & 7 \\ 5 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 45 \\ 1.751 \\ 9 \end{bmatrix}$$

Solve it on a computer using **5** significant digits with chopping

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0.625 \\ 1.5 \\ 0.99995 \end{bmatrix}$$

Is there a way to reduce the round off error?

Avoiding Pitfalls

Increase the number of significant digits

- Decreases round-off error
- Does not avoid division by zero

Avoiding Pitfalls

Gaussian Elimination with Partial Pivoting

- Avoids division by zero
- Reduces round off error

Gauss Elimination with Partial Pivoting

What is Different About Partial Pivoting?

At the beginning of the k^{th} step of forward elimination, find the maximum of

$$|a_{kk}|, |a_{k+1,k}|, \dots, |a_{nk}|$$

If the maximum of the values is $|a_{pk}|$

in the p^{th} row, $k \leq p \leq n$, then switch rows p and k .

Matrix Form at Beginning of 2nd Step of Forward Elimination

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a'_{22} & a'_{23} & \cdots & a'_{2n} \\ 0 & a'_{32} & a'_{33} & \cdots & a'_{3n} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & a'_{n2} & a'_{n3} & a'_{n4} & a'_{nn} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b'_2 \\ b'_3 \\ \vdots \\ b'_n \end{bmatrix}$$

Example (2nd step of FE)

$$\begin{bmatrix} 6 & 14 & 5.1 & 3.7 & 6 \\ 0 & -7 & 6 & 1 & 2 \\ 0 & 4 & 12 & 1 & 11 \\ 0 & 9 & 23 & 6 & 8 \\ 0 & -17 & 12 & 11 & 43 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 5 \\ -6 \\ 8 \\ 9 \\ 3 \end{bmatrix}$$

Which two rows would you switch?

Example (2nd step of FE)

$$\begin{bmatrix} 6 & 14 & 5.1 & 3.7 & 6 \\ 0 & -17 & 12 & 11 & 43 \\ 0 & 4 & 12 & 1 & 11 \\ 0 & 9 & 23 & 6 & 8 \\ 0 & -7 & 6 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 5 \\ 3 \\ 8 \\ 9 \\ -6 \end{bmatrix}$$

Switched Rows

Gauss Elimination with Partial Pivoting Example

Example

Solve the following set of equations by Gaussian elimination with partial pivoting

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

Example Cont.

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix} \Rightarrow \begin{bmatrix} 25 & 5 & 1 & \vdots & 106.8 \\ 64 & 8 & 1 & \vdots & 177.2 \\ 144 & 12 & 1 & \vdots & 279.2 \end{bmatrix}$$

1. Forward Elimination
2. Back Substitution

Forward Elimination

Number of Steps of Forward Elimination

Number of steps of forward elimination is
 $(n-1)=(3-1)=2$

Forward Elimination: Step 1

- Examine absolute values of first column, first row and below.

$$|25|, |64|, |144|$$

- Largest absolute value is 144 and exists in row 3.
- Switch row 1 and row 3.

$$\begin{bmatrix} 25 & 5 & 1 & \vdots & 106.8 \\ 64 & 8 & 1 & \vdots & 177.2 \\ 144 & 12 & 1 & \vdots & 279.2 \end{bmatrix} \Rightarrow \begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 64 & 8 & 1 & \vdots & 177.2 \\ 25 & 5 & 1 & \vdots & 106.8 \end{bmatrix}$$

Forward Elimination: Step 1 (cont.)

$$\begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 64 & 8 & 1 & \vdots & 177.2 \\ 25 & 5 & 1 & \vdots & 106.8 \end{bmatrix}$$

Divide Equation 1 by 144 and multiply it by 64, $\frac{64}{144} = 0.4444$.

$$[144 \quad 12 \quad 1 \quad \vdots \quad 279.2] \times 0.4444 = [63.99 \quad 5.333 \quad 0.4444 \quad \vdots \quad 124.1]$$

Subtract the result from
Equation 2

$$\begin{array}{r} \begin{bmatrix} 64 & 8 & 1 & \vdots & 177.2 \end{bmatrix} \\ - \begin{bmatrix} 63.99 & 5.333 & 0.4444 & \vdots & 124.1 \end{bmatrix} \\ \hline \begin{bmatrix} 0 & 2.667 & 0.5556 & \vdots & 53.10 \end{bmatrix} \end{array}$$

Substitute new equation for
Equation 2

$$\begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 0 & 2.667 & 0.5556 & \vdots & 53.10 \\ 25 & 5 & 1 & \vdots & 106.8 \end{bmatrix}$$

Forward Elimination: Step 1 (cont.)

$$\begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 0 & 2.667 & 0.5556 & \vdots & 53.10 \\ 25 & 5 & 1 & \vdots & 106.8 \end{bmatrix} \quad \begin{array}{l} \text{Divide Equation 1 by 144 and} \\ \text{multiply it by 25, } \frac{25}{144} = 0.1736. \end{array}$$

$$[144 \quad 12 \quad 1 \quad \vdots \quad 279.2] \times 0.1736 = [25.00 \quad 2.083 \quad 0.1736 \quad \vdots \quad 48.47]$$

Subtract the result from
Equation 3

$$\begin{array}{r} \begin{bmatrix} 25 & 5 & 1 & \vdots & 106.8 \end{bmatrix} \\ - \begin{bmatrix} 25 & 2.083 & 0.1736 & \vdots & 48.47 \end{bmatrix} \\ \hline \begin{bmatrix} 0 & 2.917 & 0.8264 & \vdots & 58.33 \end{bmatrix} \end{array}$$

Substitute new equation for
Equation 3

$$\begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 0 & 2.667 & 0.5556 & \vdots & 53.10 \\ 0 & 2.917 & 0.8264 & \vdots & 58.33 \end{bmatrix}$$

Forward Elimination: Step 2

- Examine absolute values of second column, second row and below.

$$|2.667|, |2.917|$$

- Largest absolute value is 2.917 and exists in row 3.
- Switch row 2 and row 3.

$$\begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 0 & 2.667 & 0.5556 & \vdots & 53.10 \\ 0 & 2.917 & 0.8264 & \vdots & 58.33 \end{bmatrix} \Rightarrow \begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 0 & 2.917 & 0.8264 & \vdots & 58.33 \\ 0 & 2.667 & 0.5556 & \vdots & 53.10 \end{bmatrix}$$

Forward Elimination: Step 2 (cont.)

$$\begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 0 & 2.917 & 0.8264 & \vdots & 58.33 \\ 0 & 2.667 & 0.5556 & \vdots & 53.10 \end{bmatrix}$$

Divide Equation 2 by 2.917 and multiply it by 2.667,
 $\frac{2.667}{2.917} = 0.9143$.

$$[0 \quad 2.917 \quad 0.8264 \quad \vdots \quad 58.33] \times 0.9143 = [0 \quad 2.667 \quad 0.7556 \quad \vdots \quad 53.33]$$

Subtract the result from
Equation 3

$$\begin{array}{r} \begin{bmatrix} 0 & 2.667 & 0.5556 & \vdots & 53.10 \end{bmatrix} \\ - \begin{bmatrix} 0 & 2.667 & 0.7556 & \vdots & 53.33 \end{bmatrix} \\ \hline \begin{bmatrix} 0 & 0 & -0.2 & \vdots & -0.23 \end{bmatrix} \end{array}$$

Substitute new equation for
Equation 3

$$\begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 0 & 2.917 & 0.8264 & \vdots & 58.33 \\ 0 & 0 & -0.2 & \vdots & -0.23 \end{bmatrix}$$

Back Substitution

Back Substitution

$$\begin{bmatrix} 144 & 12 & 1 & \vdots & 279.2 \\ 0 & 2.917 & 0.8264 & \vdots & 58.33 \\ 0 & 0 & -0.2 & \vdots & -0.23 \end{bmatrix} \Rightarrow \begin{bmatrix} 144 & 12 & 1 \\ 0 & 2.917 & 0.8264 \\ 0 & 0 & -0.2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 279.2 \\ 58.33 \\ -0.23 \end{bmatrix}$$

Solving for a_3

$$-0.2a_3 = -0.23$$

$$a_3 = \frac{-0.23}{-0.2}$$

$$= 1.15$$

Back Substitution (cont.)

$$\begin{bmatrix} 144 & 12 & 1 \\ 0 & 2.917 & 0.8264 \\ 0 & 0 & -0.2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 279.2 \\ 58.33 \\ -0.23 \end{bmatrix}$$

Solving for a_2

$$2.917a_2 + 0.8264a_3 = 58.33$$

$$\begin{aligned} a_2 &= \frac{58.33 - 0.8264a_3}{2.917} \\ &= \frac{58.33 - 0.8264 \times 1.15}{2.917} \\ &= 19.67 \end{aligned}$$

Back Substitution (cont.)

$$\begin{bmatrix} 144 & 12 & 1 \\ 0 & 2.917 & 0.8264 \\ 0 & 0 & -0.2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 279.2 \\ 58.33 \\ -0.23 \end{bmatrix}$$

Solving for a_1

$$144a_1 + 12a_2 + a_3 = 279.2$$

$$\begin{aligned} a_1 &= \frac{279.2 - 12a_2 - a_3}{144} \\ &= \frac{279.2 - 12 \times 19.67 - 1.15}{144} \\ &= 0.2917 \end{aligned}$$

Gaussian Elimination with Partial Pivoting Solution

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 0.2917 \\ 19.67 \\ 1.15 \end{bmatrix}$$

Partial Pivoting: Example 2

Consider the system of equations

$$10x_1 - 7x_2 = 7$$

$$-3x_1 + 2.099x_2 + 6x_3 = 3.901$$

$$5x_1 - x_2 + 5x_3 = 6$$

In matrix form

$$\begin{bmatrix} 10 & -7 & 0 \\ -3 & 2.099 & 6 \\ 5 & -1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 3.901 \\ 6 \end{bmatrix}$$

Solve using Gaussian Elimination with Partial Pivoting using five significant digits with chopping

Partial Pivoting: Example

Forward Elimination: Step 1

Examining the values of the first column

$|10|$, $|-3|$, and $|5|$ or 10, 3, and 5

The largest absolute value is 10, which means, to follow the rules of Partial Pivoting, we switch row1 with row1.

Performing Forward Elimination

$$\begin{bmatrix} 10 & -7 & 0 \\ -3 & 2.099 & 6 \\ 5 & -1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 3.901 \\ 6 \end{bmatrix} \Rightarrow \begin{bmatrix} 10 & -7 & 0 \\ 0 & -0.001 & 6 \\ 0 & 2.5 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 6.001 \\ 2.5 \end{bmatrix}$$

Partial Pivoting: Example

Forward Elimination: Step 2

Examining the values of the first column

$|-0.001|$ and $|2.5|$ or 0.0001 and 2.5

The largest absolute value is 2.5 , so row 2 is switched with row 3

Performing the row swap

$$\begin{bmatrix} 10 & -7 & 0 \\ 0 & -0.001 & 6 \\ 0 & 2.5 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 6.001 \\ 2.5 \end{bmatrix} \Rightarrow \begin{bmatrix} 10 & -7 & 0 \\ 0 & 2.5 & 5 \\ 0 & -0.001 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 2.5 \\ 6.001 \end{bmatrix}$$

Partial Pivoting: Example

Forward Elimination: Step 2

Performing the Forward Elimination results in:

$$\begin{bmatrix} 10 & -7 & 0 \\ 0 & 2.5 & 5 \\ 0 & 0 & 6.002 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 2.5 \\ 6.002 \end{bmatrix}$$

Partial Pivoting: Example

Back Substitution

Solving the equations through back substitution

$$\begin{bmatrix} 10 & -7 & 0 \\ 0 & 2.5 & 5 \\ 0 & 0 & 6.002 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 2.5 \\ 6.002 \end{bmatrix}$$

$$x_3 = \frac{6.002}{6.002} = 1$$

$$x_2 = \frac{2.5 - 5x_3}{2.5} = -1$$

$$x_1 = \frac{7 + 7x_2 - 0x_3}{10} = 0$$

Partial Pivoting: Example

Compare the calculated and exact solution

The fact that they are equal is coincidence, but it does illustrate the advantage of Partial Pivoting

$$[X]_{\text{calculated}} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad [X]_{\text{exact}} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

THE END