

Chapter 2: Divide-and-Conquer Algorithms

To solve a problem of size n :

- **Divide** the problem into subproblems, each of size $< n$.
- **Conquer**: Solve each subproblem recursively (and independantly of the other subproblems).
- **Combine/Merge** the solutions to the subproblems into a solution to the original problem.

Chapter 2: Divide-and-Conquer Algorithms

To solve a problem of size n :

- **Divide** the problem into subproblems, each of size $< n$.
- **Conquer**: Solve each subproblem recursively (and independantly of the other subproblems).
- **Combine/Merge** the solutions to the subproblems into a solution to the original problem.

For a given problem,

- How do we divide the problem, into how many subproblems?
- How to combine/merge?

Example: Merge Sort

To sort n numbers:

If $n \leq 1$: do nothing.

If $n \geq 2$: divide the n numbers arbitrarily into two sequences, both of size $n/2$, run Merge sort twice, once for each sequence.

Then merge the two sorted sequences into one sorted sequence.

Example: Merge Sort

To sort n numbers:

If $n \leq 1$: do nothing.

If $n \geq 2$: divide the n numbers arbitrarily into two sequences, both of size $n/2$, run Merge sort twice, once for each sequence.

Then merge the two sorted sequences into one sorted sequence.

What is the running time?

Let $T(n)$ be the running time of Merge Sort on an input of n numbers.

Let $T(n)$ be the running time of Merge Sort on an input of n numbers.

From CSI-2110, we know that the merge step takes $O(n)$ time.

Let $T(n)$ be the running time of Merge Sort on an input of n numbers.

From CSI-2110, we know that the merge step takes $O(n)$ time.

Hence, there is a constant $c > 0$ such that

$$T(n) \leq \begin{cases} c & \text{if } n = 1, \\ 2 T(n/2) + c \cdot n & \text{if } n \geq 2, \end{cases}$$

Let $T(n)$ be the running time of Merge Sort on an input of n numbers.

From CSI-2110, we know that the merge step takes $O(n)$ time.

Hence, there is a constant $c > 0$ such that

$$T(n) \leq \begin{cases} c & \text{if } n = 1, \\ 2 T(n/2) + c \cdot n & \text{if } n \geq 2, \end{cases}$$

What do we do with this?

Let $T(n)$ be the running time of Merge Sort on an input of n numbers.

From CSI-2110, we know that the merge step takes $O(n)$ time.

Hence, there is a constant $c > 0$ such that

$$T(n) \leq \begin{cases} c & \text{if } n = 1, \\ 2 T(n/2) + c \cdot n & \text{if } n \geq 2, \end{cases}$$

What do we do with this?

We solve by *unfolding*!

Solve this recurrence by unfolding:

Assume $n = 2^k$, $c = 1$

$$T(n) \leq n + 2T\left(\frac{n}{2}\right)$$

$$\leq n + 2 \cdot \left(\frac{n}{2} + 2 \cdot T\left(\frac{n}{4}\right) \right)$$

$$= 2n + 4T\left(\frac{n}{4}\right)$$

$$\leq 2n + 4 \left(\frac{n}{4} + 2 \cdot T\left(\frac{n}{8}\right) \right)$$

$$= 3n + 8T\left(\frac{n}{8}\right)$$

$$\leq 3n + 8 \left(\frac{n}{8} + 2 \cdot T\left(\frac{n}{16}\right) \right)$$

$$= 4n + 16T\left(\frac{n}{16}\right)$$

...

$$\leq kn + 2^k T\left(\frac{n}{2^k}\right)$$

$$= kn + n \cdot 1$$

$$= n \log_2(n) + n$$

$$\leq n \log_2(n) + n \cdot \log_2(n) \quad \leftarrow \text{since } \log_2(n) = \log_2(2^k) = k$$

$$\leq 2n \log_2(n)$$

In general, if we do not assume anything about the value of c , we find $T(n) \leq 2c n \log_2(n)$.

In general, if we do not assume anything about the value of c , we find $T(n) \leq 2c n \log_2(n)$.

So the running time of Merge Sort is $T(n) = O(n \log_2(n))$ if n is a power of two.

In general, if we do not assume anything about the value of c , we find $T(n) \leq 2c n \log_2(n)$.

So the running time of Merge Sort is $T(n) = O(n \log_2(n))$ if n is a power of two.

For a general n , we have

$$T(n) \leq \begin{cases} c & \text{if } n = 1, \\ T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + c \cdot n & \text{if } n \geq 2, \end{cases}$$

In general, if we do not assume anything about the value of c , we find $T(n) \leq 2c n \log_2(n)$.

So the running time of Merge Sort is $T(n) = O(n \log_2(n))$ if n is a power of two.

For a general n , we have

$$T(n) \leq \begin{cases} c & \text{if } n = 1, \\ T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + c \cdot n & \text{if } n \geq 2, \end{cases}$$

By induction, we can prove that $T(n) = O(n \log_2(n))$.