

# Chapter 6: $P$ vs $NP$

## §6.1 Introduction

An algorithm has *polynomial running time* if there is a constant  $c \geq 1$  such that for every input of length  $n$ , the algorithm takes  $O(n^c)$  time.

## §6.1 Introduction

An algorithm has *polynomial running time* if there is a constant  $c \geq 1$  such that for every input of length  $n$ , the algorithm takes  $O(n^c)$  time.

An algorithm has *exponential running time* if there is a constant  $c \geq 1$  such that for every input of length  $n$ , the algorithm takes  $O(2^{n^c})$  time.

## §6.1 Introduction

An algorithm has *polynomial running time* if there is a constant  $c \geq 1$  such that for every input of length  $n$ , the algorithm takes  $O(n^c)$  time.

An algorithm has *exponential running time* if there is a constant  $c \geq 1$  such that for every input of length  $n$ , the algorithm takes  $O(2^{n^c})$  time.

Intuitively speaking,

polynomial means good, fast, efficient, easy, ...

exponential means bad, slow, “try all possible solutions”, difficult, ...

# Complexity Class $P$

A *decision problem* is a problem for which the answer is “yes” or “no”.

$P$  = set of all decision problems that can be solved in polynomial time

# Complexity Class $P$

A *decision problem* is a problem for which the answer is “yes” or “no”.

$P$  = set of all decision problems that can be solved in polynomial time

Examples of problems that are in  $P$ :

- Is a given input graph connected?

# Complexity Class $P$

A *decision problem* is a problem for which the answer is “yes” or “no”.

$P$  = set of all decision problems that can be solved in polynomial time

Examples of problems that are in  $P$ :

- Is a given input graph connected?
- Is a given input graph bipartite?

# Complexity Class $P$

A *decision problem* is a problem for which the answer is “yes” or “no”.

$P$  = set of all decision problems that can be solved in polynomial time

Examples of problems that are in  $P$ :

- Is a given input graph connected?
- Is a given input graph bipartite?
- Is a given input sequence sorted?



# Complexity Class $P$

A *decision problem* is a problem for which the answer is “yes” or “no”.

$P$  = set of all decision problems that can be solved in polynomial time

Examples of problems that are in  $P$ :


- Is a given input graph connected?
- Is a given input graph bipartite?
- Is a given input sequence sorted?
- Does a given input graph contain an Euler cycle? An *Euler cycle* is a cycle that traverses each edge exactly once.

# Other Problems

## HAM-CYCLE

**input:** An undirected graph  $G = (V, E)$  stored using adjacency lists.

**question:** Does  $G$  contain a Hamiltonian cycle? A *Hamiltonian cycle* is a cycle that traverses each vertex exactly once.




# Other Problems

## HAM-CYCLE

**input:** An undirected graph  $G = (V, E)$  stored using adjacency lists.

**question:** Does  $G$  contain a Hamiltonian cycle? A *Hamiltonian cycle* is a cycle that traverses each vertex exactly once.




# Other Problems

## HAM-CYCLE

**input:** An undirected graph  $G = (V, E)$  stored using adjacency lists.

**question:** Does  $G$  contain a Hamiltonian cycle? A *Hamiltonian cycle* is a cycle that traverses each vertex exactly once.



Not known if this problem is in *P*!

# Other Problems

## Traveling Salesman Problem (TSP)

### input:

- A complete directed graph  $G = (V, E)$ , where each edge  $(u, v) \in E$  has a weight  $wt(u, v) > 0$ .
- An integer  $K$ .

**question:** Does  $G$  contain a Hamiltonian cycle of total weight at most  $K$ ?


# Other Problems

## Traveling Salesman Problem (TSP)

### input:

- A complete directed graph  $G = (V, E)$ , where each edge  $(u, v) \in E$  has a weight  $wt(u, v) > 0$ .
- An integer  $K$ .

**question:** Does  $G$  contain a Hamiltonian cycle of total weight at most  $K$ ?




# Other Problems

## Traveling Salesman Problem (TSP)

### input:

- A complete directed graph  $G = (V, E)$ , where each edge  $(u, v) \in E$  has a weight  $wt(u, v) > 0$ .
- An integer  $K$ .

**question:** Does  $G$  contain a Hamiltonian cycle of total weight at most  $K$ ?



Not known if this problem is in  $P$ !

# Other Problems

## SUBSET-SUM

**input:** A set  $S$  of integers together with an integer  $t$

**question:** Is there a subset  $S'$  of  $S$  such that

$$\sum_{x \in S'} x = t?$$



# Other Problems

## SUBSET-SUM

**input:** A set  $S$  of integers together with an integer  $t$

**question:** Is there a subset  $S'$  of  $S$  such that

$$\sum_{x \in S'} x = t?$$

Example:

$$S = \{5, 7, 56, 23, 2902\}$$

$$t = 2963$$

# Other Problems

## SUBSET-SUM

**input:** A set  $S$  of integers together with an integer  $t$

**question:** Is there a subset  $S'$  of  $S$  such that

$$\sum_{x \in S'} x = t?$$

Example:

$$S = \{5, 7, 56, 23, 2902\}$$

$$t = 2963$$

Yes!  $S' = \{5, 56, 2902\}$

# Other Problems

## SUBSET-SUM

**input:** A set  $S$  of integers together with an integer  $t$

**question:** Is there a subset  $S'$  of  $S$  such that

$$\sum_{x \in S'} x = t?$$

Example:

$$S = \{5, 7, 56, 23, 2902\}$$

$$t = 2963$$

$$\text{Yes! } S' = \{5, 56, 2902\}$$


Not known if this problem is in  $P$ !

# Other Problems

## CLIQUE

**input:** An undirected graph  $G = (V, E)$  together with an integer  $K$

**question:** Does  $G$  contain a clique (complete subgraph) with  $K$  vertices?




$$K = 4$$

# Other Problems

## CLIQUE

**input:** An undirected graph  $G = (V, E)$  together with an integer  $K$

**question:** Does  $G$  contain a clique (complete subgraph) with  $K$  vertices?




$$K = 4$$

# Other Problems

## CLIQUE

**input:** An undirected graph  $G = (V, E)$  together with an integer  $K$

**question:** Does  $G$  contain a clique (complete subgraph) with  $K$  vertices?



Not known if this problem is in  $P$ !

# Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.

# Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.
- If the answer to the question is YES, then
  - There is a “short” proof for this.



# Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.
- If the answer to the question is YES, then
  - There is a “short” proof for this.

*Here, “short” means the length of the proof is “polynomial in the length of the input”.*

# Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.
- If the answer to the question is YES, then
  - There is a “short” proof for this.

*Here, “short” means the length of the proof is “polynomial in the length of the input”.*

- If someone gives us such a short proof, then we can “easily” verify this proof.

# Other Problems

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.
- If the answer to the question is YES, then
  - There is a “short” proof for this.

*Here, “short” means the length of the proof is “polynomial in the length of the input”.*

- If someone gives us such a short proof, then we can “easily” verify this proof.

*Here, “easily” means “in polynomial time”.*

# HAM-CYCLE

proof/solution: sequence  $v_1, \dots, v_k$  of vertices.

# HAM-CYCLE

proof/solution: sequence  $v_1, \dots, v_k$  of vertices.

verification:    •  $k = n?$

# HAM-CYCLE

proof/solution: sequence  $v_1, \dots, v_k$  of vertices.

verification:

- $k = n$ ?
- no duplicates in  $v_1, \dots, v_k$ ?

# HAM-CYCLE

proof/solution: sequence  $v_1, \dots, v_k$  of vertices.

- verification:
- $k = n$ ?
  - no duplicates in  $v_1, \dots, v_k$ ?
  - For each  $1 \leq i < k - 1$ ,  $\{v_i, v_{i+1}\}$  is an edge?

# HAM-CYCLE

proof/solution: sequence  $v_1, \dots, v_k$  of vertices.

- verification:
- $k = n$ ?
  - no duplicates in  $v_1, \dots, v_k$ ?
  - For each  $1 \leq i < k - 1$ ,  $\{v_i, v_{i+1}\}$  is an edge?
  - $\{v_k, v_1\}$  is an edge?



# TSP

proof/solution: sequence  $v_1, \dots, v_k$  of vertices.

# TSP

proof/solution: sequence  $v_1, \dots, v_k$  of vertices.

verification:    •  $k = n?$

# TSP

proof/solution: sequence  $v_1, \dots, v_k$  of vertices.

- verification:
- $k = n$ ?
  - no duplicates in  $v_1, \dots, v_k$ ?

# TSP

proof/solution: sequence  $v_1, \dots, v_k$  of vertices.

- verification:
- $k = n$ ?
  - no duplicates in  $v_1, \dots, v_k$ ?
  - Do we have

$$\sum_{i=1}^{k-1} wt(v_i, v_{i+1}) + wt(v_k, v_1) \leq K?$$

# SUBSET-SUM

proof/solution: A set  $S'$ .

# SUBSET-SUM

proof/solution: A set  $S'$ .

verification:    • Is  $S'$  a subset of  $S$ ?

# SUBSET-SUM

proof/solution: A set  $S'$ .

verification:

- Is  $S'$  a subset of  $S$ ?
- Do we have

$$\sum_{x \in S'} x = t?$$

# CLIQUE

proof/solution: A set  $V'$ .



# CLIQUE

proof/solution: A set  $V'$ .

verification: 

- Is  $V'$  a subset of  $V$ ?

# CLIQUE

proof/solution: A set  $V'$ .

- verification:
- Is  $V'$  a subset of  $V$ ?
  - Do we have  $|V'| = K$ ?

# CLIQUE

proof/solution: A set  $V'$ .

- verification:
- Is  $V'$  a subset of  $V$ ?
  - Do we have  $|V'| = K$ ?
  - For each  $u, v \in V'$  such that  $u \neq v$ , is  $\{u, v\}$  an edge in  $E$ ?

For each of the 4 previous problems,

- Not known if it can be solved in polynomial time.
- If the answer to the question is YES, then
  - There is a “short” proof for this.

*Here, “short” means the length of the proof is “polynomial in the length of the input”.*

- If someone gives us such a short proof, then we can “easily” verify this proof.

*Here, “easily” means “in polynomial time”.*

# Complexity Class $NP$

A decision problem  $A$  is in  $NP$  if

- If for a given input  $I$ , the answer to the question  $A(I)$  is YES, then there exists a proof/solution/certificate  $C$  such that
  - $C$  is short (polynomial size in the length of  $I$ )
  - In polynomial time, we can verify that  $C$  is a correct proof for the fact that  $A(I) = YES$ .

# Complexity Class $NP$

A decision problem  $A$  is in  $NP$  if

- If for a given input  $I$ , the answer to the question  $A(I)$  is YES, then there exists a proof/solution/certificate  $C$  such that
  - $C$  is short (polynomial size in the length of  $I$ )
  - In polynomial time, we can verify that  $C$  is a correct proof for the fact that  $A(I) = \text{YES}$ .

$NP$  stands for *Nondeterministic Polynomial*.

# Complexity Class $NP$

A decision problem  $A$  is in  $NP$  if

- If for a given input  $I$ , the answer to the question  $A(I)$  is YES, then there exists a proof/solution/certificate  $C$  such that
  - $C$  is short (polynomial size in the length of  $I$ )
  - In polynomial time, we can verify that  $C$  is a correct proof for the fact that  $A(I) = \text{YES}$ .

$NP$  stands for *Nondeterministic Polynomial*.

The following problems are in  $NP$ :

HAM-CYCLE, TSP, SUBSET-SUM, CLIQUE