

IMO Shortlist 2013

— Algebra

A1 Let n be a positive integer and let a_1, \dots, a_{n-1} be arbitrary real numbers. Define the sequences u_0, \dots, u_n and v_0, \dots, v_n inductively by $u_0 = u_1 = v_0 = v_1 = 1$, and $u_{k+1} = u_k + a_k u_{k-1}$, $v_{k+1} = v_k + a_{n-k} v_{k-1}$ for $k = 1, \dots, n-1$.
Prove that $u_n = v_n$.

A2 Prove that in any set of 2000 distinct real numbers there exist two pairs $a > b$ and $c > d$ with $a \neq c$ or $b \neq d$, such that

$$\left| \frac{a-b}{c-d} - 1 \right| < \frac{1}{100000}.$$

A3 Let $\mathbb{Q}_{>0}$ be the set of all positive rational numbers. Let $f : \mathbb{Q}_{>0} \rightarrow \mathbb{R}$ be a function satisfying the following three conditions:

- (i) for all $x, y \in \mathbb{Q}_{>0}$, we have $f(x)f(y) \geq f(xy)$;
- (ii) for all $x, y \in \mathbb{Q}_{>0}$, we have $f(x+y) \geq f(x) + f(y)$;
- (iii) there exists a rational number $a > 1$ such that $f(a) = a$.

Prove that $f(x) = x$ for all $x \in \mathbb{Q}_{>0}$.

Proposed by Bulgaria

A4 Let n be a positive integer, and consider a sequence a_1, a_2, \dots, a_n of positive integers. Extend it periodically to an infinite sequence a_1, a_2, \dots by defining $a_{n+i} = a_i$ for all $i \geq 1$. If

$$a_1 \leq a_2 \leq \dots \leq a_n \leq a_1 + n$$

and

$$a_{a_i} \leq n + i - 1 \quad \text{for } i = 1, 2, \dots, n,$$

prove that

$$a_1 + \dots + a_n \leq n^2.$$

- A5** Let $\mathbb{Z}_{\geq 0}$ be the set of all nonnegative integers. Find all the functions $f : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{Z}_{\geq 0}$ satisfying the relation

$$f(f(f(n))) = f(n+1) + 1$$

for all $n \in \mathbb{Z}_{\geq 0}$.

- A6** Let $m \neq 0$ be an integer. Find all polynomials $P(x)$ with real coefficients such that

$$(x^3 - mx^2 + 1)P(x+1) + (x^3 + mx^2 + 1)P(x-1) = 2(x^3 - mx + 1)P(x)$$

for all real number x .

— Combinatorics

- C1** Let n be a positive integer. Find the smallest integer k with the following property; Given any real numbers a_1, \dots, a_d such that $a_1 + a_2 + \dots + a_d = n$ and $0 \leq a_i \leq 1$ for $i = 1, 2, \dots, d$, it is possible to partition these numbers into k groups (some of which may be empty) such that the sum of the numbers in each group is at most 1.

- C2** A configuration of 4027 points in the plane is called Colombian if it consists of 2013 red points and 2014 blue points, and no three of the points of the configuration are collinear. By drawing some lines, the plane is divided into several regions. An arrangement of lines is good for a Colombian configuration if the following two conditions are satisfied:

- i) No line passes through any point of the configuration.
- ii) No region contains points of both colors.

Find the least value of k such that for any Colombian configuration of 4027 points, there is a good arrangement of k lines.

Proposed by *Ivan Guo* from *Australia*.

- C3** A crazy physicist discovered a new kind of particle which he called an imon, after some of them mysteriously appeared in his lab. Some pairs of imons in the lab can be entangled, and each imon can participate in many entanglement relations. The physicist has found a way to perform the following two kinds of operations with these particles, one operation at a time.
- (i) If some imon is entangled with an odd number of other imons in the lab, then the physicist can destroy it.
 - (ii) At any moment, he may double the whole family of imons in the lab by

creating a copy I' of each imon I . During this procedure, the two copies I' and J' become entangled if and only if the original imons I and J are entangled, and each copy I' becomes entangled with its original imon I ; no other entanglements occur or disappear at this moment.

Prove that the physicist may apply a sequence of much operations resulting in a family of imons, no two of which are entangled.

- C4** Let n be a positive integer, and let A be a subset of $\{1, \dots, n\}$. An A -partition of n into k parts is a representation of n as a sum $n = a_1 + \dots + a_k$, where the parts a_1, \dots, a_k belong to A and are not necessarily distinct. The number of different parts in such a partition is the number of (distinct) elements in the set $\{a_1, a_2, \dots, a_k\}$. We say that an A -partition of n into k parts is optimal if there is no A -partition of n into r parts with $r < k$. Prove that any optimal A -partition of n contains at most $\sqrt[3]{6n}$ different parts.

- C5** Let r be a positive integer, and let a_0, a_1, \dots be an infinite sequence of real numbers. Assume that for all nonnegative integers m and s there exists a positive integer $n \in [m+1, m+r]$ such that

$$a_m + a_{m+1} + \dots + a_{m+s} = a_n + a_{n+1} + \dots + a_{n+s}$$

Prove that the sequence is periodic, i.e. there exists some $p \geq 1$ such that $a_{n+p} = a_n$ for all $n \geq 0$.

- C6** In some country several pairs of cities are connected by direct two-way flights. It is possible to go from any city to any other by a sequence of flights. The distance between two cities is defined to be the least possible numbers of flights required to go from one of them to the other. It is known that for any city there are at most 100 cities at distance exactly three from it. Prove that there is no city such that more than 2550 other cities have distance exactly four from it.

- C7** Let $n \geq 3$ be an integer, and consider a circle with $n+1$ equally spaced points marked on it. Consider all labellings of these points with the numbers $0, 1, \dots, n$ such that each label is used exactly once; two such labellings are considered to be the same if one can be obtained from the other by a rotation of the circle. A labelling is called *beautiful* if, for any four labels $a < b < c < d$ with $a+d = b+c$, the chord joining the points labelled a and d does not intersect the chord joining the points labelled b and c . Let M be the number of beautiful labelings, and let N be the number of ordered pairs (x, y) of positive integers such that $x + y \leq n$ and $\gcd(x, y) = 1$. Prove

that

$$M = N + 1.$$

-
- C8** Players A and B play a "painful" game on the real line. Player A has a pot of paint with four units of black ink. A quantity p of this ink suffices to blacken a (closed) real interval of length p . In every round, player A picks some positive integer m and provides $1/2^m$ units of ink from the pot. Player B then picks an integer k and blackens the interval from $k/2^m$ to $(k+1)/2^m$ (some parts of this interval may have been blackened before). The goal of player A is to reach a situation where the pot is empty and the interval $[0, 1]$ is not completely blackened.
Decide whether there exists a strategy for player A to win in a finite number of moves.
-
- Geometry
-
- G1** Let ABC be an acute triangle with orthocenter H , and let W be a point on the side BC , lying strictly between B and C . The points M and N are the feet of the altitudes from B and C , respectively. Denote by ω_1 is the circumcircle of BWN , and let X be the point on ω_1 such that WX is a diameter of ω_1 . Analogously, denote by ω_2 the circumcircle of triangle CWM , and let Y be the point such that WY is a diameter of ω_2 . Prove that X, Y and H are collinear.
Proposed by Warut Suksompong and Potcharapol Suteparuk, Thailand
-
- G2** Let ω be the circumcircle of a triangle ABC . Denote by M and N the midpoints of the sides AB and AC , respectively, and denote by T the midpoint of the arc BC of ω not containing A . The circumcircles of the triangles AMT and ANT intersect the perpendicular bisectors of AC and AB at points X and Y , respectively; assume that X and Y lie inside the triangle ABC . The lines MN and XY intersect at K . Prove that $KA = KT$.
-
- G3** In a triangle ABC , let D and E be the feet of the angle bisectors of angles A and B , respectively. A rhombus is inscribed into the quadrilateral $AEDB$ (all vertices of the rhombus lie on different sides of $AEDB$). Let φ be the non-obtuse angle of the rhombus. Prove that $\varphi \leq \max\{\angle BAC, \angle ABC\}$.
-
- G4** Let ABC be a triangle with $\angle B > \angle C$. Let P and Q be two different points on line AC such that $\angle PBA = \angle QBA = \angle ACB$ and A is located between P and C . Suppose that there exists an interior point D of segment BQ for which
-

$PD = PB$. Let the ray AD intersect the circle ABC at $R \neq A$. Prove that $QB = QR$.

G5 Let $ABCDEF$ be a convex hexagon with $AB = DE$, $BC = EF$, $CD = FA$, and $\angle A - \angle D = \angle C - \angle F = \angle E - \angle B$. Prove that the diagonals AD , BE , and CF are concurrent.

G6 Let the excircle of triangle ABC opposite the vertex A be tangent to the side BC at the point A_1 . Define the points B_1 on CA and C_1 on AB analogously, using the excircles opposite B and C , respectively. Suppose that the circumcentre of triangle $A_1B_1C_1$ lies on the circumcircle of triangle ABC . Prove that triangle ABC is right-angled.

Proposed by Alexander A. Polyansky, Russia

— Number Theory

N1 Let $\mathbb{Z}_{>0}$ be the set of positive integers. Find all functions $f : \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$ such that

$$m^2 + f(n) \mid mf(m) + n$$

for all positive integers m and n .

N2 Assume that k and n are two positive integers. Prove that there exist positive integers m_1, \dots, m_k such that

$$1 + \frac{2^k - 1}{n} = \left(1 + \frac{1}{m_1}\right) \cdots \left(1 + \frac{1}{m_k}\right).$$

Proposed by Japan

N3 Prove that there exist infinitely many positive integers n such that the largest prime divisor of $n^4 + n^2 + 1$ is equal to the largest prime divisor of $(n+1)^4 + (n+1)^2 + 1$.

N4 Determine whether there exists an infinite sequence of nonzero digits a_1, a_2, a_3, \dots and a positive integer N such that for every integer $k > N$, the number $\overline{a_k a_{k-1} \cdots a_1}$ is a perfect square.

N5 Fix an integer $k > 2$. Two players, called Ana and Banana, play the following game of numbers. Initially, some integer $n \geq k$ gets written on the blackboard. Then they take moves in turn, with Ana beginning. A player making a move erases the number m just written on the blackboard and replaces it by some

number m' with $k \leq m' < m$ that is coprime to m . The first player who cannot move anymore loses.

An integer $n \geq k$ is called good if Banana has a winning strategy when the initial number is n , and bad otherwise.

Consider two integers $n, n' \geq k$ with the property that each prime number $p \leq k$ divides n if and only if it divides n' . Prove that either both n and n' are good or both are bad.

N6

Determine all functions $f : \mathbb{Q} \rightarrow \mathbb{Z}$ satisfying

$$f\left(\frac{f(x) + a}{b}\right) = f\left(\frac{x + a}{b}\right)$$

for all $x \in \mathbb{Q}$, $a \in \mathbb{Z}$, and $b \in \mathbb{Z}_{>0}$. (Here, $\mathbb{Z}_{>0}$ denotes the set of positive integers.)

N7

Let ν be an irrational positive number, and let m be a positive integer. A pair of (a, b) of positive integers is called *good* if

$$a \lceil b\nu \rceil - b \lfloor a\nu \rfloor = m.$$

A good pair (a, b) is called *excellent* if neither of the pair $(a - b, b)$ and $(a, b - a)$ is good.

Prove that the number of excellent pairs is equal to the sum of the positive divisors of m .