

IMO Shortlist 2002

— Geometry

1 Let B be a point on a circle S_1 , and let A be a point distinct from B on the tangent at B to S_1 . Let C be a point not on S_1 such that the line segment AC meets S_1 at two distinct points. Let S_2 be the circle touching AC at C and touching S_1 at a point D on the opposite side of AC from B . Prove that the circumcentre of triangle BCD lies on the circumcircle of triangle ABC .

2 Let ABC be a triangle for which there exists an interior point F such that $\angle AFB = \angle BFC = \angle CFA$. Let the lines BF and CF meet the sides AC and AB at D and E respectively. Prove that

$$AB + AC \geq 4DE.$$

3 The circle S has centre O , and BC is a diameter of S . Let A be a point of S such that $\angle AOB < 120^\circ$. Let D be the midpoint of the arc AB which does not contain C . The line through O parallel to DA meets the line AC at I . The perpendicular bisector of OA meets S at E and at F . Prove that I is the incentre of the triangle CEF .

4 Circles S_1 and S_2 intersect at points P and Q . Distinct points A_1 and B_1 (not at P or Q) are selected on S_1 . The lines A_1P and B_1P meet S_2 again at A_2 and B_2 respectively, and the lines A_1B_1 and A_2B_2 meet at C . Prove that, as A_1 and B_1 vary, the circumcentres of triangles A_1A_2C all lie on one fixed circle.

5 For any set S of five points in the plane, no three of which are collinear, let $M(S)$ and $m(S)$ denote the greatest and smallest areas, respectively, of triangles determined by three points from S . What is the minimum possible value of $M(S)/m(S)$?

6 Let $n \geq 3$ be a positive integer. Let $C_1, C_2, C_3, \dots, C_n$ be unit circles in the plane, with centres $O_1, O_2, O_3, \dots, O_n$ respectively. If no line meets more than two of the circles, prove that

$$\sum_{1 \leq i < j \leq n} \frac{1}{O_i O_j} \leq \frac{(n-1)\pi}{4}.$$

7 The incircle Ω of the acute-angled triangle ABC is tangent to its side BC at a point K . Let AD be an altitude of triangle ABC , and let M be the midpoint of the segment AD . If N is the common point of the circle Ω and the line KM (distinct from K), then prove that the incircle Ω and the circumcircle of triangle BCN are tangent to each other at the point N .

8 Let two circles S_1 and S_2 meet at the points A and B . A line through A meets S_1 again at C and S_2 again at D . Let M, N, K be three points on the line segments CD, BC, BD respectively, with MN parallel to BD and MK parallel to BC . Let E and F be points on those arcs BC of S_1 and BD of S_2 respectively that do not contain A . Given that EN is perpendicular to BC and FK is perpendicular to BD prove that $\angle EMF = 90^\circ$.

— Number Theory

1 What is the smallest positive integer t such that there exist integers x_1, x_2, \dots, x_t with

$$x_1^3 + x_2^3 + \dots + x_t^3 = 2002^{2002} ?$$

2 Let $n \geq 2$ be a positive integer, with divisors $1 = d_1 < d_2 < \dots < d_k = n$. Prove that $d_1 d_2 + d_2 d_3 + \dots + d_{k-1} d_k$ is always less than n^2 , and determine when it is a divisor of n^2 .

3 Let p_1, p_2, \dots, p_n be distinct primes greater than 3. Show that $2^{p_1 p_2 \dots p_n} + 1$ has at least 4^n divisors.

4 Is there a positive integer m such that the equation

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{abc} = \frac{m}{a+b+c}$$

has infinitely many solutions in positive integers a, b, c ?

5 Let $m, n \geq 2$ be positive integers, and let a_1, a_2, \dots, a_n be integers, none of which is a multiple of m^{n-1} . Show that there exist integers e_1, e_2, \dots, e_n , not all zero, with $|e_i| < m$ for all i , such that $e_1 a_1 + e_2 a_2 + \dots + e_n a_n$ is a multiple of m^n .

6 Find all pairs of positive integers $m, n \geq 3$ for which there exist infinitely many positive integers a such that

$$\frac{a^m + a - 1}{a^n + a^2 - 1}$$

is itself an integer.

Laurentiu Panaitopol, Romania

— Algebra

1 Find all functions f from the reals to the reals such that

$$f(f(x) + y) = 2x + f(f(y) - x)$$

for all real x, y .

2 Let a_1, a_2, \dots be an infinite sequence of real numbers, for which there exists a real number c with $0 \leq a_i \leq c$ for all i , such that

$$|a_i - a_j| \geq \frac{1}{i + j} \quad \text{for all } i, j \text{ with } i \neq j.$$

Prove that $c \geq 1$.

3 Let P be a cubic polynomial given by $P(x) = ax^3 + bx^2 + cx + d$, where a, b, c, d are integers and $a \neq 0$. Suppose that $xP(x) = yP(y)$ for infinitely many pairs x, y of integers with $x \neq y$. Prove that the equation $P(x) = 0$ has an integer root.

4 Find all functions f from the reals to the reals such that

$$(f(x) + f(z))(f(y) + f(t)) = f(xy - zt) + f(xt + yz)$$

for all real x, y, z, t .

5 Let n be a positive integer that is not a perfect cube. Define real numbers a, b, c by

$$a = \sqrt[3]{n}, \quad b = \frac{1}{a - [a]}, \quad c = \frac{1}{b - [b]},$$

where $[x]$ denotes the integer part of x . Prove that there are infinitely many such integers n with the property that there exist integers r, s, t , not all zero, such that $ra + sb + tc = 0$.

- 6 Let A be a non-empty set of positive integers. Suppose that there are positive integers b_1, \dots, b_n and c_1, \dots, c_n such that

- for each i the set $b_i A + c_i = \{b_i a + c_i : a \in A\}$ is a subset of A , and
- the sets $b_i A + c_i$ and $b_j A + c_j$ are disjoint whenever $i \neq j$

Prove that

$$\frac{1}{b_1} + \dots + \frac{1}{b_n} \leq 1.$$

– Combinatorics

- 1 Let n be a positive integer. Each point (x, y) in the plane, where x and y are non-negative integers with $x + y < n$, is coloured red or blue, subject to the following condition: if a point (x, y) is red, then so are all points (x', y') with $x' \leq x$ and $y' \leq y$. Let A be the number of ways to choose n blue points with distinct x -coordinates, and let B be the number of ways to choose n blue points with distinct y -coordinates. Prove that $A = B$.

- 2 For n an odd positive integer, the unit squares of an $n \times n$ chessboard are coloured alternately black and white, with the four corners coloured black. A tromino is an L -shape formed by three connected unit squares. For which values of n is it possible to cover all the black squares with non-overlapping trominos? When it is possible, what is the minimum number of trominos needed?

- 3 Let n be a positive integer. A sequence of n positive integers (not necessarily distinct) is called **full** if it satisfies the following condition: for each positive integer $k \geq 2$, if the number k appears in the sequence then so does the number $k - 1$, and moreover the first occurrence of $k - 1$ comes before the last occurrence of k . For each n , how many full sequences are there ?

- 4 Let T be the set of ordered triples (x, y, z) , where x, y, z are integers with $0 \leq x, y, z \leq 9$. Players A and B play the following guessing game. Player A chooses a triple (x, y, z) in T , and Player B has to discover A 's triple in as few moves as possible. A *move* consists of the following: B gives A a triple (a, b, c) in T , and A replies by giving B the number $|x + y - a - b| + |y + z - b - c| + |z + x - c - a|$. Find the minimum number of moves that B needs to be sure of determining A 's triple.



Art of Problem Solving

2002 IMO Shortlist

- 5 Let $r \geq 2$ be a fixed positive integer, and let F be an infinite family of sets, each of size r , no two of which are disjoint. Prove that there exists a set of size $r - 1$ that meets each set in F .
-
- 6 Let n be an even positive integer. Show that there is a permutation (x_1, x_2, \dots, x_n) of $(1, 2, \dots, n)$ such that for every $i \in \{1, 2, \dots, n\}$, the number x_{i+1} is one of the numbers $2x_i, 2x_i - 1, 2x_i - n, 2x_i - n - 1$. Hereby, we use the cyclic subscript convention, so that x_{n+1} means x_1 .
-
- 7 Among a group of 120 people, some pairs are friends. A *weak quartet* is a set of four people containing exactly one pair of friends. What is the maximum possible number of weak quartets ?
-