

USAMO 2003

Day 1

April 29th

1 Prove that for every positive integer n there exists an n -digit number divisible by 5^n all of whose digits are odd.

2 A convex polygon \mathcal{P} in the plane is dissected into smaller convex polygons by drawing all of its diagonals. The lengths of all sides and all diagonals of the polygon \mathcal{P} are rational numbers. Prove that the lengths of all sides of all polygons in the dissection are also rational numbers.

3 Let $n \neq 0$. For every sequence of integers

$$A = a_0, a_1, a_2, \dots, a_n$$

satisfying $0 \leq a_i \leq i$, for $i = 0, \dots, n$, define another sequence

$$t(A) = t(a_0), t(a_1), t(a_2), \dots, t(a_n)$$

by setting $t(a_i)$ to be the number of terms in the sequence A that precede the term a_i and are different from a_i . Show that, starting from any sequence A as above, fewer than n applications of the transformation t lead to a sequence B such that $t(B) = B$.

Day 2

April 30th

4 Let ABC be a triangle. A circle passing through A and B intersects segments AC and BC at D and E , respectively. Lines AB and DE intersect at F , while lines BD and CF intersect at M . Prove that $MF = MC$ if and only if $MB \cdot MD = MC^2$.

5 Let a, b, c be positive real numbers. Prove that

$$\frac{(2a + b + c)^2}{2a^2 + (b + c)^2} + \frac{(2b + c + a)^2}{2b^2 + (c + a)^2} + \frac{(2c + a + b)^2}{2c^2 + (a + b)^2} \leq 8.$$

6 At the vertices of a regular hexagon are written six nonnegative integers whose sum is 2003^{2003} . Bert is allowed to make moves of the following form: he may pick a vertex and replace the number written there by the absolute value of the



Art of Problem Solving

2003 USAMO

difference between the numbers written at the two neighboring vertices. Prove that Bert can make a sequence of moves, after which the number 0 appears at all six vertices.



—

These problems are copyright © Mathematical Association of America (<http://maa.org>).
