



Art of Problem Solving

2010 USA Team Selection Test

USA Team Selection Test 2010

Day 1

- 1 Let P be a polynomial with integer coefficients such that $P(0) = 0$ and

$$\gcd(P(0), P(1), P(2), \dots) = 1.$$

Show there are infinitely many n such that

$$\gcd(P(n) - P(0), P(n+1) - P(1), P(n+2) - P(2), \dots) = n.$$

-
- 2 Let a, b, c be positive reals such that $abc = 1$. Show that

$$\frac{1}{a^5(b+2c)^2} + \frac{1}{b^5(c+2a)^2} + \frac{1}{c^5(a+2b)^2} \geq \frac{1}{3}.$$

-
- 3 Let h_a, h_b, h_c be the lengths of the altitudes of a triangle ABC from A, B, C respectively. Let P be any point inside the triangle. Show that

$$\frac{PA}{h_b + h_c} + \frac{PB}{h_a + h_c} + \frac{PC}{h_a + h_b} \geq 1.$$

Day 2

- 4 Let ABC be a triangle. Point M and N lie on sides AC and BC respectively such that $MN \parallel AB$. Points P and Q lie on sides AB and CB respectively such that $PQ \parallel AC$. The incircle of triangle CMN touches segment AC at E . The incircle of triangle BPQ touches segment AB at F . Line EN and AB meet at R , and lines FQ and AC meet at S . Given that $AE = AF$, prove that the incenter of triangle AEF lies on the incircle of triangle ARS .

-
- 5 Define the sequence a_1, a_2, a_3, \dots by $a_1 = 1$ and, for $n > 1$,

$$a_n = a_{\lfloor n/2 \rfloor} + a_{\lfloor n/3 \rfloor} + \dots + a_{\lfloor n/n \rfloor} + 1.$$

Prove that there are infinitely many n such that $a_n \equiv n \pmod{2^{2010}}$.

- 6 Let T be a finite set of positive integers greater than 1. A subset S of T is called *good* if for every $t \in T$ there exists some $s \in S$ with $\gcd(s, t) > 1$. Prove that the number of good subsets of T is odd.

Day 3

- 7 In triangle ABC , let P and Q be two interior points such that $\angle ABP = \angle QBC$ and $\angle ACP = \angle QCB$. Point D lies on segment BC . Prove that $\angle APB + \angle DPC = 180^\circ$ if and only if $\angle AQC + \angle DQB = 180^\circ$.

- 8 Let m, n be positive integers with $m \geq n$, and let S be the set of all n -term sequences of positive integers (a_1, a_2, \dots, a_n) such that $a_1 + a_2 + \dots + a_n = m$. Show that

$$\sum_S 1^{a_1} 2^{a_2} \dots n^{a_n} = \binom{n}{n} n^m - \binom{n}{n-1} (n-1)^m + \dots + (-1)^{n-2} \binom{n}{2} 2^m + (-1)^{n-1} \binom{n}{1}.$$

- 9 Determine whether or not there exists a positive integer k such that $p = 6k + 1$ is a prime and

$$\binom{3k}{k} \equiv 1 \pmod{p}.$$



These problems are copyright © Mathematical Association of America (<http://maa.org>).