

India
International Mathematical Olympiad Training Camp
2007

Day 1

- [1] Show that in a non-equilateral triangle, the following statements are equivalent: (a) The angles of the triangle are in arithmetic progression. (b) The common tangent to the Nine-point circle and the Incircle is parallel to the Euler Line.
- [2] Prove that the equation $\frac{x^7-1}{x-1} = y^5 - 1$ doesn't have integer solutions!
- [3] Let \mathbb{X} be the set of all bijective functions from the set $S = \{1, 2, \dots, n\}$ to itself. For each $f \in \mathbb{X}$, define

$$T_f(j) = \begin{cases} 1, & \text{if } f^{(12)}(j) = j, \\ 0, & \text{otherwise} \end{cases}$$

Determine $\sum_{f \in \mathbb{X}} \sum_{j=1}^n T_f(j)$. (Here $f^{(k)}(x) = f(f^{(k-1)}(x))$ for all $k \geq 2$.)

India
International Mathematical Olympiad Training Camp
2007

Day 2

- [1] Let $ABCD$ be a trapezoid with parallel sides $AB > CD$. Points K and L lie on the line segments AB and CD , respectively, so that $\frac{AK}{KB} = \frac{DL}{LC}$. Suppose that there are points P and Q on the line segment KL satisfying $\angle APB = \angle BCD$ and $\angle CQD = \angle ABC$. Prove that the points P , Q , B and C are concyclic.

Proposed by Vyacheslav Yasinskiy, Ukraine

- [2] Let a, b, c be non-negative real numbers such that $a + b \leq c + 1$, $b + c \leq a + 1$ and $c + a \leq b + 1$. Show that

$$a^2 + b^2 + c^2 \leq 2abc + 1.$$

- [3] Given a finite string S of symbols X and O , we denote $\Delta(s)$ as the number of X 's in S minus the number of O 's (For example, $\Delta(XOOXOOX) = -1$). We call a string S **balanced** if every sub-string T of (consecutive symbols) S has the property $-1 \leq \Delta(T) \leq 2$. (Thus $XOOXOOX$ is not balanced, since it contains the sub-string $OOXOO$ whose Δ value is -3 . Find, with proof, the number of balanced strings of length n .

India
International Mathematical Olympiad Training Camp
2007

Day 3

- 1 a_0, a_1, a_2, \dots is a sequence of real numbers such that

$$a_{n+1} = [a_n] \cdot \{a_n\}$$

prove that exist j such that for every $i \geq j$ we have $a_{i+2} = a_i$.

Proposed by Harmel Nestra, Estonia

- 2 Let S be a finite set of points in the plane such that no three of them are on a line. For each convex polygon P whose vertices are in S , let $a(P)$ be the number of vertices of P , and let $b(P)$ be the number of points of S which are outside P . A line segment, a point, and the empty set are considered as convex polygons of 2, 1, and 0 vertices respectively. Prove that for every real number x :

$\sum_P x^{a(P)}(1-x)^{b(P)} = 1$, where the sum is taken over all convex polygons with vertices in S .

Alternative formulation:

Let M be a finite point set in the plane and no three points are collinear. A subset A of M will be called round if its elements is the set of vertices of a convex A -gon $V(A)$. For each round subset let $r(A)$ be the number of points from M which are exterior from the convex A -gon $V(A)$. Subsets with 0, 1 and 2 elements are always round, its corresponding polygons are the empty set, a point or a segment, respectively (for which all other points that are not vertices of the polygon are exterior). For each round subset A of M construct the polynomial

$$P_A(x) = x^{|A|}(1-x)^{r(A)}.$$

Show that the sum of polynomials for all round subsets is exactly the polynomial $P(x) = 1$.

Proposed by Federico Ardila, Colombia

India
International Mathematical Olympiad Training Camp
2007

Day 4

- [1] Circles w_1 and w_2 with centres O_1 and O_2 are externally tangent at point D and internally tangent to a circle w at points E and F respectively. Line t is the common tangent of w_1 and w_2 at D . Let AB be the diameter of w perpendicular to t , so that A, E, O_1 are on the same side of t . Prove that lines AO_1 , BO_2 , EF and t are concurrent.
- [2] Find all integer solutions (x, y) of the equation $y^2 = x^3 - p^2x$, where p is a prime such that $p \equiv 3 \pmod{4}$.
- [3] Find all function(s) $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfying the equation

$$f(x+y) + f(x)f(y) = (1+y)f(x) + (1+x)f(y) + f(xy);$$

For all $x, y \in \mathbb{R}$.