

APMO 2009

- 1** Consider the following operation on positive real numbers written on a blackboard:
Choose a number r written on the blackboard, erase that number, and then write a pair of positive real numbers a and b satisfying the condition $2r^2 = ab$ on the board.
- Assume that you start out with just one positive real number r on the blackboard, and apply this operation $k^2 - 1$ times to end up with k^2 positive real numbers, not necessarily distinct. Show that there exists a number on the board which does not exceed kr .
-
- 2** Let a_1, a_2, a_3, a_4, a_5 be real numbers satisfying the following equations:

$$\frac{a_1}{k^2+1} + \frac{a_2}{k^2+2} + \frac{a_3}{k^2+3} + \frac{a_4}{k^2+4} + \frac{a_5}{k^2+5} = \frac{1}{k^2} \text{ for } k = 1, 2, 3, 4, 5$$
Find the value of $\frac{a_1}{37} + \frac{a_2}{38} + \frac{a_3}{39} + \frac{a_4}{40} + \frac{a_5}{41}$ (Express the value in a single fraction.)
-
- 3** Let three circles $\Gamma_1, \Gamma_2, \Gamma_3$, which are non-overlapping and mutually external, be given in the plane. For each point P in the plane, outside the three circles, construct six points $A_1, B_1, A_2, B_2, A_3, B_3$ as follows: For each $i = 1, 2, 3$, A_i, B_i are distinct points on the circle Γ_i such that the lines PA_i and PB_i are both tangents to Γ_i . Call the point P exceptional if, from the construction, three lines A_1B_1, A_2B_2, A_3B_3 are concurrent. Show that every exceptional point of the plane, if exists, lies on the same circle.
-
- 4** Prove that for any positive integer k , there exists an arithmetic sequence $\frac{a_1}{b_1}, \frac{a_2}{b_2}, \frac{a_3}{b_3}, \dots, \frac{a_k}{b_k}$ of rational numbers, where a_i, b_i are relatively prime positive integers for each $i = 1, 2, \dots, k$ such that the positive integers $a_1, b_1, a_2, b_2, \dots, a_k, b_k$ are all distinct.
-
- 5** Larry and Rob are two robots travelling in one car from Argovia to Zillis. Both robots have control over the steering and steer according to the following algorithm: Larry makes a 90 degrees left turn after every ℓ kilometer driving from start, Rob makes a 90 degrees right turn after every r kilometer driving from start, where ℓ and r are relatively prime positive integers.
- In the event of both turns occurring simultaneously, the car will keep going without changing direction. Assume that the ground is flat and the car can move in any direction. Let the car start from Argovia facing towards Zillis. For



Art of Problem Solving

2009 APMO

which choices of the pair (ℓ, r) is the car guaranteed to reach Zillis, regardless of how far it is from Argovia?
