



Sorbonne Université

École doctorale Physique en Île-de-France

EDPIF n°564

Laboratoire de Physique Nucléaire et des Hautes Energies

Towards precision cosmology with Type Ia supernova peculiar velocities: from SN modeling to field level inference

Par Mahmoud AHMED EMAM OSMAN

Thèse de doctorat de Cosmologie Physique

Dirigée par Nicolas REGNAULT
et Pauline ZARROUK

Présentée et soutenue publiquement le *(XX Septembre 2026)*

Devant un jury composé de:

Guilhem Lavaux, *Directeur de recherche, Institut d'Astrophysique de Paris, Chef de Jury - Rapporteur*

Anne Ealet, *Directrice de recherche, Institut de Physique des 2 Infinis, Rapportrice*

Julian Bautista, *Professeur, Centre de Physique des Particules de Marseille, Examinateur*

Ariel Goobar, *Professeur, établissement, Examinateur*

Céline Combet, *Directrice de recherche, Laboratoire de Physique Subatomique et Cosmologie, Examinatrice*

Nicolas Regnault, *Directeur de recherche, Laboratoire de Physique Nucléaire et des Hautes Energies, Encadrant de thèse*

Invités:

Pauline Zarrouk, *Directrice de recherche, Laboratoire de Physique Nucléaire et des Hautes Energies, Encadrante de thèse*

Contents

Contents	3
1 Introduction	7
1 Introduction to Modern Cosmology	9
1.1 General Relativity	9
1.2 The expanding, homogenous Universe	11
1.3 Friedmann-Lemaitre-Robertson-Walker metric	11
1.4 Friedmann equations	14
1.5 Energy contents of the Universe	14
1.6 The Λ CDM Model	16
2 Large Scale Structure Formation	19
2.1 Newtonian approach to structure formation	19
2.2 Relativistic structure formation	23
2.3 Inflation	24
3 Cosmological distances	27
4 Cosmological Observables	28
4.1 The Cosmic Microwave Background	28
4.2 Galaxy Surveys	30
4.2.1 The Matter Power Spectrum	31
4.2.2 Baryon Acoustic Oscillations	32
4.2.3 Redshift Space Distortions	32
4.3 Type Ia Supernovae	33
4.4 Gravitational Lensing	33
4.5 The Lyman- α Forest	33
5 Tensions in The Cosmological Model	36
5.1 H_0 Tension	36
5.2 S_8 Tension	37
5.3 (w_0, w_a) Tension	37
6 Physical explanations for the accelerated Universe	39
6.1 Cosmological constant	39
6.2 Dynamical dark energy	39
6.3 Quintessence field	39
6.4 Modified Theories of Gravity	40
2 Cosmology with type Ia Supernova	43
1 Type Ia Supernova	44
1.1 Classification	44

1.2	Explosion mechanisms	44
1.3	Explosion rate	47
1.4	Observations	47
2	Cosmology with distances	52
2.1	Hubble diagram	52
2.1.1	Measurement of $\Omega_m - w$	52
2.1.2	Measurement of H_0	53
2.2	Peculiar velocities	53
2.2.1	Extracting peculiar velocities	54
2.2.2	Measurement of $f\sigma_8$	55
3	Distance measurements from observed fluxes	57
4	Standardization	58
4.1	Brighter-slower effect	58
4.2	Brighter-bluer effect	58
4.3	Environmental effects	59
5	Spectrophotometric models of supernovae	61
5.1	SALT2/SALT3	61
5.2	Models trained on the SNfactory dataset	64
5.2.1	SNEMO	65
5.2.2	SUGAR	66
5.2.3	Twins Embedding	67
5.2.4	SuPAErnova	68
6	Main goal of this PhD	69
3	LEMAITRE project	71
1	LEMAITRE: Hubble diagram project overview	72
2	Datasets: ZTF, SNLS, HSC	72
2.1	ZTF survey	72
2.1.1	ZTF photometry	73
2.1.2	Spectroscopic follow up	73
2.1.3	ZTF DR2	74
2.2	SNLS survey	74
2.3	HSC survey	76
3	Pipeline	78
3.1	Pipeline overview	78
3.2	Cosmological analyses with LEMAITRE	78
3.3	Uncertainties in SN measurements?	78
4	DESI survey	79
4.1	Spectroscopic instrument	79
4.2	Tracers	79
4.3	ZTF and DESI Synergies	80
4	NaCl	81
1	NaCl framework	82
1.1	The idea behind the framework	82
1.2	Differences with the SALT models	82
1.2.1	Practical implementation of the SALT2 model	82
1.3	Constraints	84

1.4	Regularization	85
1.5	Error models	86
1.5.1	Error Snake	86
1.5.2	Color Scatter	87
1.5.3	Calibration Uncertainties	87
1.6	NaCl output	87
2	Improving NaCl	90
2.1	Implementing a model for spectrophotometric data	90
2.2	Regularization	90
2.3	NaCl training recipe	92
3	NaCl on real data	94
3.1	NaCl training results	94
3.1.1	Data coverage	94
3.1.2	Fitting the spectra	95
3.1.3	Training residuals	95
5	LEMAITRE Data Challenges	99
1	LEMAÎTRE Data Challenges	100
1.1	General description of the DCs	100
1.2	Georges analysis pipeline	100
1.2.1	PeTS & miniPeTS	100
1.2.2	Data compression	101
1.2.3	NaCl	102
1.2.4	EDRIS	102
1.2.5	Cosmologix	102
1.3	DC1	103
1.4	DC2	103
1.5	DC3	103
1.6	DC4	103
1.7	DC5	104
2	pre-DC1	105
2.1	Data generation : simulation	105
2.2	Goals	106
3	Results : Potential biases & systematics	107
3.1	Model & data agreement	107
3.2	Parameter reconstruction and covariance matrix	107
3.3	Distance modulus and cosmological parameters reconstruction	108
6	Field level inference with peculiar velocities from ZTF SN Ia	113
1	The BORG forward model	114
1.1	The BORG algorithm	114
1.2	Utility of field level inferences	115
1.2.1	BORG's Bayesian inference chain	115
1.2.2	Particle mass assignment	117
1.2.3	Gravity Model	118
1.2.4	Likelihood components of BORG	121
1.2.5	Advantages over summary-statistics analyses	122
1.2.6	Numerical implementation	122

2	Self-consistent simulation analysis	123
7	BORG applications on ZTF simulations	125
1	ZTF peculiar velocity data challenge (DC PV)	126
2	Supernova forward-model analysis	127
2.1	Bias model	127
3	Results of the field level inference	128
4	Comparison with other methods for $f\sigma_8$	128
	Bibliography	131

CHAPTER 1

Introduction

Contents

1	Introduction to Modern Cosmology	9
1.1	General Relativity	9
1.2	The expanding, homogenous Universe	11
1.3	Friedmann-Lemaitre-Robertson-Walker metric	11
1.4	Friedmann equations	14
1.5	Energy contents of the Universe	14
1.6	The Λ CDM Model	16
2	Large Scale Structure Formation	19
2.1	Newtonian approach to structure formation	19
2.2	Relativistic structure formation	23
2.3	Inflation	24
3	Cosmological distances	27
4	Cosmological Observables	28
4.1	The Cosmic Microwave Background	28
4.2	Galaxy Surveys	30
4.3	Type Ia Supernovae	33
4.4	Gravitational Lensing	33
4.5	The Lyman- α Forest	33
5	Tensions in The Cosmological Model	36
5.1	H_0 Tension	36
5.2	S_8 Tension	37
5.3	(w_0, w_a) Tension	37
6	Physical explanations for the accelerated Universe	39
6.1	Cosmological constant	39

6.2	Dynamical dark energy	39
6.3	Quintessence field	39
6.4	Modified Theories of Gravity	40

1 Introduction to Modern Cosmology

The quest to understand the natural world and how we are placed in it has existed for many centuries. It has influenced many religious, cultural and scientific ideas throughout time and their evolutions lead us to how we see our Universe today. Physical cosmology (cosmology just from now on) is the study of the nature of our Universe through observations in order to understand its origins, dynamics and components. We consider the beginning of modern cosmology was about a century ago with the discovery of the expansion of the Universe initially predicted by Lemaitre in 1927 ([Lemaître, 1927], [Lemaître, 1931]) and observed by Hubble in 1929 ([Hubble, 1929]). The current cosmological model named Λ CDM (Λ Cold Dark Matter) best describes of our Universe's history, dynamics and components. In this section I will detail the necessary tools that allows us to describe the large scale Universe as we know it today.

1.1 General Relativity

According to modern belief and understanding, the Universe contains four fundamental forces dominating at different scales: at large scales, that force is gravity. The basis of our understanding of modern cosmology is centered around Einstein's gravitational theory: General Relativity (GR) ([Einstein, 1916]), the most tested theory of gravity to date. GR's views on the nature of gravity differs a lot from the classical Newtonian view. In GR, massive objects do not directly exert a force on other massive objects, but instead cause the distortions in spacetime. The theory was established after the success of the theory of Special Relativity, also theorized by Einstein in 1905, where the speed of light was shown to be constant in all observation frames.

Understanding the mathematical foundation of GR requires a couple of fundamental notions. GR treats coordinate transformations in a 4-dimensional space (spacetime) (t, x, y, z) using Lorentz transformations. The formalism will be described using tensorial notations and the Einstein summing convention will also be assumed. We invite the reader to familiarise themselves with these notions as well as an understanding of classical field theory. We will also not be going into a lot of details regarding some mathematical and geometrical notions (like geodesics, covectors, covariant derivatives, etc..) and physical concepts found in the Einstein equations such as the Riemann tensor $R^\mu_{\nu\kappa\lambda}$, Ricci tensor $R_{\mu\nu}$, the Ricci scalar R scalar and Christoffel symbols $\Gamma^\mu_{\nu\lambda}$. If you are interested in learning more on those concepts we recommend David Tong's lectures on General Relativity ¹.

Equivalence principle

Einstein's equivalence principle builds up upon previous results obtained in Newtonian physics. The principle states that the gravitational mass m_g defined due to gravitational interactions is equal to the inertial mass m_i subdued to acceleration as a result of a force. This previously lead to the idea that free falling objects have the same velocity regardless of their masses. Einstein's principle extends this idea to accelerated frames and not just inertial frames. This principle has been heavily tested on Earth and in the Solar System and is yet to be invalidated (REFS).

Einstein equations

¹<https://www.damtp.cam.ac.uk/user/tong/gr.html>

In order to define the equations of motions of any object in GR, the first thing that must be defined is a metric. Physically, a metric is an infinitesimal "separation" that is conserved. In classical physics, the length between two points ΔL would be identical for two observers even if both are moving. In GR this is no longer true and only a combination of a spatial separation and a temporal one becomes invariant. If we write down the spacetime displacement as dx^μ with μ and index taking the values 0 if it's a time displacement and 1, 2, 3 if it's spatial displacement along the x, y, z , we can define the invariant displacement ds^2 as :

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu \quad (1.1)$$

where $g_{\mu\nu}$ is the metric that can depend on the coordinates x^μ . Essentially, the metric expresses the distance between two events in spacetime. An example of a metric is the Minkowski metric used to describe events on Special Relativity and is written as:

$$\eta_{\mu\nu} = \begin{pmatrix} +1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad (1.2)$$

Since in GR spacetime can go through a series of curvatures, we do not expect that an object exhibiting no forces to move in a straight line. What we see instead is that they follow *geodesics*. One can think of geodesics as the straightest possible path in a curved space. The geodesic equation, which corresponds to the equation of motion for a point-like object, is written as:

$$\frac{d^2 x^\mu}{d\lambda^2} = \Gamma_{\alpha\beta}^\mu \frac{dx^\alpha}{d\lambda} \frac{dx^\beta}{d\lambda} \quad (1.3)$$

where $\Gamma_{\alpha\beta}^\mu$ are the Christoffel symbols and λ is the evolution parameter which increases along the particle's path. One can derive the geodesic equation by applying the equivalence principle on a free falling particle that does not accelerate (such that $\frac{d^2 x^\mu}{d\lambda^2} = 0$).

The Einstein field equations ([Einstein, 1916]) close the theory by relating the geometry of spacetime, encoded in the metric, to the distribution of matter and energy within it. In this framework, gravity is not a force but a manifestation of spacetime curvature, and the motion of particles follows the geodesics of that curved geometry. We note $G_{\mu\nu}$ the Einstein tensor describing the geometry and $T_{\mu\nu}$ the energy-momentum tensor describing the energy content, we get the following Einstein equations:

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = \frac{8\pi G}{c^4} T_{\mu\nu} \quad (1.4)$$

where $R_{\mu\nu}$ is the Ricci tensor, R the Ricci scalar, G Newton's gravitational constant and c the speed of light. It's important to note that for a symmetric metric, this represents a set of 10 equations where the coefficients of $g_{\mu\nu}$ are unknowns.

Mathematically, an equivalent way to write the Einstein equations would be:

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} \quad (1.5)$$

where Λ is a constant. Physically however, this can lead to an accelerated expanding Universe (see Section 6.1).

The Einstein equations could also be determined by varying the Einstein-Hilbert action which is:

$$S_{EH} = \frac{c^4}{16\pi G} \int d^4x \sqrt{-g} (R - 2\Lambda) + \int d^4x \sqrt{-g} \mathcal{L}_m \quad (1.6)$$

$$\frac{\delta S_{EH}}{\delta g_{\mu\nu}} = 0 \quad (1.7)$$

where g is the determinant of the metric and $\mathcal{L}_m$ is the Lagrangian density of a field that we include in our theory (such as matter, radiation ...). This is a mathematically rigorous approach to deriving the field equations and it is commonly used when exploring modified theories of gravity (see Section 6.4).

1.2 The expanding, homogenous Universe

Cosmological principle

A key principle in modern cosmology is the cosmological principle, stating that the Universe on large scales is homogenous and isotropic. This is understood in the sense that the Universe does not possess a privileged center and that the structures in it remain identical regardless of the direction of observation. Observations from the SDSS (Sloan Digital Sky Survey [Sarkar et al., 2009]) and DESI (Dark Energy Spectroscopic Instrument [Adame et al., 2025]) have shown strong evidence for homogeneity in the Universe at large scales. We represent this in detail with large scale dark matter simulations such as the Uchuu simulation [Ishiyama et al., 2021] seen in Figure 1.1. We can see that on large scales the matter distribution is homogeneous. CMB measurements from the Planck collaboration ([Akrami et al., 2020]) have shown strong evidence for isotropy as shown in Figure 1.2.

Expanding Universe and Hubble relation

Following the publication of GR, discoveries from [Lemaître, 1927]’s work on the Einstein field equations showed that the proper velocity of a celestial object was proportional to its distance. Two years later, [Hubble, 1929] manages to show that the radial velocities of 32 nearby galaxies increase with their distances indicating an expanding Universe, which is shown in Figure 1.3. The Hubble-Lemaitre relation is established as :

$$v = H_0 d \quad (1.8)$$

where v is the velocity, d the distance of the object and H_0 the Hubble constant. Initially, H_0 was measured to be around $500 \text{ km s}^{-1} \text{ Mpc}^{-1}$, today more precise measurements show that it is around $70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. We may also define a *Hubble diagram* as the diagram that relates the distance of celestial objects from an observer to their velocities.

An expanding Universe entails that objects emitting any kind of waves will be stretched in space making their wavelengths longer. We can call this the **cosmological redshift** z and it is defined as :

$$1 + z = \frac{\nu_{rest}}{\nu_{obs}} = \frac{\lambda_{obs}}{\lambda_{rest}} \quad (1.9)$$

where ν_{rest} and λ_{rest} are the frequency and wavelength emitted at restframe and ν_{obs} and λ_{obs} are the observed ones. Other sources that can the cosmological are the peculiar motion of celestial objects which can shift the emitted colour to either bluer or redder frequencies (see Section 2.2 where we explain the analysis done with peculiar velocities).

1.3 Friedmann-Lemaitre-Robertson-Walker metric

We present here a metric that satisfies the isotropic homogeneity of the Universe as well as an expanding one called Friedmann-Lemaitre-Robertson-Walker (FLRW). It is more convenient to express the metric




Fig. 1.1 Uchuu dark matter simulation of a box $2000\text{Mpc}.h^{-1} \times 1333\text{Mpc}.h^{-1}$ with a thickness of $25\text{Mpc}.h^{-1}$. The blue boxes highlight the homogeneity at large scales while the white boxes highlight the inhomogeneity at some scales due to structure formation. Adapted from [Ishiyama et al., 2021].

in spherical coordinates (t, r, ϕ, θ) . Working in natural units $c = 1$, we obtain:

$$ds^2 = dt^2 - a^2(t) \left[\frac{dr^2}{1 + kr^2} + r^2(d\theta^2 + \sin^2(\theta)d\phi^2) \right] \quad (1.10)$$

where $a(t)$ is a dimensionless parameter called the "scale" factor, defined such that today at t_0 it is equal to 1, and used to describe the expansion which can be written as :

$$H(t) = \frac{\dot{a}(t)}{a(t)} \quad (1.11)$$

where the dot denotes the derivative with respect to time and $H(t)$ the Hubble parameter. k is a global curvature term such that it can take one of three values $0, -1, 1$ to describe a Universe that is flat, open/hyperbolic or closed/spherical respectively. It is important to note that we can obtain the redshift of an object that emitted a wave at a time t_e from the scale factor as the following relation holds true:

$$\frac{a(t_0)}{a(t_e)} = 1 + z \quad (1.12)$$




Fig. 1.2 CMB map provided by ESA and the Planck Collaboration.




Fig. 1.3 Original Hubble diagram ([Hubble, 1929]) portraying the distance velocity diagram of nearby galaxies.

Often we apply a change of coordinates such that any object would maintain its coordinates regardless of the expansion. These are called comoving coordinates. We define the comoving distance χ such that:

$$r = S_k(\chi) = \begin{cases} S_0(\chi) = \chi \\ S_{-1}(\chi) = \sin(\chi) \\ S_1(\chi) = \sinh(\chi) \end{cases} \quad (1.13)$$

$$d\chi = \frac{dr}{\sqrt{1 - kr^2}} \quad (1.14)$$

The metric then becomes :

$$ds^2 = dt^2 - a^2(t) [d\chi^2 + S_k^2(\chi)(d\theta^2 + \sin^2(\theta)d\phi^2)] \quad (1.15)$$

1.4 Friedmann equations

Since the Universe is assumed to be homogenous and isotropic, we can express the energy-momentum tensor $T^{\mu\nu}$ only as a time-dependent object describing the density ρ and pressure P of every component present in the Universe. We assume that these can be described as a perfect fluid with no viscosity. For each fluid present indexed by i , we can then write down the pressure term as $P_i = w\rho_i c^2$ where w is the equation of state parameter. This gives us the following form for $T^{\mu\nu}$:

$$T_i^{\mu\nu} = \rho_i [(w_i + 1)u^\mu u^\nu - w_i g^{\mu\nu}] \quad (1.16)$$

with u^μ being the velocity vector of the fluid.

Using this, the Einstein equations and the FLRW metric, Friedmann derived the following equations known as the Friedmann equations in 1924 (REF):

$$H^2 = \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho_i - \frac{kc^2}{a^2} \quad (1.17)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho_i + 3\frac{p_i}{c^2}\right) \quad (1.18)$$

The first equation describes how fast the Universe is expanding knowing the Universe's composition ρ_i and curvature k , while the second describes its acceleration. Combining equations 1.17 and 1.18 we get:

$$\frac{\dot{\rho}_i}{\rho_i} = -3(w_i + 1)\frac{\dot{a}}{a} \quad (1.19)$$

And the solution to this differential equation is:

$$\boxed{\rho_i(t) = \rho_i(t_0)a(t)^{-3(w_i+1)}} \quad (1.20)$$

This entails that the evolution of the scale factor is entirely determined by the nature of the fluids present at a time t . Knowing w of each fluid informs us on the behaviour of $a(t)$. If we consider one fluid that dominates over, the scale factor can be written as:

$$a(t) \sim t^{\frac{2}{3(1+w)}} \quad (1.21)$$

Consequently, the value of w of the dominant fluid determines the behaviour of the expansion:

$$\begin{array}{lll} \ddot{a} > 0 & w < -\frac{1}{3} & \text{accelerated expansion} \\ \ddot{a} = 0 & w = -\frac{1}{3} & \text{constant expansion} \\ \ddot{a} < 0 & w > -\frac{1}{3} & \text{decelerated expansion} \end{array} \quad (1.22)$$

1.5 Energy contents of the Universe

Within the current cosmological framework, the total energy density of the Universe is commonly divided into the following main components:

- **Radiation** ($w_r = \frac{1}{3}$): a relativistic component composed primarily of photons and neutrinos. Its contribution is negligible in the present-day Universe, but it dominated the energy budget during the earliest cosmological epochs.

- **Matter** ($w_m = 0$): a non-relativistic component including both baryonic matter and dark matter. While baryonic matter constitutes the visible content of galaxies, the dominant fraction is dark matter, whose existence was first inferred from galaxy cluster dynamics by Zwicky ([Zwicky, 1933], [Andernach and Zwicky, 2017]). In the standard cosmological model, dark matter is assumed to be cold, meaning that its particles were non-relativistic during the epoch of structure formation.
- **Cosmological constant / Dark energy** ($w_\Lambda = -1$): in the standard Λ CDM model, cosmic acceleration is described by the cosmological constant Λ , interpreted as an intrinsic property of spacetime with equation of state $w = -1$. More generally, dark energy may be viewed as an additional fluid component driving accelerated expansion. Possible deviations from Λ CDM include models with a constant but free equation of state (w CDM), or a time-evolving equation of state such as the CPL parametrization ([Chevallier and Polarski, 2001]).

COMMENT : might be worth mentioning the quantum vacuum fluctuations and the order of magnitude issue

The equations of state for matter and radiation can be determined using a statistical physics framework. For simplicity, we will not be going into the details of the derivations for each component but simply give the end results.

For non relativistic species, such as matter, we obtain :

$$w_m = 0 \quad (1.23)$$

For relativistic species, such as radiation, we obtain :

$$w_r = \frac{1}{3} \quad (1.24)$$

For the cosmological constant this can be determined without going through the statistical physics approach where by simply applying the Friedmann equations on a constant naturally leads to ([**Review this part**]):

$$w_\Lambda = -1 \quad (1.25)$$

We introduce the critical density ρ_c defined as the energy density of the Universe for $k = 0$ and at $t = 0$:

$$\rho_c = \frac{3H_0^2}{8\pi G} \quad (1.26)$$

This critical density is used to normalize the densities of the other fluids such that the normalized density is written as:

$$\Omega_i = \frac{\rho_i}{\rho_c} \quad (1.27)$$

At $t = 0$, the Friedmann equation yields $\Omega_m + \Omega_r + \Omega_\Lambda + \Omega_k = 1$ with Ω_k being the contribution from the curvature by construction:

$$\Omega_k = \frac{-kc^2}{a(t_0)^2 H_0^2} \quad (1.28)$$

Equation 1.17 can be then rewritten as :

$$H(z) = H_0 \sqrt{\Omega_{m,0} a(t)^{-3} + \Omega_{r,0} a(t)^{-4} + \Omega_{\Lambda,0} + \Omega_{k,0} a(t)^{-2}} \quad (1.29)$$

Current cosmological observations indicate that the present-day Universe is remarkably close to spatial flatness, corresponding to $\Omega_{k,0} \simeq 0$ [Aghanim et al., 2020]. The dominant contribution to the total energy budget is consistent with a cosmological constant (or dark energy), representing approximately 70% of the total density. Matter contributes roughly 30%, of which about 25% corresponds to dark

matter and only $\sim 5\%$ to baryonic matter. The radiation component today is subdominant, contributing less than 1% of the total energy density.

The relative importance of these components evolves with cosmic expansion according to their different scalings with the scale factor. The evolution of each matter component is represented in Figure 1.4. Since radiation scales as a^{-4} , matter as a^{-3} , and the cosmological constant remains constant, the early Universe was initially radiation dominated. As the Universe expanded and cooled, a transition to matter domination occurred, enabling the growth of gravitational instabilities and the formation of large-scale structure. At late times, the cosmological constant became dominant, driving the current phase of accelerated expansion.




Fig. 1.4 Evolution of the main components of the Λ CDM model with respect to the scale factor. This highlights where each component dominates as well as when they become equal.

The acceleration of the Universe was confirmed by [Riess et al., 1998] and [Perlmutter et al., 1999] when they constructed a Hubble diagram with distant Type Ia Supernovae. Each paper has shown that the supernova Hubble diagram cannot be solely explained with a Universe containing only matter and that an additional energetic component must be accounted for. This is seen in Figure 1.6.

1.6 The Λ CDM Model

The Λ CDM model is the standard model of cosmology. It is built upon the framework of GR and assumes that the Universe began in an extremely hot and dense state, known as the Hot Big Bang, and has




Fig. 1.5 Hubble diagram constructed using 50 type Ia supernovae from [Riess et al., 1998] which confirmed the accelerated expansion. The dotted line represents the prediction for a Universe containing only matter that accounts for 20% of the energy content. The dashed line is for a Universe containing matter that accounts for 100% of the energy content of the Universe. The solid line is for a Universe containing matter energy content of 24% and Λ accounting for 76% of the energy content. The data fits the the solid line best.

been expanding ever since. As the Universe expanded and cooled, it underwent three successive epochs of domination by different energy components, each governing the dynamics of expansion through the Friedmann equations: a radiation dominated era at early times, followed by a matter dominated era, and finally the current dark energy dominated era driven by the cosmological constant Λ .

The energy content of the Universe in Λ CDM consists of the components introduced in the previous section: radiation, baryonic matter, cold dark matter and a cosmological constant Λ , with their present-

day fractions measured to be approximately $\Omega_r \approx 9 \times 10^{-5}$, $\Omega_b \approx 0.05$, $\Omega_{\text{CDM}} \approx 0.26$ and $\Omega_\Lambda \approx 0.69$ respectively ([Aghanim et al., 2020]), in a spatially flat Universe where $\sum_i \Omega_i = 1$. The model makes the additional assumptions that the Universe is homogeneous and isotropic on large scales (the cosmological principle), and that the primordial density perturbations seeding all structures are adiabatic and nearly Gaussian ([Collaboration et al., 2019]).

The Λ CDM model is fully characterised by six independent cosmological parameters:

- A_s — the amplitude of primordial scalar perturbations
- n_s — the spectral index of the primordial power spectrum
- $\Omega_b h^2$ — the physical baryon density
- $\Omega_{\text{CDM}} h^2$ — the physical cold dark matter density
- H_0 — the Hubble constant, parametrising the present-day expansion rate
- τ — the optical depth to reionisation

All other cosmological quantities, such as Ω_Λ , σ_8 and S_8 , are derived from these six parameters within the model. Despite its simplicity, Λ CDM has proven remarkably successful in describing a wide range of observations, from the CMB anisotropies ([Aghanim et al., 2020]) to the large-scale distribution of galaxies ([DESI Collaboration et al., 2025]), the BAO standard ruler ([Eisenstein et al., 2005]), and the accelerated expansion of the Universe ([Riess et al., 1998, Perlmutter et al., 1999]). Nevertheless, as discussed in Section 5, emerging tensions between different observational probes suggest that Λ CDM may require extensions or modifications.

[Transition to next section: structure formation theory. In Pauline’s notes three pillars of HBB model]

2 Large Scale Structure Formation

Now that we have understood how modern cosmology came to be, we can move on to understanding the formation of structures. Observations from galaxy surveys such as DESI show that although on large scales the Universe looks homogenous, the smaller scales indicate the existence of an underlying non-homogenous structure. This can be seen through the formation of galaxy clusters and the cosmic web. Understanding how these structures are formed gives us a lot of information regarding the strength of gravity at large scales as well as the amount of matter in the Universe. Since the full derivation of each equation in structure formation is too long to write for this thesis, we recommend reading [Knobel, 2013] and [Dodelson, 2003] for a more detailed derivation

2.1 Newtonian approach to structure formation

In this introductory section, we will be starting off by describing structure formation by going only to linear scales. This will be the model on which we build up more complex structure formation model.

Theory of structure formation

We start by assuming a Universe is filled with an inhomogeneous, dissipationless, ideal fluid with matter density $\rho(\mathbf{r}, t)$, velocity field $\vec{v}(\mathbf{r}, t)$, pressure $P(\mathbf{r}, t)$ and entropy per unit mass $S(\mathbf{r}, t)$ with $\mathbf{r}$ being the fluid's coordinates. We obtain the following hydrodynamical equations:

$$\begin{cases} \frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot (\rho \vec{v}) = 0 & \text{Continuity equation} \\ \frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} + \frac{\vec{\nabla} P}{\rho} + \vec{\nabla} \Phi = 0 & \text{Euler equation} \\ \nabla^2 \Phi = 4\pi G \rho & \text{Poisson equation} \\ \frac{\partial S}{\partial t} + (\vec{v} \cdot \vec{\nabla}) S = 0 & \text{Entropy conservation} \end{cases} \quad (1.30)$$

with Φ the gravitational potential. Since matter is approximated as a single-stream fluid, neighboring particles follow nearly parallel trajectories without "shell crossing" (collision of streams). This description breaks down once nonlinear collapse occurs, when multiple streams overlap and complex dynamics emerge. A fully accurate treatment of structure formation would instead require solving the general relativistic Boltzmann equation including all cosmic components and their interactions. This is in itself a difficult problem due to the nonlinear coupling between the fluid equations. To overcome that issue, we assume a FLRW Universe containing cold dark matter (treated as a fluid) and perturb the fluid around the Hubble flow, meaning we separate each variable as a homogenous part and an inhomogeneous perturbation. The cold dark matter assumption justifies the use of Newtonian mechanics since we are dealing with non-relativistic components. Moreover we assume that the perturbations that arise are very small. We can then write down the expansion of the quantities used in the hydrodynamical equation at first order as:

$$\begin{aligned} \rho(\mathbf{r}, t) &= \bar{\rho}(t) + \delta\rho(\mathbf{r}, t) \\ \mathbf{v}(\mathbf{r}, t) &= \bar{\mathbf{v}}(\mathbf{r}, t) + \delta\mathbf{v}(\mathbf{r}, t) \\ P(\mathbf{r}, t) &= \bar{P}(t) + \delta P(\mathbf{r}, t) \\ \Phi(\mathbf{r}, t) &= \bar{\Phi}(\mathbf{r}, t) + \delta\Phi(\mathbf{r}, t) \\ S(\mathbf{r}, t) &= \bar{S}(t) + \delta S(\mathbf{r}, t) \end{aligned}$$

We can then re-introduce our new variables in Equations 1.30 although to slightly simplify the equations we will be working in comoving coordinates $\mathbf{x}$ such that

$$\begin{aligned}\mathbf{r} &= a\mathbf{x} \\ \nabla_{\mathbf{r}} &= \frac{1}{a}\nabla_{\mathbf{x}} \\ \left.\frac{\partial}{\partial t}\right|_{\mathbf{r}} &= \left.\frac{\partial}{\partial t}\right|_{\mathbf{x}} - \frac{1}{a}\mathbf{v} \cdot \nabla_{\mathbf{x}}\end{aligned}$$

We will also be working in Fourier space where we get for any perturbation $\delta q(\mathbf{x}, t)$:

$$\delta q(t, \mathbf{k}) = \int \delta q(t, \mathbf{x}) e^{-i\mathbf{k}\mathbf{x}} d^3x \quad (1.31)$$

$$\delta q(t, \mathbf{x}) = \frac{1}{(2\pi)^3} \int \delta q(t, \mathbf{k}) e^{-i\mathbf{k}\mathbf{x}} d^3k \quad (1.32)$$

With $\mathbf{k}$ are the comoving Fourier modes. We get the following hydrodynamical equations at first order:

$\begin{cases} \delta\dot{\rho} + 3H\delta\rho + \frac{i\bar{\rho}\mathbf{k}}{a} \cdot \delta\mathbf{v} = 0 \\ \delta\dot{\mathbf{v}} + H\delta\mathbf{v} + \frac{i\mathbf{k}}{a\bar{\rho}} (c_s^2\delta\rho + \sigma\delta S) + \frac{i\mathbf{k}}{a}\delta\Phi = 0 \\ k^2\delta\Phi + 4\pi G a^2\delta\rho = 0 \\ \delta\dot{S} = 0 \end{cases}$	Continuity equation Euler equation Poisson equation Entropy conservation	(1.33)
--	--	--------

with $c_s^2 = (\partial\bar{P}/\partial\bar{\rho})_{\bar{S}}$ the speed of sound in the fluid, $\sigma = (\partial\bar{P}/\partial\bar{S})_{\bar{\rho}}$ and $k = |\mathbf{k}|$.

COMMENT : how do we get the 3h delta rho?

What we are interested in is essentially how do matter perturbations grow as they will give birth to structure by gravitational attraction. The first thing to note is that the entropy fluctuations must exist at early times in order for them to keep existing otherwise they would simply dissipate. Moreover, results from the simplest models of inflation do not support the fluctuations of entropy. So from now on we will set $\delta S = 0$. We can also assume that the velocity perturbations would follow the overdense regions without experiencing any vorticities giving us modes that are parallel to the velocity $\mathbf{k}||\delta\mathbf{v}$. We call the modes that result from these conditions the adiabatic modes.

COMMENT : say no vorticity to justify parallel and separate from adiabatic modes.

We will be denoting the overdensity $\delta(\mathbf{k}, t)$ as the matter density contrast which is written as:

$$\delta(\mathbf{k}, t) = \frac{\rho(\mathbf{k}, t)}{\bar{\rho}(t)} - 1 = \frac{\delta\rho(\mathbf{k}, t)}{\bar{\rho}(t)} \quad (1.34)$$

In the adiabatic modes, the continuity equation becomes:

$$\dot{\delta} + \frac{ik}{a}\delta v = 0 \quad (1.35)$$

If we then differentiate this equation and combining its results with Euler and Poisson equations we obtain:

$$\ddot{\delta} + 2H\dot{\delta} + \left(\frac{c_s^2 k^2}{a^2} - 4\pi G\bar{\rho}\right)\delta = 0 \quad (1.36)$$

Here we obtain a linear second order differential equation with two solutions to it. We denote $\lambda = \frac{2\pi a}{k}$ the physical wavelength of the mode, $\lambda_J = c_s \sqrt{\frac{\pi}{G\bar{\rho}}}$ the Jeans length and $M_J = \frac{\pi}{6}\lambda_J^3\bar{\rho}$ the Jeans mass.

Equation 1.36 tells us that if $\lambda < \lambda_J$, the solution to the equation is an oscillating wave leaving no room for structures to grow. So a condition we must have is $\lambda > \lambda_J$. We place ourselves in the condition where the wavelength is much larger than the Jeans length $\lambda \gg \lambda_J$, Equation 1.36 is now reduced to:

$$\ddot{\delta} + 2H\dot{\delta} - 4\pi G\bar{\rho}\delta = 0 \quad (1.37)$$

COMMENT : important to specify this reduction for collisionless matter so DM but not baryonic

which has the following general solution:

$$\delta(\mathbf{k}, t) = \delta_+(\mathbf{k})D_+(t) + \delta_-(\mathbf{k})D_-(t) \quad (1.38)$$

with the + and - denoting growing and decaying modes. We call the $D_+(t)$ term the **linear growth function**. In the radiation dominated era, the Mészáros effect suppresses the growth of sub-Hubble matter perturbations, which grow only logarithmically rather than as a power law. In the matter dominated era the expansion rate grows as $H(t) = \frac{2}{3t}$ which gives us:

$$D_+ \propto t^{2/3} \propto a(t) \quad (1.39)$$

$$D_- \propto t^{-1} \quad (1.40)$$

This means that we can get grown modes for matter perturbations in the matter dominated era with D_+ while D_- goes to 0. A general formula for D_+ can be obtained ([Carroll et al., 1992]) and it gives us:

$$D_+(z) = \frac{1}{1+z} \frac{g(z)}{g(0)} \quad (1.41)$$

$$g(z) = \frac{\Omega_m(z)}{\Omega_m^{4/7}(z) - \Omega_\Lambda(z) + (1 + \Omega_m(z)/2)(1 + \Omega_\Lambda(z)/70)} \quad (1.42)$$

The growth of perturbations described by D_+ can be characterised by the *growth rate* f , defined as the logarithmic derivative of the growth factor with respect to the scale factor:

$$f = \frac{d \ln D_+}{d \ln a} \quad (1.43)$$

In the matter dominated era, since $D_+ \propto a$, we get $f = 1$. A useful approximation valid across different epochs and cosmologies is given by ([Peebles, 1980]):

$$f \approx \Omega_m(z)^\gamma \quad (1.44)$$

where γ is obtained from the underlying gravitational theory. For a flat Λ CDM cosmology and GR it is $\gamma \simeq 0.55$. Deviations from this value would indicate modifications to GR on cosmological scales. In practice, f is not directly observable, and is instead measured in combination with σ_8 , the root mean square of matter fluctuations on scales of $8h^{-1}\text{Mpc}$. The combination $f\sigma_8$ is therefore the key observable quantity that encodes the rate of growth of large-scale structure and provides a direct test of gravity on cosmological scales.

COMMENT : not just Peebles more ref and there is an expression for gamma wrt w
COMMENT : can relate here or later As connecton

Statistical analysis of overdensities

Practically, we perform a statistical analysis when it comes to the study of the structures of the Universe. δ is looked at as a realization of a stochastic process and by analyzing that we can compare our observations to theories.

COMMENT : primordial fluctuations expected from inflation described in later section implies stochastic nature etc... can relate to CMB observations today (just to make the link)

We introduce $\langle \dots \rangle$ as the ensemble average of the stochastic process underlying a random field. The large-scale structure we observe is only one realization of an underlying random cosmic process, so estimating its true statistical properties requires additional assumptions. A common assumption is ergodicity, meaning that averages measured over a sufficiently large volume of the Universe approximate the averages over many hypothetical realizations. In practice, this is related to the fair sample hypothesis, which assumes that widely separated regions of the Universe can be treated as nearly independent samples of the same process. If the survey volume is large enough and correlations become weak on large scales, these approximations are reasonable. However, real galaxy surveys cover finite volumes and therefore their measured averages are affected by statistical fluctuations known as **sample variance**, or **cosmic variance** when limited by the size of the observable Universe.

We denote the 2-point correlation function as:

$$\xi(\mathbf{x}, \mathbf{x}') = \langle \delta(\mathbf{x})\delta(\mathbf{x}') \rangle \quad (1.45)$$

Our field is homogenous and isotropic therefore the only quantity that affects the correlation function is $r = |\mathbf{x} - \mathbf{x}'|$, $\xi(\mathbf{x}, \mathbf{x}') = \xi(r)$. If at $r = +\infty$, the correlations go to 0, then we can decompose it in Fourier modes: $\delta(\mathbf{k})$:

$$\langle \delta(\mathbf{k})\delta^*(\mathbf{k}') \rangle = (2\pi)^3 \delta_D(\mathbf{k} - \mathbf{k}') \mathcal{P}(\mathbf{k}) \quad (1.46)$$

$$\mathcal{P}(\mathbf{k}) = \int \xi(\mathbf{x}) e^{-i\mathbf{k}\mathbf{x}} d^3x \quad (1.47)$$

where the $\mathcal{P}(k)$ is the **power spectrum**, which is a real function, and δ_D is the Dirac delta function. The works from [Harrison, 1970], [Zel'dovich, 1970b] and [Peebles and Yu, 1970] show that there is a preferable form for the initial primordial power spectrum that takes the form:

$$\mathcal{P}(k) \propto k^{n_s} \quad (1.48)$$

with n_s the **spectral index**. If $n_s = 1$ the power spectrum becomes scale invariant. It is possible to predict the shape of the power spectrum of the initial conditions by studying an inflationary model. Most common models and results from [Collaboration et al., 2019] indicate that the initial perturbations are Gaussian with $n_s \approx 1$.

The power spectrum for the linear overdensity field is:

$$\mathcal{P}_{\text{lin}}(t, k) = A_s k^{n_s} T^2(k) D_+^2(t) \quad (1.49)$$

with A_s the normalization parameter and $T(k)$ is the transfer function which characterizes how each Fourier mode of the density field evolved from the initial conditions to a later time.

COMMENT : can link sigma8 and A_s here

2.2 Relativistic structure formation

Similarly to the previous section, we will be introducing structure formation from a general relativistic framework with a focus on linear perturbation. We will not be deriving every single quantity, however explanation of how we get there and how it relates to the Newtonian approximation will be present.

General description of perturbations

Firstly, we will be working in the FLRW metric for $k = 0$. We will also work in conformal time τ which simplifies this problem. It is defined as $ad\tau = dt$. We will also work in natural units setting $c = \hbar = 1$. We will denote the FLRW metric as $\bar{g}_{\mu\nu}$ so that for a flat spacetime we have:

$$ds^2 = \bar{g}_{\mu\nu} dx^\mu dx^\nu = a^2(\tau) (-d\tau^2 + \bar{\gamma}_{ij} dx^i dx^j) \quad (1.50)$$

where $\bar{\gamma}_{ij}$ is the metric for the spatial components and (i, j) can take the values 1, 2, 3 (in general we will put a bar on top of the parameter related to a flat FLRW metric).

In a relativistic framework, the perturbations are done on the metric in itself such that when we choose a metric $g_{\mu\nu}$ the deviations from the homogenous isotropic background are close enough to $\bar{g}_{\mu\nu}$ ². We then obtain:

$$\delta g_{\mu\nu} = g_{\mu\nu}(x) - \bar{g}_{\mu\nu}(x) \quad (1.51)$$

Physically, we are splitting $g_{\mu\nu}$ into a background $\bar{g}_{\mu\nu}$ and perturbation $\delta g_{\mu\nu}$. However this perturbation is not a 4-tensor which will require us to set a gauge transformation. Similarly, the perturbation for a general scalar-like object and a vector-like objects are defined as:

$$\delta q(x) = q(x) - \bar{q}(x) \quad (1.52)$$

$$\delta q^\mu = q^\mu - \bar{q}^\mu = \delta q_\nu \bar{g}^{\mu\nu} + \bar{q}_\nu \delta g^{\mu\nu} \quad (1.53)$$

We can also perform the same for the energy-momentum tensor.

$$\delta T_{\mu\nu} = T_{\mu\nu} - \bar{T}_{\mu\nu} \quad (1.54)$$

with

$$T_{\mu\nu} = (\rho + P)u_\mu u_\nu + P g_{\mu\nu} + \Pi_{\mu\nu} \quad (1.55)$$

and $\Pi_{\mu\nu}$ is the anisotropic stress which accounts for deviations from an ideal fluid.

The general form for the metric perturbation is written as:

$$\delta g_{\mu\nu} dx^\mu dx^\nu = a^2(\tau) [-2A d\tau^2 + B_i d\tau dx^i + 2(H_L \delta_{ij} + H_{ij}) dx^i dx^j] \quad (1.56)$$

where $A(\tau, \mathbf{x})$, $B_i(\tau, \mathbf{x})$, $H_L(\tau, \mathbf{x})$ and $H_{ij}(\tau, \mathbf{x})$ are perturbation quantities and δ_{ij} a delta function.

Newtonian gauge

These perturbation quantities intervene will then intervene in the field equations. Determining them necessitates a gauge transformation. We will be showing the results of the Newtonian gauge as it is useful

²Mathematically, this change in metric means we also the manifolds being worked on so simple mathematical operations no longer apply. One way to overcome that issue is to define $\bar{g}_{\mu\nu}$'s functional form as invariant under coordinate transformations.

for analyzing the density fluctuations inside of the Hubble horizon ($D_H = 1/H$) as well as it is simple to interpret its results in terms of Newtonian physics. The gauge is defined by:

$$B = H = 0 \quad (1.57)$$

and we will be renaming the other perturbations as:

$$\Psi = A \quad (1.58)$$

$$\Phi = -H_L \quad (1.59)$$

The field equations for the Newtonian gauge are:

$$\begin{cases} -k^2\Phi - 3H(\dot{\Phi} + H\Psi) = 4\pi G a^2 \delta\rho \\ \dot{\Phi} + H\Psi = 4\pi G a^2 \frac{v}{k}(\bar{\rho} + \bar{P}) \\ \ddot{\Phi} + H(\dot{\Psi} + 2\dot{\Phi}) + (2\dot{H} + H^2)\Psi + \frac{k^2}{3}(\Phi - \Psi) = 4\pi G a^2 \delta P \\ k^2(\Phi - \Psi) = 8\pi G a^2 \Pi \end{cases} \quad (1.60)$$

where the dots are derivatives with respect to the conformal time. If we consider scale much smaller than the Hubble horizon $H/k \ll 1$, the first field equation then becomes:

$$-k^2\Phi = 4\pi G a^2 \delta\rho \quad (1.61)$$

In this limit we recover the Poisson equation in the Newtonian formalism. Therefore, we can say that well within the horizon the Newtonian gauge reduces to Newtonian mechanics. This also means we can have a physical interpretation for Φ it acting as the Newtonian gravitational potential (called generalized Newtonian potential). The last equation also informs us that in the matter dominated epoch we have $\Phi \simeq \Psi$ reducing the metric perturbations to only one parameter.

One can only also derive the equations of motion in the Newtonian gauge, however for this thesis we will only be showing the final equations of motions obtained for matter fluctuations in the matter dominated era. We note the matter overdensity $\delta_m = \frac{\delta\rho_m}{\bar{\rho}}$, we obtain:

$$\ddot{\delta}_m + H\dot{\delta}_m - 4\pi G a^2 \bar{\rho}_m \delta_m = 0 \quad (1.62)$$

2.3 Inflation

Inflation is a period of time at the very early stages of the Universe where an accelerated expansion of the Universe occurred ([Guth, 1981]). It was theorized to tackle known issues regarding the fine tuning of the flatness of the Universe and the horizon problem. What we see as well is that through inflation we can have a predict for the initial density fluctuations. This following condition is imposed on inflation to obtain an accelerated early Universe:

$$\ddot{a} > 0 \quad (1.63)$$

During inflation, we assume that the Universe is dominated by a quantum scalar field ϕ called the inflaton, experiencing a "slow roll" down its potential $V(\phi)$ ([Mukhanov et al., 1992], [Albrecht and Steinhardt, 1982]). The action for a scalar field minimally coupled to gravity is given by:

$$S = \int d^4x \sqrt{-g} \left[\frac{M_{\text{Pl}}^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right] \quad (1.64)$$

where M_{pl} is the Planck mass defined as $M_{\text{pl}}^2 = (\frac{1}{8\pi G})^2$. Varying the action with the respect to the metric allows to obtain the energy momentum tensor $T_{\mu\nu}$ and its components P_ϕ and ρ_ϕ :

$$P_\phi = \frac{1}{2}\dot{\phi}^2 - V(\phi) \quad (1.65)$$

$$\rho_\phi = \frac{1}{2}\dot{\phi}^2 + V(\phi) \quad (1.66)$$

where the kinetic energy of the field is given by $\frac{1}{2}\dot{\phi}^2$.

The dynamics can be recovered with the Klein Gordon equation giving us:

$$\ddot{\phi} + 3H\dot{\phi} + \frac{dV(\phi)}{d\phi} = 0 \quad (1.67)$$

We can also apply the Friedmann equations to the inflaton field where we get:

$$H^2 = \frac{1}{M_{\text{pl}}^2} \left(\frac{1}{2}\dot{\phi}^2 + V(\phi) \right) \quad (1.68)$$

$$\dot{H} = -\frac{1}{2} \frac{\dot{\phi}^2}{M_{\text{pl}}^2} \quad (1.69)$$

Assuming a slow rolling field entails that kinetic term is small compared to the potential term V . We can show that given this assumption $a \sim e^{Ht}$ which is a necessary condition to inflation. Additionally we assume coherent oscillations around a minimum that are damped by a $H\dot{\phi}$ as seen in the Klein-Gordon equation. The field is then expected to decay into matter and radiation, although the exact physical process behind that is unknown. The inflaton field is then treated as a perturbation around a mean field, that these perturbation give rise to perturbation amplitudes that we can identify in the observations we make today. Some useful parameters to define for a slow-roll inflation are :

$$\epsilon = -\frac{\dot{H}}{H^2} = \frac{3}{2} \frac{\dot{\phi}^2}{V} \quad (1.70)$$

$$\eta = \epsilon - \frac{\ddot{\phi}}{H\dot{\phi}} \quad (1.71)$$

where the slow rolling conditions are $\epsilon \ll 1$ and $\eta \ll 1$.

We define q as:

$$q(x) = \delta\phi + \frac{\dot{\phi}}{H}\Phi \quad (1.72)$$

This quantity q is known as the Mukhanov-Sasaki variable ([Mukhanov, 1988], [Sasaki, 1986]) and is the canonical variable for quantizing inflaton perturbations.

In order to fully treat the inflaton field we have to apply the canonical quantization of the field (quantum field theory) by elevating q to an operator $\hat{q}$ as well as introducing its canonical conjugate momentum $\hat{\pi}$. In the Fourier space, they are expressed as function of the quantum harmonic operator $a_{\mathbf{k}}$ and $a_{\mathbf{k}}^\dagger$. With these operators we can calculate the expected value of the field as well as power spectrum as a function of k . Once the power spectrum during inflation is determined it is possible to extend this to the primordial power spectrum of matter during the matter dominated era. Outside the Hubble horizon, each mode is frozen with a nearly scale-invariant amplitude. Before writing the primordial power spectrum, we introduce the comoving curvature perturbation $\mathcal{R}$, a gauge-invariant quantity that

measures the spatial curvature of constant-density hypersurfaces. In the context of inflation it is related to the inflaton perturbation $\delta\phi$ by:

$$\mathcal{R} = -\frac{H}{\dot{\phi}}\delta\phi \quad (1.73)$$

The key property of $\mathcal{R}$ is that it is conserved on super-Hubble scales for adiabatic perturbations ([Wands et al., 2000]). This means its amplitude is frozen once a mode exits the Hubble radius during inflation and remains unchanged until it re-enters the horizon during the hot Big Bang, directly connecting inflationary predictions to late-time observables. The dimensionless power spectrum of the comoving curvature perturbation $\mathcal{R}$ evaluated at Hubble crossing $k = aH$ is:

$$\mathcal{P}_{\mathcal{R}}(k) = \frac{H^4}{4\pi^2\dot{\phi}^2} \Big|_{k=aH} \approx \frac{H^2}{8\pi^2 M_{\text{pl}}^2 \epsilon} \Big|_{k=aH} \quad (1.74)$$

which is approximately scale-invariant for slow-roll inflation where H and ϵ vary slowly. The spectral index for a slow-roll inflation is given by:

$$n_s = 1 - 6\epsilon + 2\eta \quad (1.75)$$

Once modes re-enter the Hubble horizon during the matter and radiation dominated eras, they begin to evolve dynamically. Their evolution is captured by the transfer function $T(k)$ and the linear growth factor D_+ . The linear matter power spectrum at a time τ is therefore:

$$\mathcal{P}_{\text{lin}}(\tau, k) = A_s \left(\frac{k}{k_*} \right)^{n_s} T^2(k) D_+^2(\tau) \quad (1.76)$$

where A_s is the primordial amplitude defined at the pivot scale $k_* = 0.05 \text{ Mpc}^{-1}$ ([Planck Collaboration et al., 2020]), $T(k)$ encodes the suppression of small-scale modes that entered the horizon during radiation domination ([Eisenstein and Hu, 1998]), and $D_+(\tau)$ describes the growth of perturbations during matter domination ([Carroll et al., 1992]). The k^{n_s} factor reflects the nearly scale-invariant primordial spectrum, with $n_s \approx 0.965$ ([Planck Collaboration et al., 2020]) consistent with slow-roll inflation.

3 Cosmological distances

Observables such as the redshift or distances are used in order to constrain the matter contents of the Universe as well as its expansion history and cosmological parameters. There are multiple ways one can define a distance in cosmology each relying on a different set of parameters.

Physical distances

Under a FLRW metric, we can define the physical distance as the distance separating the observer and an event happening at χ :

$$d_p(\chi) = a_0 S_k(\chi) \quad (1.77)$$

with a_0 being the scale factor at t_0 . It is more practical to express χ as a function of the redshift since it is not a direct observable. If we consider a photon emitted at a time t_e from the source object present at χ and recieved by the observer at t_0 then we get:

$$\chi = c \int_{t_e}^{t_0} \frac{dt}{a(t)} \quad (1.78)$$

By performing the following two variable changes:

$$H = \frac{\dot{a}}{a} = \frac{1}{a} \frac{da}{dt} \iff dt = \frac{da}{aH} \quad (1.79)$$

$$1 + z = \frac{a_0}{a} \iff dz = -\frac{a_0}{a^2} da \quad (1.80)$$

We get:

$$\chi = \frac{c}{a_0} \int_0^z \frac{dz'}{H(z')} \quad (1.81)$$

Which gives us the following equation for the physical distance:

$$d_p(z) = a_0 S_k \left(\frac{c}{a_0} \int_0^z \frac{dz'}{H(z')} \right) \quad (1.82)$$

In a flat Universe, this expression is reduced to:

$$d_p(z) = c \int_0^z \frac{dz'}{H(z')} \quad (1.83)$$

In practice, this distance isn't directly measurable as it requires to measure the values H at every redshift along the line of sight. However, other observable distances exist to which we can relate to the physical distance.

Angular distances

We can firstly define the distance of an object based on its apparent size D and angle $\Delta\theta$. We call this the angular distance and is defined as:

$$d_A = \frac{D}{\delta\theta} = a(t) S_k(\chi) = \frac{1}{1+z} a_0 S_k \left(\frac{c}{a_0} \int_0^z \frac{dz'}{H(z')} \right) \quad (1.84)$$

Therefore knowing the redshift, its apparent angle and size of an object gives us access to its distance.

Luminosity distances

We can also determine the distance of an object by linking its intrinsic luminosity to its emitted flux at the surface. This defined as the luminosity distance d_L and is given by:

$$f = \frac{L}{4\pi d_L^2} \quad (1.85)$$

where f is the surface flux and L is the object's luminosity.

We need to account for the fact that the photons emitted are scattered on the surface S of the object $S = 4\pi a_0^2 S_k^2(\chi)$ as well as the fact that its energy ($\frac{hc}{\lambda}$) is reduced by $1+z$ due to the cosmic expansion. The photons emitted during a duration of Δt will be received during a diluted duration $\Delta t(1+z)$. This means that the observed luminosity will differ from the emitted luminosity giving us:

$$L' = \frac{L}{(1+z)^2} \quad (1.86)$$

where L' is the luminosity seen by the observer. By definition, the surface flux is seen at an observer frame meaning:

$$f = \frac{L'}{S} = \frac{L}{(1+z)^2 4\pi a_0^2 S_k^2(\chi)} \quad (1.87)$$

We can then identify a formula for the luminosity distance that is redshift dependent:

$$d_L(z) = (1+z)a_0 S_k \left(\frac{c}{a_0} \int_0^z \frac{dz'}{H(z')} \right) \quad (1.88)$$

We can see from the previous equations that:

$$d_L(z) = d_A(z)(1+z)^2 = d_p(z)(1+z) \quad (1.89)$$

At low redshift ($z \lesssim 0.15$) all three distance expression become equivalent and we can approximate it to:

$$d_L(z) \approx \frac{cz}{H_0} \Leftrightarrow cz = v = H_0 d_L \quad (1.90)$$

where we can see that we recover the Hubble law.

4 Cosmological Observables

In this section we will be briefly presenting some of the main observables that cosmologists are concerned with.

4.1 The Cosmic Microwave Background

During the radiation dominated era, the Universe was hot and dense, filled with a plasma of baryons and free electrons. Atoms could not form at that time because photons were energetic enough to ionise any neutral hydrogen immediately upon formation. As a result, baryonic matter remained tightly coupled to photons through Thomson scattering off free electrons. As the Universe expanded and cooled, photons eventually became insufficiently energetic to ionise hydrogen, allowing neutral atoms to form




Fig. 1.6 Different distances in cosmology plotted as a function of redshift for a flat Λ CDM with $H_0 = 70 \text{ km.s}^{-1} \text{ Mpc}^{-1}$ and $\Omega_m = 0.3$.

for the first time. This epoch is referred to as **recombination** and is associated with the decoupling of baryons and photons. The mean free path of photons then exceeded the Hubble radius, allowing them to travel freely across space without scattering. Today, these photons have been redshifted drastically due to the expansion of the Universe and are observed as a background of microwave radiation. We refer to this relic radiation as the Cosmic Microwave Background (CMB), and it constitutes our primary observational window onto the primordial Universe. The CMB was first discovered by Arno Penzias and Robert Wilson accidentally while calibrating an antenna for telecommunications and astronomy ([Penzias and Wilson, 1965a, Penzias and Wilson, 1965b]).

The CMB has a nearly perfect blackbody spectrum with a mean temperature of $T_{\text{CMB}} = 2.7255 \pm 0.0006 \text{ K}$ ([Fixsen, 2009]) and temperature anisotropies at the level of $\Delta T/T \sim 10^{-5}$. The two-point correlation function of these anisotropies, equivalently described by the angular power spectrum C_ℓ , encodes the imprint of primordial density perturbations consistent with a Gaussian random field ([Collaboration et al., 2019]), which later evolved into the large-scale structures we observe today. Secondary anisotropies, arising from effects such as gravitational lensing by galaxy clusters and inverse Compton scattering of CMB photons off hot intracluster gas (the Sunyaev-Zel'dovich effect), provide additional information on the late-time Universe. The polarisation of the CMB, decomposed into curl-free E-modes and divergence-free B-modes, further constrains inflationary models and provides a potential window onto primordial gravitational

waves ([Kamionkowski et al., 1997]).

Through the detailed structure of the acoustic peaks in C_ℓ , CMB observations allow precise constraints on multiple cosmological parameters, including Ω_m , Ω_b , H_0 , n_s , and A_s ([Aghanim et al., 2020]).




Fig. 1.7 Compilation of CMB angular power spectrum measurements from [Collaboration et al., 2020]. The upper panel shows the power spectra of the temperature and E -mode and B -mode polarization signals, the next panel the cross-correlation spectrum between T and E , while the lower panel shows the lensing deflection power spectrum

4.2 Galaxy Surveys

Galaxy surveys map the three-dimensional distribution of galaxies across large volumes of the Universe. Galaxies are biased tracers of the underlying matter distribution, and their spatial statistics allow us to infer a vast amount of information about the structure, dynamics, and history of the Universe. Surveys such as SDSS ([York et al., 2000]) and DESI ([DESI Collaboration et al., 2016]) measure tens of millions of

galaxies across a wide redshift range, from which significant cosmological constraints have been extracted.

4.2.1 The Matter Power Spectrum

A key statistical tool for analysing galaxy surveys is the matter power spectrum $P(k)$, defined as the variance of density fluctuations as a function of scale k :

$$\langle \delta(\mathbf{k}) \delta^*(\mathbf{k}') \rangle = (2\pi)^3 \mathcal{P}(k) \delta_D^{(3)}(\mathbf{k} - \mathbf{k}') \quad (1.91)$$

The shape and amplitude of $\mathcal{P}(k)$ depend directly on cosmological parameters. The primordial power spectrum can be related to late-time observations through the transfer function ([Eisenstein and Hu, 1998]):

$$\mathcal{P}(k) = A_s \left(\frac{k}{k_*} \right)^{n_s-1} T^2(k) D_1^2(z) \quad (1.92)$$

where $k_* = 0.05 \text{ Mpc}^{-1}$ is the pivot scale at which the primordial amplitude A_s is defined ([Planck Collaboration et al., 2020]). $T(k)$ is the transfer function encoding the scale-dependent evolution of perturbations through radiation domination, and $D_1(z)$ is the linear growth factor describing the growth of structure during matter domination.




Fig. 1.8 Linear-theory matter power spectrum (at $z = 0$) inferred from different cosmological probes taken from [Collaboration et al., 2020].




Fig. 1.9 BAO signals in the measured correlation functions from SDD ([Alam et al., 2017]). On the left is the smoothed prediction of the best-fitted model. On the right the redshift bins are decomposed into the component of the separations transverse to and along the line of sight at $0.4 < z < 0.6$.

4.2.2 Baryon Acoustic Oscillations

Baryon Acoustic Oscillations (BAO) are a feature in the matter clustering that corresponds to the imprint of sound waves propagating through the baryon-photon plasma in the early Universe. The radiation pressure of the hot plasma drove acoustic oscillations at a sound speed of $c_s \simeq c/\sqrt{3}$. These oscillations were frozen in at the **drag epoch** $z_d \approx 1060$, shortly after photon decoupling at $z_* \approx 1100$, when baryons decoupled from the photon fluid and the sound waves could no longer propagate. The comoving distance these sound waves had travelled by that time defines the **sound horizon**:

$$r_s(z_d) = \int_{z_d}^{\infty} \frac{c_s(z)}{H(z)} dz \approx 147 \text{ Mpc} \quad (1.93)$$

This characteristic scale is imprinted in the distribution of matter as a preferred separation between galaxies, appearing as a distinct peak in the two-point correlation function $\xi(r)$ at $r \approx 147$ Mpc, or equivalently as oscillations in $P(k)$. The BAO scale thus defines a **standard ruler** that can be measured at different redshifts to constrain the expansion history of the Universe through $H(z)$ and the angular diameter distance $D_A(z)$ ([Eisenstein et al., 2005], [Blake and Glazebrook, 2003]).

4.2.3 Redshift Space Distortions

Galaxies are observed in redshift space rather than real space. Their peculiar velocities along the line of sight distort their apparent positions, producing anisotropies in the observed galaxy clustering. On large scales, coherent infall of galaxies into overdensities causes an apparent compression of structures along the line of sight, known as the **Kaiser effect** ([Kaiser, 1987]). On small scales, the virial motions of galaxies within collapsed structures produce an elongation of structures along the line of sight, known as the **Fingers of God** effect ([Jackson, 1972]).

The amplitude of these distortions depends on the rate of structure growth, encapsulated in the parameter $f\sigma_8$, where $f = d\ln D_1/d\ln a \approx \Omega_m(z)^{0.55}$ is the growth rate predicted by General Relativity and σ_8 is the amplitude of matter fluctuations on scales of $8h^{-1}$ Mpc. Modified gravity theories predict different values of f , making measurements of $f\sigma_8$ a direct test of GR on cosmological scales ([Guzzo et al., 2008], [Aghanim et al., 2020]).

4.3 Type Ia Supernovae

As discussed in Section 3, it is possible to obtain the distance of an object through its luminosity. However, measuring the intrinsic luminosity of celestial objects is a difficult task in itself. Type Ia Supernovae are thermonuclear explosions involving white dwarf stars that are **standardizable candles**, meaning they're characterized by a reproducible intrinsic luminosity. By measuring their apparent brightness as a function of redshift, one can reconstruct the luminosity distance-redshift relation $d_L(z)$ and constrain the expansion history of the Universe. This led to the discovery of the accelerated expansion of the Universe ([Riess et al., 1998], [Perlmutter et al., 1999]). A detailed discussion of Type Ia Supernovae and their use as cosmological probes is presented in Chapter 2.

4.4 Gravitational Lensing

According to general relativity, mass curves spacetime and light follows the geodesics of that curved geometry. As a result, the path of photons is deflected when passing near massive objects, a phenomenon known as gravitational lensing. In a cosmological context, this deflection can be derived from the geodesic equation applied to the perturbed FLRW metric, and it provides a direct probe of the total matter distribution, both baryonic and dark, along the line of sight.

Two main regimes of gravitational lensing are relevant for cosmology. **Strong lensing** occurs in the vicinity of objects with deep gravitational potentials, such as massive galaxy clusters, producing dramatic and easily identifiable distortions such as multiple images, giant arcs, and Einstein rings. It provides precise measurements of the mass distribution of individual clusters. **Weak lensing** operates in the regime where the distortion of any individual background source is too small to be detected — the induced ellipticity is much smaller than the intrinsic ellipticity of a galaxy [Bartelmann and Schneider, 2001]. However, the lensing signal can be recovered statistically by measuring the coherent distortion of the shapes of large samples of background galaxies, an effect known as **cosmic shear** γ .

Since lensing responds to all matter along the line of sight, cosmic shear surveys provide a direct measurement of the projected matter distribution. The amplitude of the cosmic shear signal is sensitive to both the amplitude of matter fluctuations σ_8 and the matter density Ω_m , but these two parameters are largely degenerate in lensing observations. This degeneracy is characterised by the parameter $S_8 = \sigma_8(\Omega_m/0.3)^{0.5}$, which current surveys constrain to percent-level precision ([Heymans et al., 2021a] [DES Collaboration et al., 2022a]).

4.5 The Lyman- α Forest

When observing the spectra of distant quasars, a dense series of absorption lines is observed, a feature known as the **Lyman- α forest**. These absorption lines are produced by neutral hydrogen clouds in the




Fig. 1.10 A comparison of the marginalized parameter constraints in the Λ CDM model from the Dark Energy Survey ([DES Collaboration et al., 2022a]) with predictions from Planck CMB data (no lensing; green). There is also the fiducial cosmology 3x2pt (black) and the combined Y3 3x2pt and Planck (orange) results.

intergalactic medium (IGM) through which the quasar photons pass on their way to the observer, typically at redshifts $2 \lesssim z \lesssim 5$. Each absorbing cloud at redshift z produces an absorption line at an observed wavelength:

$$\lambda_{obs} = 1216(1+z) \text{ \AA} \quad (1.94)$$

where $1216 \text{ \AA}$ is the restframe Lyman- α wavelength. It's possible to show that the neutral hydrogen fraction in the IGM traces the underlying total matter density field, making the Lyman- α forest an effective map of the matter distribution along the line of sight ([Gunn and Peterson, 1965]).

The cosmological information in the Lyman- α forest is extracted through the power spectrum of

the transmitted flux fluctuations, which can be directly related to the three-dimensional matter power spectrum ([Croft et al., 1998]). A particularly powerful feature of this probe is that it is sensitive to scales $k \sim 0.1\text{--}10 h\text{Mpc}^{-1}$ at high redshift, well beyond the reach of galaxy surveys, making it highly complementary to both CMB and galaxy clustering measurements. This sensitivity to small scales and high redshifts allows the Lyman- α forest to constrain the sum of neutrino masses $\sum m_\nu$ through the small-scale suppression of the power spectrum ([Palanque-Delabrouille et al., 2015]), the spectral index n_s and the BAO standard ruler at $z \sim 2.3$ ([Delubac et al., 2015]).




Fig. 1.11 High resolution spectrum of the Lyman- α forest part of a quasar measured at $z = 3.63$ taken with the HIRES spectrograph on the 10 m Keck telescope in Hawaiï.

5 Tensions in The Cosmological Model

The previous sections served to introduce and demonstrate the power of the Λ CDM model in describing the Universe. Nonetheless, tensions have emerged between cosmological parameter measurements obtained from different probes, suggesting that the discrepancies may be of physical or systematic origin. Here we present the three main tensions.

5.1 H_0 Tension

The Hubble constant H_0 can be inferred in two ways: through direct measurements via the cosmic distance ladder, or indirectly through model-dependent fitting of early-Universe observables. The direct measurement is obtained by calibrating standard candles such as Cepheid variables and Type Ia Supernovae. The latest results from the SH0ES collaboration give $H_0 = 73.04 \pm 1.04 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ([Riess et al., 2022]). The indirect method relies on fitting the acoustic peak structure of the CMB angular power spectrum within the Λ CDM framework. The Planck 2018 collaboration reports $H_0 = 67.4 \pm 0.5 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ([Aghanim et al., 2020]), yielding a $\sim 5\sigma$ discrepancy with the SH0ES result. This tension persists across multiple independent measurements of H_0 ([Di Valentino et al., 2021]), raising the question of whether it reflects unknown systematic errors, or new physics beyond Λ CDM present at different epochs of the Universe.




Fig. 1.12 Probability density functions of H_0 from different measurements probed at different redshifts, taken from [Freedman, 2021]. The CMB measurements prefer a lower H_0 value while Cepheids prefer higher values.

5.2 S_8 Tension

The parameter $S_8 = \sigma_8 \sqrt{\Omega_m/0.3}$ combines the amplitude of matter fluctuations σ_8 with the matter density Ω_m , and characterises the growth of large-scale structure. It can be measured both through the spatial correlations of galaxies and weak gravitational lensing, and through the CMB anisotropy power spectrum, where it is related to the primordial amplitude A_s through the linear growth of structure. The Planck 2018 collaboration measures $S_8 = 0.832 \pm 0.013$ ([Aghanim et al., 2020]), while the Dark Energy Survey Year 3 measures $S_8 = 0.776^{+0.017}_{-0.017}$ from a combination of cosmic shear, galaxy clustering, and galaxy-galaxy lensing ([DES Collaboration et al., 2022b]), indicating a 2.5σ discrepancy.




Fig. 1.13 S_8 and Ω_m contours for different galaxy surveys and CMB taken from [Heymans et al., 2021b] enclosing the 68% and 95% posterior probability. It is highlighted that CMB measurements prefer higher S_8 values to galaxy surveys.

5.3 (w_0, w_a) Tension

Within the Λ CDM paradigm, the accelerated expansion of the Universe is driven by a cosmological constant with equation of state $w = -1$, corresponding to a constant dark energy density $\rho_\Lambda = \text{const.}$ If instead the dark energy density evolves with time, the simplest extension is the Chevallier-Polarski-Linder

(CPL) parametrisation ([Chevallier and Polarski, 2001], [Linder, 2003]):

$$w(a) = w_0 + w_a(1 - a) \quad (1.95)$$

where w_0 is the present-day equation of state and w_a parametrises its time evolution. The Λ CDM limit is recovered for $w_0 = -1$ and $w_a = 0$.

Recent results from DESI DR2, when combined with Planck CMB data and Type Ia Supernova datasets, show deviations from Λ CDM in the (w_0, w_a) plane ([DESI Collaboration et al., 2025]). The significance of the deviation depends on the supernova sample used: 2.8σ when combined with Pantheon+, 3.8σ with UNION3, and 4.2σ with DES Year 5 ([DESI Collaboration et al., 2025]). A subsequent recalibration of the DES Year 5 supernova sample ([Popovic et al., 2025]) reduced this DES discrepancy to 3.2σ .




Fig. 1.14 w_0 and w_a contours from the latest [DESI Collaboration et al., 2025] measurements enclosing the the 68% and 95% posterior probability when combined with Pantheon+, UNION3 and DES year 5 data and CMB.

6 Physical explanations for the accelerated Universe

Since the discovery of the acceleration of the Universe by [Riess et al., 1998] and [Perlmutter et al., 1999], the quest to understand the underlying nature of this acceleration has seen many theories arise on what it could be. We've seen one way this can be done by via the addition of a cosmological constant that introduces a new type of energy in the Einstein equations. Due to the fact that we do not know enough about the nature of this energy it is called "dark energy". In the next subsections we will be discussing some of the theoretical models that were proposed to explain this acceleration.

6.1 Cosmological constant

The simplest way to account for the acceleration as we have mentioned before is by adding a cosmological constant Λ to the Einstein equations 1.5, which is what is done in the Λ CDM model. We can look at Λ in two different lights, either a modification to the geometry of spacetime or as a part of the energy content of the Universe. Mathematically they are equivalent since the energy-momentum tensor $T_{\mu\nu}$ can include the $\Lambda g_{\mu\nu}$ and their equations of state are indistinguishable in Λ CDM.

6.2 Dynamical dark energy

One could naturally assume that if the acceleration of the Universe is driven by a fluid that its energy density would change throughout time. The common way to model this is with the Chevallier-Polarski-Linder (CPL) model ([Chevallier and Polarski, 2001]) characterized by two parameters w_0 and w_a such that:

$$w(z) = w_0 + w_a \frac{z}{1+z} = w_0 + w_a(1-a) \quad (1.96)$$

Therefore the Hubble parameter becomes:

$$H(z) = H_0 \sqrt{\Omega_m(1+z)^3 + \Omega_r(1+z)^4 + \Omega_k(1+z)^2 + e^{\left(-3w_a \frac{z}{1+z}\right)} \Omega_\Lambda(1+z)^{3(1+w_0+w_a)}} \quad (1.97)$$

Λ CDM then becomes a particular case where $w_0 = -1$ and $w_a = 0$.

6.3 Quintessence field

The quintessence model uses an approach similar to inflation where one assumes the accelerated expansion is driven by a scalar field ϕ , here called the quintessence field. The action associated to that field is written as:

$$S(\phi) = \int d^4x \sqrt{-g} \left(-\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) \right) \quad (1.98)$$

The equations of motion of ϕ are determined by the Klein-Gordon equation:

$$\ddot{\phi} + 3H\dot{\phi} + \frac{d}{d\phi} V = 0 \quad (1.99)$$

For a FLRW metric, the pressure and energy density become:

$$P_\phi = \frac{1}{2} \dot{\phi}^2 - V(\phi) \quad (1.100)$$

$$\rho_\phi = \frac{1}{2} \dot{\phi}^2 + V(\phi) \quad (1.101)$$

Giving us the following equation of state:

$$w = \frac{P_\phi}{\rho_\phi} = \frac{\frac{1}{2}\dot{\phi}^2 - V(\phi)}{\frac{1}{2}\dot{\phi}^2 + V(\phi)} \quad (1.102)$$

Under a slow-roll approximation, the field evolves slowly such that $\dot{\phi} \simeq 0$ giving us $w \approx -1$ where we recover the results of a cosmological constant. If we consider a fast rolling field where the potential is negligible in front of the evolution, the equation of state becomes $w \approx 1$ however this is not physical since we know that for an accelerating Universe $w < -\frac{1}{3}$. Results from [DESI Collaboration et al., 2025] show a preference for a dynamical dark energy in the CPL parameterization.

6.4 Modified Theories of Gravity

An alternative to dark energy is to modify the theory of GR on cosmological scales. We call the models that do so Modified Gravity models (MG). Multiple classes of MG theories exist each modifying a certain aspect of GR. We will be presenting some of those models in each class.

Additional scalar fields

The first class consists of adding an additional scalar field to the Einstein-Hilbert action. These fields need to have equations of motions of second-order time derivatives at most to avoid any unphysical solutions.

One of the models that do that are the $f(R)$ models ([Sotiriou and Faraoni, 2010]) which replaces the Ricci scalar R by a general function of R . The action for an $f(R)$ gravity becomes:

$$S = \frac{c^4}{16\pi G} \int \sqrt{-g} f(R) d^4x + \int \sqrt{-g} \mathcal{L}_m d^4x \quad (1.103)$$

And the field equations are:

$$f'(R)R_{\mu\nu} - \frac{1}{2}f(R)g_{\mu\nu} + [g_{\mu\nu}\square - \nabla_\mu\nabla_\nu]f'(R) = \frac{8\pi G}{c^4}T_{\mu\nu} \quad (1.104)$$

For the simplest case $f(R) = R - 2\Lambda$ we recover GR. For a FLRW metric, the Friedmann equations become (with $c = 1$):

$$3f'H^2 = 8\pi G\rho_m + \frac{1}{2}(f'R - f) - 3H\dot{f}' \quad (1.105)$$

$$-2f'\dot{H} = 8\pi G(\rho_m + P_m) + \ddot{f}' - H\dot{f}' \quad (1.106)$$

If one rewrite Equation 1.105 to match what we get in GR $3H^2 = 8\pi G\rho$, we can identify a and effective "curvature fluid" that under certain conditions would cause an accelerated Universe. This gives us an effective equation of state on w_{eff} :

$$w_{eff} = -1 + \frac{1}{3f'H^2} [8\pi G(\rho_m + P_m) + \ddot{f}' - H\dot{f}'] \quad (1.107)$$

Since acceleration is possible for $w < -\frac{1}{3}$ this means that a condition on any $f(R)$ theory would be:

$$8\pi G(\rho_m + P_m) + \ddot{f}' - H\dot{f}' < 2f'H^2 \quad (1.108)$$

$f(R)$ must also be able to reproduce GR during the early Universe and at high matter density regions to remain consistent with observational results. This screening effect is called the chameleon mechanism

and the most well known $f(R)$ theories do possess it.

Massive gravity

In quantum field theory, forces are mediated by particles — for instance photons for electromagnetism and gluons for the strong force. By analogy, gravity is expected to be mediated by a spin-2 particle called the graviton. The graviton is predicted to be massless particle with an infinite interaction range, that can recover GR in a quantum field theory formalism. If we now assume that the graviton does in fact possess a mass, that would decrease the range of its gravitational interaction as well as its strength between distant objects ([Fierz and Pauli, 1939], [de Rham et al., 2011]). The gravitational potential is then modified as follows:

$$\Phi = -\frac{GM}{r}e^{-mr} \quad (1.109)$$

where r is the distance to an object, M the mass of the object and m the mass of the graviton. However, the introduction of a massive gravitons produces multiple instabilities such as the inability to recover GR when m goes to 0. It also suffers to produce flat FLRW solutions.

Breaking assumptions

The last class of models we will present here are the ones that break physical assumptions. Here we present an example of a model proposed by Gia Dvali, Gregory Gabadadze and Massimo Porrati ([Dvali et al., 2000]) where gravity can propagate in a 5-dimensional space time with an additional spatial dimension. We can think of it as our 4D world embedded in an infinite volume 5D space. The idea being that the gravity on large scales appears to be in 5D while on small scales in 4D, this is due to the Vainshtein effect. The action for such a theory looks like :

$$S = 2M_5^3 \int d^5X \sqrt{-\tilde{g}} \tilde{R} + 2M_{pl}^2 \int d^4x \sqrt{-g} R + \int \sqrt{-g} \mathcal{L}_m + S_{GH-} + S_{GH+} \quad (1.110)$$

where M_5 is the Planck mass M_{pl} in 5D, $\tilde{g}$ and $\tilde{R}$ are the norm of the metric and the Ricci scalar in 5D. The S_{GH-} and S_{GH+} are Gibbons-Hawking terms that serve as boundary terms commonly found in, but not limited to, quantum gravity theories.

The theory does predict a late time acceleration ([Deffayet, 2001]), however it suffers theoretical instabilities and unphysical solutions ([Koyama, 2008]). Observations have also ruled the theory out DGP theories ([Fang et al., 2008]).

CHAPTER 2

Cosmology with type Ia Supernova

Contents

1	Type Ia Supernova	44
1.1	Classification	44
1.2	Explosion mechanisms	44
1.3	Explosion rate	47
1.4	Observations	47
2	Cosmology with distances	52
2.1	Hubble diagram	52
2.2	Peculiar velocities	53
3	Distance measurements from observed fluxes	57
4	Standardization	58
4.1	Brighter-slower effect	58
4.2	Brighter-bluer effect	58
4.3	Environmental effects	59
5	Spectrophotometric models of supernovae	61
5.1	SALT2/SALT3	61
5.2	Models trained on the SNfactory dataset	64
6	Main goal of this PhD	69

1 Type Ia Supernova

For many centuries, astronomers all over the globe would observe the emergence of a bright light in the sky thinking it was the birth of a new star in the Universe. In 1572, an astronomer by the name of Tycho Brahe developed an interest in such event and named the phenomenon *nova stella*, latin for new star. His work inspired later astronomers such as Kepler and eventually led to the Copernican revolution.

It wasn't until 1934 when Fritz Zwicky and Walter Baade when the term supernova (from the latin "super" and "novae" meaning new bright stars) was coined to describe the brightest novae observed. The supernova classifies phenomena related to the end of a star's life cycle when they go through explosions. This process is temporary therefore taking part in the class of **transients**. Their fluxes are initially weak and then increases over time to a point where it becomes detectable from the Earth, then it diminishes slowly till it disappears.

In the next sections we will be explaining the different classifications of supernovae as well as our understanding behind their explosion mechanisms. A particular focus will be on a specific class of supernovae called "type Ia" and why they are heavily studied by cosmologists.

1.1 Classification

Two main categories of supernovae exist: type I and type II each with their own subcategories of supernovae. These main categories were discovered by Rudolph Minkowski and Walter Baade in 1941 when studying the spectral features of 14 supernovae. Type I supernovae's absorption spectra is characterized by having no Hydrogen, contrary to type II. Type I supernovae are then divided into three classes ([Wheeler and Levreault, 1985, Elias et al., 1985], [Wheeler and Harkness, 1990]):

- **Ia**: contains Silicon line at 615.0 nm
- **Ib**: does not contain Silicon and contains Helium line at 587.6 nm
- **Ic**: does not contain Silicon or Helium lines

Type II supernovae are divided in following way ([Hillier, Desmond John and Dessart, Luc, 2019]):

- **IIn**: contains narrow absorption lines
- **IIf**: contains Helium like Ib
- **IIP**: Hydrogen dominates the spectra and its integrated flux becomes constant
- **IIl**: Hydrogen dominates the spectra and its integrated flux decreases in time

The classification of supernovae can be summarized in Figure 2.2.

1.2 Explosion mechanisms

Different explosion mechanisms and progenitors exist for the different classes of supernovae. We will be separating them into two cases: core-collapse and thermonuclear.

Core collapse

This explosion concerns supernovae of types Ib, Ic and all types II. The occurrence of this explosion happens at the end of the life cycle of a star after it has completed the fusion of its light elements. The star




Fig. 2.1 Photometric and spectroscopic curves for different classes taken from [Baumont, 2007]. The left are the schematics for the integrated fluxes (lightcurves) for SN Ia (blue), SN Ib or Ic (red), SN IIP (green) and SN IIL (cyan). On the right we have the spectra of three supernovae observed in 1999, SN Ia (blue), SN IIP (red) and SN Ic (green).

then does generate enough energy to compensate for the gravitational pull and then collapses on itself. The material of the star collapses until it reaches scales at which it is stopped by the strong nuclear interaction, which defines the limiting distance between atoms. This causes a collision between the different layers of the star creating a shock wave that propulses the star material and ignites the supernova explosion. If the star has a mass inferior to $\sim 3.3M_{\odot}$ (Oppenheimer-Volkoff limit [Oppenheimer and Volkoff, 1939]), where $\odot$ denotes the Solar mass, the star will become neutron star. Larger mass stars turn to black holes.

Thermonuclear

Thermonuclear explosions concern type Ia supernovae. This occurs in white dwarf stars comprised primarily of Oxygen and Carbon when their masses approach the Chandrasekhar limit of $\sim 1.44M_{\odot}$ ([Chandrasekhar, 1931]). White dwarf stars form from the remains of low mass stars (typically under $8M_{\odot}$) and are dense degenerated systems. This term comes from the fact that the density of a white dwarf is limited by electron degeneracy pressure that comes from the Pauli exclusion principle which states that two electron within the same energy levels cannot share the same quantum state. Accreting more mass from external sources induces Carbon and Oxygen fusion which increases the temperature of the white dwarf stars. If the accretion keeps on happening, the fusion of Carbon and Oxygen becomes a chain reaction at a constant pressure causing a thermonuclear explosion ([Röpke and Hillebrandt, 2004]). The explosion ends up releasing around 10^{51}J of energy, mainly kinetic energy. Only a fraction of that is turned into luminous energy of 10^{44}J ([Khokhlov et al., 1993]) meaning type Ia supernovae have a luminosity at the same order of magnitude of its host galaxy.

Observations such as SN2011fe have demonstrated that type Ia supernovae are originated from compact and dense stars, which are white dwarfs. It also important to note that type Ia supernova spectra are characterized by Silicium and Calcium absorption lines which are the product of the Carbon and Oxygen fusion. Due to the fact that type Ia supernovae occur in quasi-identical environments, we tend to observe similar luminosities between them. Finally, they are observed in environments with older stars which is justified due to the fact that white dwarfs happen at the final stages of a star.




Fig. 2.2 Classification of supernovae. The left column are type I (blue) and the right column are type II (red).

White dwarf stars aren't very luminous to be observed at long distances. Usually they are detected a short while after their explosions. Therefore the progenitors of type Ia supernovae have never been observed. Multiple scenarios exist to explain the explosions. We will only be presenting the two main scenarios:

- **Single degenerate** ([Nomoto, 1982]): This occurs when white dwarfs accrete mass from nearby more massive non-degenerate stars with low surface gravity, like red giants. Arguments against that scenario is that the accreted Helium and Hydrogen must all be transformed into Carbon and Oxygen with a fusion reaction, which would explain why they do not appear as absorption line in the spectrum of the supernova. The other is that the speed of the accretion can neither be too slow nor too fast or the explosion doesn't happen. Lastly, the companion stars weren't observed near the explosion area ([Li et al., 2011]).

- **Double degenerate** ([Iben and Tutukov, 1984]): This occurs when a white dwarf star accretes its mass from a nearby less massive white dwarf star in close orbit. This scenario does explain the absence of Helium and Hydrogen absorption lines and the observation of superluminous supernovae (like [Andrew Howell et al., 2006]). It also allows for the case of a double detonation which when it happens for two white dwarfs whose masses are inferior to the Chandrasekhar mass can explain the existence of subluminous supernovae ([Tanikawa et al., 2019]).

1.3 Explosion rate

The explosion rate of type Ia supernovae is the measurement of the number of explosion at a certain redshift in year. It can be divided into two categories: low redshift up to $z \sim 0.3$ and high redshift. At low redshift the explosion is nearly constant and barely evolves. Low redshift samples such as ZTF allow us to measure that rate. A study done by [Perley et al., 2020] on the ZTF sample, assuming an isotropic distribution, have measured a rate of

$$R_{ZTF} = (2.35 \pm 0.24) \times 10^4 \text{Gpc}^{-3} \cdot \text{yr}^{-1} \quad (2.1)$$

Where each supernova was weighted by the volume it probes with a varying limiting absolute magnitude. However, a recent paper by [Lordet et al., 2026] showed that the rate of supernova explosion might be more complex than anticipated and is much more dependent on its environment. Figure 2.3 highlights that the assumption that the rate is homogenous at low redshift isn't sufficient to describe supernova as we observe an excess at $z \sim 0.035$ while Figure 2.4 shows that the excess of supernova at that redshift seems to be located around extremely dense regions.

At high redshift the rate becomes very dependent on the redshift. Its behaviour can be described by a power law with respect to $(1+z)$. Estimates by combining a multitude of probes in [Frohmaier et al., 2019] have shown that the rate can be described as:

$$R(z) = (2.27 \pm 0.19) \times 10^4 \times (1+z)^{(1.70 \pm 0.21)} \text{Gpc}^{-3} \cdot \text{yr}^{-1} \quad (2.2)$$

1.4 Observations

Spectroscopic

Type Ia supernova spectra have a distinct spectra. Most of the light emitted from a type Ia is in the visible wavelengths with some parts of their spectra in the infrared and ultraviolet. During the initial phases of the explosion the external layers of the explosion are opaque. Near the maximum emission of light, we expect to see absorption lines of Silicon and Calcium caused by the fusion of Carbon and Oxygen. The days that follow, the opacity decreases and we start to see absorption lines from Nickel and Cobalt. Finally, in the following weeks the Nickel and Cobalt will disintegrate into Iron for which we see the absorption lines.

Photometric

When describing the photometric evolution of type Ia supernovae, we are describing the integrated flux in a observation band or filter that covers a specific wavelength. Generally, we use the standard filters U, B, V, R and I (also known as the Bessel filters [Bessell, 1990]) that cover the ultraviolet, visible and near-infrared range. We observe in all bands an increase the lightcurve that lasts approximately 20 days




Fig. 2.3 Distribution of simulated supernova observed as a function of redshift z vs the ZTF DR2 sample of observed supernovae (circle with black uncertainty). The different simulations are done using different priors on their positions. Red assumes a uniform sampling like [Perley et al., 2020]. Blue uses a uniform sampling weighted by halo masses from Manticore-local. Orange assumes a redshift dependent rate. Green assumes a redshift dependent rate and weights by the halo masses.

with an absolute magnitude in the B band of -19.5mag that occurs following the disintegration of Nickel. After that, the visible and ultraviolet bands measure an exponential decrease due to the disintegration into Iron. In the infrared bands, we observe a second peak somewhere between 20 to 25 days after the first peak.




Fig. 2.4 Distribution of supernova observed vs the dark matter distribution posterior from Manticore-local as well as the local supernova rate. Top left corner we have a zoom in on the Hercules supercluster. Top right corner we have a zoom in on the Leo supercluster.




Fig. 2.5 Spectral series of SN 2011fe supernova taken by with the SNIFS instrument ([Pereira et al., 2013]). The spectra were captured between -15 days to 50 days after the maximum emission data. They are ordered from top to bottom, or blue to red. The magnitude shown are relative to the maximum magnitude.




Fig. 2.6 Synthetic lightcurves of the SN 2009ig supernova in the UBVRI bands made for the SNfactory collaboration taken from [Chotard, 2011].

2 Cosmology with distances

2.1 Hubble diagram

The main tool used to constrain the expansion of the Universe with type Ia supernovae is the Hubble diagram. For supernovae, the Hubble diagram maps their luminosity distances to their redshifts. In practice what we attempt to do is relating the type Ia supernova's magnitude in the standard B band denoted as m_B . This magnitude is defined as:

$$m_B = -2.5 \log_{10} \left(\frac{f}{f_0} \right) \quad (2.3)$$

where f is the integrated flux in the observer frame B band and f_0 the flux of a reference star. We also define the absolute magnitude M of an object with an intrinsic luminosity L as the apparent magnitude if it was located at 10pc from the observer. Given the definition of the luminosity distance we can get:

$$M = -2.5 \log_{10} \left(\frac{L}{f_0 4\pi (10\text{pc})^2} \right) \quad (2.4)$$

We can finally define the distance modulus of a type Ia supernova as:

$$\mu = m_B - M \quad (2.5)$$

and from Equation 1.85 we can identify:

$$\mu = 5 \log_{10} \left(\frac{d_L}{10\text{pc}} \right) = 5 \log_{10}(d_L) + 25 \quad (2.6)$$

where d_L is the luminosity distance measured in Mpc. If we manage to determine μ from observations, we can construct a Hubble diagram by plotting μ against z .

2.1.1 Measurement of $\Omega_m - w$

Once we have access to the supernova Hubble diagram, we can start deducing cosmological parameters. We recall that under a flat w -CDM cosmology, Equation 1.88 becomes:

$$d_L = \frac{c(1+z)}{H_0} \int_0^\infty \frac{dz'}{E(z')} \quad (2.7)$$

with $E(z) = \frac{H(z)}{H_0}$ being the dimensionless Hubble parameter. Supernova measurements can go up to a redshift of 2, where the only contribution to the Hubble parameter are Ω_m and Ω_Λ . Given that $\Omega_m + \Omega_\Lambda = 1$ at that redshift, we can deduce that:

$$E(z) = \Omega_m(1+z)^3 + (1-\Omega_m)^{3(1+w)} \quad (2.8)$$

We can see that if we then adjust the Ω_m and w parameters and integrate Equation 2.7 we can compare our observations with the predicted values and constrain w and Ω_m . The value of H_0 for such analyses does not directly affect the values of w and Ω_m and it is usually absorbed in a generalized constant that contains M as well. Ω_m and w are also degenerated so with supernova alone we do a joint inference of the parameters. In order to break this degeneracy, combinations with other probes such as BAO measurements are done. Finally, doing this requires supernova accross a large range of redshift (a low- z sample at $z \lesssim 0.1$ and a high- z sample $0.1 \lesssim z \lesssim 2.0$) in order to observe the evolutions of Ω_m and Ω_Λ .

2.1.2 Measurement of H_0

We recall that for objects at a low redshift the distance luminosity is written as Equation 1.90. We can then express the magnitude as a function of the redshift with:

$$m \approx A - 2.5 \log_{10}(LH_0^2) + 5 \log_{10}(z) \quad (2.9)$$

where A is a constant. We can see that the relationship between the magnitude and redshift naturally introduces the Hubble parameter H_0 , however it is degenerated with the intrinsic luminosity of supernovae L . Breaking this degeneracy requires us to calibrate the intrinsic supernova luminosity. We commonly use the "distance ladder" method that uses Cepheid stars in the Milky-Way as distance calibrators.

Cepheids are a class stars that have variable periodic luminosities, which was firstly inferred by Henrietta Swan Leavitt thus establishing the Leavitt relation ([Leavitt, 1908]). Most recent confirmations of this relation were observed by [Breuval et al., 2025a]. Distances of Cepheids can be determined using "parallax": a method that measures the apparent displacement angle of an object with respect to a static background due to the motion of the observer. The application of this method is done by measuring the positions of the Cepheids with respect to a background of distant stars 6 months apart. The displacement angle of the observed Cepheid measured due to the half-rotation of the Earth around the Sun can be expressed as $\theta = \frac{r}{d}$ where r is the distance to the Cepheid and d the Earth-Sun distance. This means that the measurement of the distance luminosity of Cepheids can be calibrated using objects sufficiently close to the Milky-Way for their distance to be measured by parallax.

If we can identify nearby galaxies that contain both Cepheids and type Ia supernovae, we can use the Leavitt relation to measure the Cepheid distance modulus μ_{Cepheid} , which in turn calibrates the absolute magnitude M of the supernova:

$$M = m_B - \mu_{\text{Cepheid}} \quad (2.10)$$

where m_B is the observed peak magnitude of the type Ia supernova in the rest-frame B band. This breaks the degeneracy between L and H_0 , as M is now determined independently of the Hubble expansion. Applying this calibrated M to type Ia supernovae at low redshift, we can compute the luminosity distance d_L from the observed magnitude and directly infer H_0 from Equation 1.90, averaged over a large sample of supernovae. The most recent measurement using this method yields $H_0 = 73.04 \pm 1.04 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ([Riess et al., 2022]).

2.2 Peculiar velocities

We define the peculiar velocity of a celestial object as the velocity due to its proper motion independent from the Hubble expansion. This is particularly important for cosmological analyses as it can be used as a test for GR. If we assume peculiar velocities originate from the infall into gravitational potential wells then the detection of correlations between peculiar velocities directly probes the "strength" of gravity at cosmological scales. The parameter probed here is the growth rate of structures $f(z)\sigma_8(z)$ which combines the growth rate function f as well as the matter fluctuations in a 8 Mpc box σ_8 . A common parameterization of the growth rate of structures in Λ CDM is:

$$f(z)\sigma_8(z) = \Omega_m(z)^\gamma \sigma_8(z) \quad (2.11)$$

Where γ is the growth index. For GR γ is expected to be 0.55 therefore deviations from this value will indicate that GR is not the most suitable gravitational theory for the description of the large scale Universe.




Fig. 2.7 Original measurements by Leavitt establishing the magnitude period relation adjusted in [Breuval et al., 2025b].

2.2.1 Extracting peculiar velocities

Supernova peculiar velocity information is encoded in the observed redshift. If we define z_{pv} as the redshift due to the proper motion of the supernova and z_{cosmo} as the redshift due to the cosmic expansion, we get the following formula for the observed redshift z_{obs} :

$$1 + z_{obs} = (1 + z_{cosmo})(1 + z_{pv}) = (1 + z_{cosmo})\left(1 + \frac{v}{c}\right) \quad (2.12)$$

Where v is the peculiar velocity. Measuring z_{pv} requires knowledge of z_{cosmo} which is not directly measurable. It can be deduced however once the expansion history has been determined. To put simply, if you know exactly the expansion history, then one can directly deduce the cosmological redshift of an object based on how far away it is. This is illustrated in Figure 2.8. This entails that it is crucial to be able to accurately measure the distances of supernovae. Moreover, high redshift supernovae redshifts are almost entirely dominated by the cosmological redshift so analyses on supernova peculiar velocities will be done on low redshift samples.




Fig. 2.8 Illustration of the impact of peculiar velocities on the distance modulus on a Hubble diagram from [Carreres et al., 2023]. The velocities were drawn randomly from a Gaussian distribution. The blue and red colors indicate the radial velocity of the supernova.

2.2.2 Measurement of $f\sigma_8$

Multiple methods in inferring $f\sigma_8$ exist. We present here four different methods that exist.

Maximum likelihood estimation [Gordon et al., 2007, Carreres et al., 2023]

The maximum likelihood estimation (MLE) is a statistical method of estimating parameters given a model and a statistical distribution. The likelihood function measures how well the model explains the data by assimilating the statistical distribution. For MLE, the distribution is assumed to be Gaussian as the variable is the distance modulus of the supernova, a continuous parameter. The likelihood can be written as:

$$\mathcal{L} \propto |\mathbf{C}(\theta)|^{-\frac{1}{2}} \exp\left(-\frac{1}{2} \mathbf{v}(\theta)^T \mathbf{C}(\theta)^{-1} \mathbf{v}(\theta)\right) \quad (2.13)$$

where θ are the model parameters, $\mathbf{v}$ is the peculiar velocity calculated as the difference between the observed distance modulus and the predicted distance modulus defined by a model, $\mathbf{C}$ is the covariance matrix which can be divided into two parts $\mathbf{C} = \mathbf{C}_{\text{noise}} + \mathbf{C}_{\text{PV}}$ where the second term directly depends on the value of $f\sigma_8$ and is left as a free parameter to be determined. The goal is to either maximize $\mathcal{L}$ or minimize its logarithmic value $\log \mathcal{L}$ which can be numerically beneficial.

Compressed two-point statistics

The compressed two-point statistics is done by adjusting the amplitude of the theoretical 2 point velocity-velocity power spectrum to the data. If we would assume a Λ CDM cosmology, the power spectrum is given by:

$$\mathcal{P}_{vv}(k) \propto (f\sigma_8)^2 \int dk \mathcal{P}(k) j_1(kr)^2 \hat{\mathbf{r}} \hat{\mathbf{r}}' \quad (2.14)$$

where j_1 is the first order spherical Bessel function and $\mathcal{P}$ is the matter power spectrum. For a linear approximation we obtain:

$$\mathcal{P}_{vv}(k) \propto (f\sigma_8)^2 \frac{\mathcal{P}(k)}{k^2} \quad (2.15)$$

What remains is to then adjust the $f\sigma_8$ value to match the observed power spectrum. If the density field is indeed Gaussian, then the full power spectrum would contain all of the information on the distribution. However at small scales it is not the case.

Density field reconstruction

The density field reconstruction infers the velocity field of matter from a galaxy density field. If we call δ_g the galaxy density field and b the galaxy bias (term that models how well do galaxies trace matter) then the reconstructed velocity field becomes:

$$\mathbf{v} = \frac{aH\beta}{4\pi} \int d^3\mathbf{x}' \frac{\mathbf{x}' - \mathbf{x}}{|\mathbf{x}' - \mathbf{x}|} \delta_g(\mathbf{x}') \quad (2.16)$$

where $\beta = \frac{f}{b}$. Comparing this to the velocity field inferred from type Ia supernovae, one can determine β . If then this is combined with external measurements of $\sigma_{8,g} = b\sigma_8$, we can compute the growth rate of structures with $f\sigma_8 = \beta\sigma_{8,g}$. This has been done in [Lavaux and Hudson, 2011].

Field level forward modeling

The field level forward model treats the density and velocity field in itself as a free parameter. The idea is to evolve an initial simulated density field using a gravity model to obtain a final density field. The observables are modelled from the obtained field to compare to actual observations such as galaxies or supernovae.

$$\text{Initial field} \longrightarrow \text{Final field} \longrightarrow \text{Model observables} \Longleftrightarrow \text{Compare to observations}$$

Due to the nature of this method, the growth rate of structures can be directly determined via the final constrained simulation. The challenge of this method relies in multiple stages such as: determining an accurate gravity model while maintaining reasonable computational time and accurately modelling the relation between the observables and the density field.

3 Distance measurements from observed fluxes

The key information in doing a cosmological analysis with type Ia supernova is found in their restframe fluxes. The photometric fluxes as mentioned previously is a broadband flux, meaning it is integrated in a observation band. Due to the redshift of each supernova being different, these integrated zones will be different for an observer on Earth, which makes it difficult to compare the same physical quantities. This can be seen in Figure 2.9. The key is to be able to reconstruct the restframe flux from the observed flux in the same band. We use the restframe B band since it contains the maximum light emission from supernova. From now on we will call this band the B^* band. Once we have obtained the flux in the B^* , we can obtain the distance modulus.




Fig. 2.9 Examples of four spectra at different redshifts 0.1, 0.4, 0.7 and 1.0. The highlighted parts of the spectra represent restframe band would measure. The highlighted areas on the graph are the regions where the observer bands measure.

Although supernova lightcurves look alike, there are still a couple of variations between them. This variability can be largely accounted for using standardization parameters, hence type Ia supernovae are standardizable candles. Without taking those into effect, the dispersion that exists around peak magnitude in the standard B filter is around 40%. The most well known standardization relation is the Tripp relation ([Tripp and Branch, 1999]) which uses two standardization parameters named stretch and color. After standardization, the dispersion can go down to 15%. More complex standardization methods have

been explored, however they necessitate spectral knowledge as well.

4 Standardization

4.1 Brighter-slower effect

The brighter-slower effect is an effect observed in type Ia supernovae where the lightcurves tend to have a slower decline the higher the magnitude of the supernova ([Phillips, 1993]). We define the decline parameter Δm_{15} as the difference of magnitude of the supernova in B band at the date of maximum emission and 15 days later:

$$\Delta m_{15} = M_B(t = t_{max}) - M_B(t = t_{max} + 15) \quad (2.17)$$

which quantifies that effect. We can also define a highly correlated parameter named the *stretch*, or X_1 ([Guy et al., 2007]), when modelling the spectrophotometric evolution with empirical models that measures the width of the lightcurve.

This effect has been revised in [Ginolin et al., 2025b] where this effect was measured on a sample of over 900 supernovae from the ZTF survey. It was demonstrated that this effect in fact presents two regimes that depend on the value of the stretch. In other words, the scaling of the brighter-slower effect varies based on the value of X_1 . We refer to this variation as the "broken-alpha" effect.




Fig. 2.10 Binned Hubble residuals as a function of the stretch. The data is fitted to highlight the two regimes of the broken alpha relation [Ginolin et al., 2025b].

4.2 Brighter-bluer effect

In a similar way, the brighter-bluer effect characterizes how bluer supernovae tend to have a higher magnitude ([Hamuy et al., 1995, Riess et al., 1996]). We can define an associated parameter, the *color* or c ,

as the deviation of the difference of magnitude at the time of maximum emission between the B and V from the average supernova. When c is negative we call the supernova blue, and if it is positive, red. Part of the reason this effect exist is due to the absorption of light by the host galaxy and the intergalactic dust in the line of sight of the observer. Another reason is due to intrinsic properties related to the physical properties of the progenitor.

This effect was confirmed in [Ginolin et al., 2025a] where it was studied on the ZTF DR2 sample of over 900 supernovae. A scale dependent effect was proposed in the paper but found no statistical significance for it.




Fig. 2.11 Binned Hubble residuals as a function of color. The data is fitted to highlight the brighter-bluer effect ([Ginolin et al., 2025a]).

The Tripp relation combines both effects to standardize the observed magnitude with two parameters α and β :

$$m_{Tripp} = m_{obs} - \alpha X_1 + \beta c \quad (2.18)$$

where m_{Tripp} is the standardized magnitude in the B band using the Tripp relation.

4.3 Environmental effects

We have already mentioned that galaxy properties can affect the luminosity of supernovae as seen in the brighter-blue effect. However, studies conducted on large samples of supernovae have shown that the standardized luminosity also depends multiple additional properties of the host galaxy. This is seen as a "step" in the Hubble residuals that depend on a certain property of the galaxy. One possible explanation would be the mass of the galaxies([Sullivan et al., 2010]). This dependancy presents itself as an offset in the standardized when the population of host galaxies' masses are divided into two categories: more or less than $10^{10} M_{\odot}$. The mass step has shown a 3.9σ significance after correcting for stretch and color ([Sullivan et al., 2010]).

Another possible dependency would be the star formation rate in those galaxies ([Rigault et al., 2020]). Galaxies can be divided into two subgroups: high star formation galaxies and low star formation galaxies. This step factor has shown a 2.6σ significance after correcting for stretch and color.

The [Roman, M. et al., 2018] paper studied this dependency on the supernovae local $U - V$ restframe colors (limited to a certain radius around the supernova). This step factor has shown a 7.0σ significance. When the same study was conducted on 927 ZTF supernova with the local colors $g - z$, the significance was found to be around 18.2σ significance.

We can then add an extra nuisance parameter to the Tripp relation γ that accounts for that offset:

$$m_{Tripp} = m_{obs} - \alpha X1 + \beta c + \gamma \quad (2.19)$$

where γ is fitted alongside the standardization parameters.




Fig. 2.12 Binned Hubble residuals as a function of the star formation rate ([Rigault et al., 2020]).

5 Spectrophotometric models of supernovae

Given that the B^* fluxes of supernovae are not directly observable, we rely on spectrophotometric models that allow us to reconstruct the emitted fluxes. Theoretical supernova explosion models haven't been able to replicate the observed fluxes with a good precision as seen in [Blinnikov et al., 2006], [Röpke and Sim, 2018] and [Pakmor et al., 2024]. Instead we rely on empirical models to describe the spectrophotometric data. Here we present the main models that exist.

5.1 SALT2/SALT3

The Spectral Adaptive Lightcurve Template ([Guy et al., 2007]) class of spectrophotometric models are the current state of the art for cosmological analyses. The idea is to parameterize the model with two parameters that would represent the stretch and color in the Tripp relation. These models are particularly useful since they are determined and used on photometric data. Spectra are used to determine the spectral features of the model. Once the model has been obtained, one can directly use it on lightcurves to determine its parameters.

We define the restframe spectral flux density surface $\mathcal{S}(\lambda, p)$ the spectrophotometric evolution of type Ia supernova where p is the phase of the supernova, defined as the difference between the observation date and the date of maximum emission $p = t - t_{max}$ in days, and λ the wavelength in Å. The goal is to then obtain S .

We can define the spectral flux density as:

$$\phi_{spec}(\lambda, p) = \frac{1}{4\pi d_L^2(z)} \frac{1}{1+z} \mathcal{S}\left(\frac{\lambda}{1+z}, \frac{p}{1+z}\right) \quad (2.20)$$

With that we can also deduce the broadband flux observed through a pass-band $T(\lambda)$ as:

$$\phi_{phot}(p) = \frac{1}{4\pi d_L^2(z)} \frac{1}{1+z} \int \mathcal{S}\left(\frac{\lambda}{1+z}, \frac{p}{1+z}\right) \frac{\lambda}{hc} T(\lambda) d\lambda \quad (2.21)$$

Since the goal of doing supernova cosmology is reconstructing the restframe fluxes, we can rewrite the expression in the restframe of the supernova as:

$$\phi_{phot}(p) = \frac{1}{4\pi d_L^2(z)} (1+z) \int \mathcal{S}(\lambda, \frac{p}{1+z}) \frac{\lambda}{hc} T((1+z)\lambda) d\lambda \quad (2.22)$$

As mentioned previously, the luminosity distances of supernovae are defined using the emitted flux in the standard B band at the time of maximum emission, observed at 10pc from the supernova:

$$\Phi_B = \frac{1}{4\pi (10\text{pc})^2} \int \mathcal{S}(\lambda, p=0) \frac{\lambda}{hc} B(\lambda) d\lambda \quad (2.23)$$

By imposing the normalization of the model in the B^* band at phase 0:

$$\int \mathcal{S}(\lambda, p=0) \frac{\lambda}{hc} B(\lambda) d\lambda = 1 \quad (2.24)$$

And defining the following parameter:

$$X_0 = \Phi_B \frac{(10\text{pc})^2}{d_L^2(z)} \quad (2.25)$$

We obtain:

$$\phi_{spec}(\lambda, p) = X_0 \frac{1}{1+z} \mathcal{S}\left(\frac{\lambda}{1+z}, \frac{p}{1+z}\right) \quad (2.26)$$

$$\phi_{phot}(\lambda, p) = X_0(1+z) \int \mathcal{S}\left(\lambda, \frac{p}{1+z}\right) \frac{\lambda}{hc} T((1+z)\lambda) d\lambda \quad (2.27)$$

The SALT parameterization of $\mathcal{S}$ is ([Guy et al., 2007]):

$$\mathcal{S}(\lambda, p) = [M_0(\lambda, p) + X_1 M_1(\lambda, p)] 10^{0.4cCL(\lambda)} \quad (2.28)$$

This parameterization is a classic principal component analysis, or PCA, with the addition of a color law. The M_0 and M_1 surfaces along with the CL color law are common to all supernovae. M_0 is the average spectral surface of a type Ia supernova, which represents what your average spectrophotometric evolution of a type Ia supernova looks like. M_1 is a surface representing the main variations to M_0 . CL represents the color diversity of supernovae. In practice, the measured spectra obtained are only used to determine the spectral features of $\mathcal{S}$ as they are not as well calibrated as photometric data and are obtained from many different instruments for the same survey, making parameters such as X_0 determined primarily on the photometric data.




Fig. 2.13 The SALT2.4 M_0 surface with respect to the restframe wavelength and phase




Fig. 2.14 The SALT2.4 M_1 surface with respect to the restframe wavelength and phase

Determining the global parameters is what we called "training the model" while determining the individual supernova parameters is called "fitting the lightcurve parameters". The model training is iteratively on a sample of supernovae, the training dataset. The date of maximum emission of light is the first thing that is determined (t_{max}). The global surfaces are then fitted and the residuals are analyzed. The variability not captured by the model is then accounted for using an error model. The last two steps are then repeated until the model converges. This includes accounting for calibration uncertainties, where for each passband in the dataset requires it's own model training in order which made the propagation of the calibration uncertainties a difficult task. Once the model has been trained, it is published. It is then fitted on a cosmological dataset, different from the one used in the training, to obtain the X_0 , X_1 and c parameters.

The initial SALT2 model was trained on a dataset containing 52 low- z supernova from [Hamuy et al., 1996] and 121 high- z supernova from the first two years of observation from SNLS ([Astier et al., 2006]). A later retraining of the model was done in [Guy et al., 2010] where 207 new supernovae were added from the SDSS-II sample, giving us the SALT2.4 model. This model was used in the [Betoule et al., 2014] JLA cosmological analysis.

The SALT3 model is the current state of the art iteration of the SALT models ([Kenworthy et al., 2021]).




Fig. 2.15 The SALT2.4 color law CL surface with respect to the restframe wavelength

The entire training procedure was written in Python3 coding language and is publicly available for the community. The model was trained on a sample of 1083 supernovae, 2.5 times larger than SALT2. SALT3 improves the training by having consistent photometric handling between surveys and improved cross-calibration between surveys in order to reduce survey dependent systematics. The model was also trained on more high- z supernova adding data from PanSTARRS-1, DES and Pantheon, improving the handling of UV data regions. The color law was also readjusted to be a higher order polynomial than SALT2. The SALT3 was used for the DES cosmological analysis ([DES Collaboration et al., 2024]).

5.2 Models trained on the SNfactory dataset

The two-parameter standardization scheme introduced by [Tripp and Branch, 1999] manages to greatly reduce the dispersion between supernovae but still misses on some of the variability that is observed. In order to tackle this issue, spectrophotometric models have been developed with the aim of finding standardization parameters based on the spectroscopic features of supernovae. To do so, these models have been trained on the SNfactory dataset ([Aldering et al., 2002]). A survey that ran from 2004 to 2014 collecting a sequences of spectra of nearby supernovae at low- z (max $z \approx 0.11$) over a long period of time. This is the only dataset that contains a fine spectroscopic description in a large phase range. The well calibrated spectra in the SNfactory dataset will also mean that it doesn't need to be accompanied by photometric measurements to acquire distances.

These features makes this dataset unique and very interesting to use in order to study the properties of supernova.

5.2.1 SNEMO

SNEMO model (SuperNova Empirical MODEls, [Saunders et al., 2018]) generalises the PCA used in SALT2. In SNEMO, the $\mathcal{S}(\lambda, p)$ components are extracted using an Expectation Minimization for Factor Analysis or EMFA that takes into account the measurement uncertainties that affects the model. With SNEMO multiple model trainings can be done at different orders. The model is expressed at:

$$S_{SNEMO}(\lambda, p) = c_{SN,0} \left(e_0(\lambda, p) + \sum_{i=1}^N c'_{SN,i} e_i(\lambda, p) \right) 10^{0.4A_s C(\lambda)} \quad (2.29)$$

where c_{SN} are the individual supernova parameters, e_i the model surfaces, A_s is the supernova extinction and C the extragalactic absorption law introduced by [Cardelli et al., 1989] (Cardelli extinction law). This method expands on the traditional SALT parameterization by adding as many extra surfaces as needed to fit the data. We can see that at second order the model becomes roughly the same as SALT2 with 2 parameters that represent the color and stretch of the supernova. Comparisons with the SALT2 color and stretch parameters have shown strong correlations. The [Saunders et al., 2018] paper shows that going up to the 7th order the supernovae variability decreases and at the 15th order the model fits the data best.




Fig. 2.16 Model reconstruction of SNEMO on SNfactory spectra ([Saunders et al., 2018]). On the left we see the fluxes of the spectra in comparison with 3 SNEMO models containing 2 (green), 7 (blue) and 15 (red) parameters. The right figures shows the weighted residuals of each model and spectrum.

5.2.2 SUGAR

The SUGAR (Supernova Generator And Reconstructor, [Léget et al., 2020]) model uses EMFA like SNEMO but applied in a different space with the idea that supernova variability is encoded in their spectral data. Therefore, instead of adding surfaces that would explore the entire space for the standardization features, the model has a pre-selected set of spectral features to look out for when determining the model. The founders of SUGAR have chosen 13 spectral features to identify the supernova variability, for which they show accounts for its entirety. The model for each supernova i is expressed as:

$$M_i(\lambda, p) = M_0(\lambda, p) + \sum_{j=1}^3 h_{i,j} \alpha_j(\lambda, p) + A_{V,i} f(\lambda, R_v) + \Delta M_{\text{gray},i} \quad (2.30)$$

with $M_0(\lambda, p)$ being the magnitude of the average supernova spectrum, $\alpha_j(\lambda, p)$ is the intrinsic variation related to the factor $h_{i,j}$, $A_{V,i}$ the extinction in the V band, $f(\lambda, R_v)$ is the Cardelli extinction law and $\Delta M_{\text{gray},i}$ is a normalization factor. The parameters $h_{i,1}$ and $h_{i,3}$ are correlated with the SALT stretch while the $A_{V,i}$ parameter is correlated with the color. The work with SUGAR has essentially showed that the supernova variability can be captured by 3 parameters, one more than the traditional stretch and color parameters.




Fig. 2.17 SUGAR model (blue) vs SALT2.4 (red) on top of the SNfactory data (black) taken from [Léget et al., 2020].

5.2.3 Twins Embedding

The Twins Embedding model ([Boone et al., 2021]) uses a non-linear parameterization to describe the supernova diversity. It uses a new standardization method based on the spectral differences between supernova at specific regions. If the spectral differences between two supernovae is small enough they are considered as "Twins" (Figure 2.18). This is also called the spectral distance, denoted as γ_{ij} . The idea is to firstly determine the average spectrum near t_{max} , called f_{mean} , and account for any redshift uncertainties and galactic extinction using the Cardelli extinction law. The variability is then modeled by a non-linear model and adjusted using an algorithm that does a non-linear dimensionality reduction called Isomap which is a generalization of PCA while accounting for non-linearities. The spectral distance between two Twins i and j is written as:

$$\gamma_{ij} = \left[\sum_k \frac{f_i(\lambda_k) - f_j(\lambda_k)}{f_{mean}(\lambda_k)} \right]^{\frac{1}{2}} \quad (2.31)$$

with f_i and f_j the spectra of i and j . The model is described using three parameters. The first parameter ξ_1 highlights the pseudo-equivalent widths (pEWs) of the Ca II features and it describes the most important variability. The second parameter ξ_2 affects the pEWs of the Si II lines and is correlated with the SALT2 stretch parameter X_1 . The third parameter ξ_3 affects the ejecta velocities of the supernova.




Fig. 2.18 Two examples of sets of supernova evenly spaced and considered as "Twins". Taken from [Boone et al., 2021].




Fig. 2.19 Variations of the Twins Embedding model near maximum light when varying each one of its components ξ_1 , ξ_2 and ξ_3 . For each component, the spectrum shown is the median of 20 evenly populated bins of the coordinates of that component. Taken from [Boone et al., 2021].

5.2.4 SuPAErnova

The Super Probabilistic Autoencoder model (SuPAErnova) ([Stein et al., 2022]) uses a probabilistic autoencoder in order to assess the supernova diversity. The use of machine learning (ML) in this case leaves the model free to explore different ways to determine the necessary parameterization in order to describe the data. It Takes as input the spectra from SNfactory and encodes the parameterization of the model. The general definition of the model is:

$$F_{SN}(\lambda, p) = \mathbf{g}([z_1, \dots, z_n], p) \times 10^{-0.4[CL(\lambda)\Delta A_V + \Delta M]} \quad (2.32)$$

where $\mathbf{g}$ is a decoder that learns to reconstruct the spectral series, z_i are the latent parameters of the model, CL is the color law, ΔA_V is the relative extrinsic extinction and ΔM the magnitude residuals. The results shown in the paper indicate that the with three latent parameters (z_1 , z_2 and z_3) the reconstruction errors on the spectral series are minimal. The only setback is that it then becomes difficult to have a physical interpretation of the final parameters.




Fig. 2.20 Model comparisons from SuPAErnova (red) and SALT2 (blue) on the spectral series of the SN2010dt supernova observed by SNfactory (black) as well as ratios between model and data. Taken from [Stein et al., 2022]

6 Main goal of this PhD

We present here the overall goal of this PhD: the measurement of $f\sigma_8$ at $z \approx 0$ using low redshift supernovae and setup for a combined analysis using galaxy clustering. Two primary avenues in order to conduct such analysis were explored. The first involved the improvement of how we measure supernova distances. As we have mentioned before the measurements of supernova distances play a roll in multiple

cosmological analyses (dark energy equation of state, H_0 and $f\sigma_8$), so one way we can improve the constraining power on the growth rate of structures is by improving our distance measurements. The second avenue was regarding the analysis method itself. Current measurements of $f\sigma_8$ using type Ia supernovae have been done using the Maximum likelihood method. This method falls short in reconstructing the non linearities in the velocity fields since it assumes a linear model. My goal will be to run a field level forward model inference to extract the information on $f\sigma_8$ while accounting for non linearities. This method will also serve as a benchmark for a cross survey analysis which can be done naturally in this formalism.

CHAPTER 3

LEMAITRE project

Contents

1	LEMAITRE: Hubble diagram project overview	72
2	Datasets: ZTF, SNLS, HSC	72
2.1	ZTF survey	72
2.2	SNLS survey	74
2.3	HSC survey	76
3	Pipeline	78
3.1	Pipeline overview	78
3.2	Cosmological analyses with LEMAITRE	78
3.3	Uncertainties in SN measurements?	78
4	DESI survey	79
4.1	Spectroscopic instrument	79
4.2	Tracers	79
4.3	ZTF and DESI Synergies	80

1 LEMAITRE: Hubble diagram project overview

The LEMAITRE project aims at constructing the largest Hubble diagram by using an independent set of supernovae. One its objectives is to tackle the $(w_0 - w_a)$ tension observed by recent results from [Brout et al., 2022], [Rubin et al., 2023] and [DES Collaboration et al., 2024] supernova analyses as well as the combination with BAO results from [DESI Collaboration et al., 2025]. It is important to note that the low redshift supernova sample used in all three analyses is the same and that it seems to be the main motivating factor for the tension. By analyzing a completely new dataset for LEMAITRE, an independent assessment of this tension will be provided. Additionally with the LEMAITRE project comes a new analysis pipeline from end-to-end, starting from the lightcurve extraction and calibration up to the cosmological analysis. My work during the PhD involved the continued development and testing of a portion of the LEMAITRE pipeline named NaCl (see Chapter 3 and 4) which aims at training a spectrophotometric model.

2 Datasets: ZTF, SNLS, HSC

There are three datasets used in the LEMAITRE project: ZTF, SNLS and HSC. The three supernova datasets cover a large redshift from $z_{\min} \sim 0.02$ to $z_{\max} \sim 1.5$. They amount to roughly 3000 type Ia supernovae that can be used for a cosmological analysis most of which come from the ZTF survey at low redshift. All three surveys also observe the same portion of the sky which will allow for an improved handling of calibration systematics. In this section we present the three datasets in greater detail.

2.1 ZTF survey

We start off with the Zwicky Transient Facility survey as well as the the supernova sample that was primarily studied in this PhD the ZTF DR2. The ZTF camera is mounted on the Samuel Oschin 48-inch (1.2m) telescope in the Mount Palomar Observatory, California, USA. The primary purpose of the survey is the study of nearby astrophysical transients, which makes it really good for studying supernovae in the local Universe.



Fig. 3.1 Mount Palomar Observatory. Credits: Caltech <https://www.ztf.caltech.edu/>

2.1.1 ZTF photometry

The ZTF camera is made up of 16 CCDs (model e2v CCD231-C6) each possessing 6144×6160 pixels. The Samuel Oschin telescope contains a 1.2m mirror (called the "primary mirror" M1) that redirects the photons to the focal plane of the camera. The mirror is a spherical pyrex-glass mirror coated with aluminium which allows for a large observation field of view. The telescope also contains an aspherical lens at the entrance to correct for geometric aberrations. It covers a solid angle of $\sim 47\text{deg}^2$ of the extra-galactical northern sky and manages to conduct a full sky coverage twice per night. The field of view can be seen in Figure 3.2 while the cadence is presented in Figure 3.3. Each image covers approximately 235 times the area of the full moon. The photometry is done using 3 filters g , r and i covering the ranges of $\approx 4000 - 5500\text{\AA}$, $\approx 5500 - 7000\text{\AA}$ and $\approx 7000 - 9000\text{\AA}$ respectively. It is important to highlight that the transmissions of the filters are sensor-dependent and we display them by averaging over the sensor transmissions (Figure 3.4). The magnitude limits for each band are $m_g \sim 20.8 - 21.1\text{mag}$, $m_r \sim 20.6 - 20.9\text{mag}$, $m_i \sim 19.9 - 20.2\text{mag}$ where the variations are largely due to the phases of the moon. The full details on the ZTF photometry are in [Dekany et al., 2020].




Fig. 3.2 Comparisons between the fields of view of ZTF, PS1, LSST, PTF, SDSS and DES. In the background we see in comparison the size of the moon and the Andromeda galaxy. Credits: Caltech <https://www.ztf.caltech.edu/>

2.1.2 Spectroscopic follow up

ZTF is accompanied by a spectrograph named the Spectral Energy Distribution Machine (SEDM) located in the same observatory on the P60 telescope. The purpose of the object to conduct a follow up on the photometry in order to properly classify the transients observed. Its limit magnitude is 20.5mag. This follow up played an important role in acquiring the 60% spectra for the ZTF Bright Transient Survey (BTS) for which the magnitude limit was 18.5mag, a limited fraction of the ZTF total survey.




Fig. 3.3 ZTF cadence on the sky. Average number of visits per month for each field of ZTF in each band g , r and i . Taken from [Carreres, 2023]

2.1.3 ZTF DR2

We present in this subsection the type Ia supernova data release of ZTF named DR2. This encompasses data taken from March 2018 and December 2020. Figure 3.5 shows the image a supernova observed by ZTF. The sample contains 3628 spectroscopically identified supernovae up to a redshift of $z \approx 0.2$. Figure 3.6 shows examples of the sampling of the supernova observed by ZTF in DR2. We must note that not all supernova data is "good" enough to be used for a cosmological analysis. For such an analysis, quality cuts on sampling around the maximum emission of light, the color and the stretch must be applied. The details on the cosmology quality cuts can be found in [Rigault et al., 2020] and we remind them later in Chapter 4. After the cuts, the sample maintains 2667 supernovae that pass. An example of what the ZTF DR2 Hubble diagram looks like can be seen in Figure 3.7. A subset of the sample named the "volume-limited" sample contains all supernovae up to a redshift of $z = 0.06$ represents a complete sample of supernovae i.e. the sample is not influenced by any selection effects due to the magnitude limits of the telescope. This subset contains 994 supernovae.

2.2 SNLS survey

The second dataset is the SuperNova Legacy Survey (SNLS) (Astier et al., 2006), a type Ia supernova survey conducted with the MegaCam (Boulade et al., 2003) observing system. The camera is located on the Canada-France-Hawaii 3.6m telescope (CFHT) on top of the Maunakea volcano, Hawaii. The focal plane of the MegaCam camera is made up of 40 CCDs (model e2v CCD42-90) each having 2048×4612 pixels. The CFHT is constructed with two mirrors, the primary mirror has a parabolic shape and is made from a glass-ceramic material which makes it immune to thermal distortion. The camera is placed in the focal plane of the primary mirror. In addition, a Wide Field Corrector (WFC) is placed in front of the camera to guarantee a large field of view.




Fig. 3.4 ZTF filters. Dotted lines represent transmissions for different sensors. Solid lines represent the average transmissions over all sensors.




Fig. 3.5 Type Ia supernova detected by ZTF. Photos taken accross 2 months (from left to right 20th of Jan 2020, 25th of Jan 2020, 7th of Feb 2020, 20th of Feb 2020, 20th of March 2020). Credits: Mikaël Rigault.

The spectroscopic follow-up used to classify the supernovae and identify their redshifts are ensured by the European Southern Observatory Very Large Telescope (ESOVL), the Gemini-North and South and the Keck-I and II instruments (Balland et al., 2009; Astier et al., 2006). The number of candidates for type Ia supernova observed photometrically exceed the number of events that can be followed up spectroscopically. This creates a need for a target selection process to try and eliminate transients that are not type Ia. The SNLS target selection is done by lightcurve fitting in real time to eliminate the majority of the type II supernovae. For the remaining candidates a limit magnitude is applied in the MegaCam *i* band at 24.5mag where supernovae are less likely to be spectroscopically observed.

The MegaCam footprint is a 0.90deg^2 rectangle, a much smaller footprint than ZTF that limits the survey from doing wide scans. Instead, deep observations are done in four fields equally distributed in right ascension up to a redshift of 1. (7ot figure). MegaCam contains 5 photometric filters used to observe transients: *g*, *r*, *i*, *i2* (The *i* filter was broken during SNLS observation period and was replaced a similar filter) and *z* (figure). This covers the visible and near-infrared wavelength ranges. As seen in




Fig. 3.6 Three supernova Ia lightcurves examples highlighting the worst (top), average (middle) and best (bottom) sampling in DR2. The colors indicate the band they were measured in and the solid lines are the best SALT2 fits. Taken from [Rigault et al., 2020]

Figure 3.4 for ZTF earlier, the sensor-dependent transmissions are also present in the MegaCam filters. The SNLS data used in the LEMAITRE project consists of five years of observation (SNLS 5y). The observations started in late 2003 and ended in 2009. The 4th and 5th years have not yet been published and will be in (Betoule et al. in prep.). The number is expected to be around 300 supernovae from SNLS that will be used in LEMAITRE.

2.3 HSC survey

The final dataset comes from the Hyper Suprime-Cam Subaru Strategic Program (Aihara et al., 2022), also denoted HSCSSP. It is an astronomical observation program which dedicated some telescope time to the observation of type Ia supernovae at the end of 2010. It uses the Hyper Suprime Cam (HSC), a camera mounted on the Subaru 8.2m telescope. The Subaru telescope is also located at CFHT on top of the Maunakea volcano, Hawaii.

The redshifts of the HSC supernovae are determined by measuring the redshift of their host galaxies. They're collected by the AAT/AAOmega and DESI instruments and completed by observations from the Subaru/FOCAS and VLT instruments (Suzuki et al. in prep). The HSCSSP project also determines redshifts directly from photometry using several complementary methods. The Direct Empirical Photo-




Fig. 3.7 Hubble diagram created with standardized ZTF DR2 type Ia supernovae compared to fiducial Λ CDM cosmology anchored on Planck results. Taken from [Rigault et al., 2020].

metric method (DEmP, [?]) and the TPZ method ([?]) both rely on machine learning to infer redshifts. DEmP does it by fitting a polynomial function relating redshift to magnitude, while TPZ is through random forest algorithms. Alternatively, the LePhare ([?, ?]) and MIZUKI ([?]) methods adopt a template fitting approach with priors, where LePhare uses redshift distribution priors from the 30-band photometric redshifts of the COSMOS survey, while MIZUKI uses Bayesian priors.

The Hyper Suprime-Cam camera footprint is a 1.8deg^2 disc. The main focus of HSCSSP is to build a galaxy catalog for weak lensing analysis, however the observation cadence of its deep fields were adapted for supernovae observation. The observation strategy of the HSC supernova survey is restricted to four deep fields (XMM-LSS, E-COSMOS, ELAIS-N1, DEEP2-F3) and two ultra-deep fields (SXDS and COSMOS) partially overlapping with the SNLS D1 and D2 fields, which go up to a redshift of $z \sim 1.6$.

HSC observation are done in 7 photometric filters: g , r , $r2$, i , $i2$, z and y spanning the visible and near-infrared wavelength ranges. The transmissions are sensor-dependent like the ZTF and SNLS filters. The data used in the LEMAITRE project consists of observations from a first session in 2017 up to observations from a second session in 2020. The number is expected to be around 300 supernovae from HSC that will be used in LEMAITRE.

3 Pipeline

3.1 Pipeline overview

3.2 Cosmological analyses with LEMAITRE

3.3 Uncertainties in SN measurements?

4 DESI survey

In this section, we briefly present the Dark Energy Spectroscopic Instrument (DESI). This novel, fully robotic instrument, has allowed us to observe the Universe in great detail and proven to being capable of adding insight to our understanding of the accelerated Universe ([DESI Collaboration et al., 2025]). In april 2026, DESI completed its main observation sequence and covered 14000deg^2 with over 60 million objects (with 47 million galaxies and quasars) observed within 5 years.

4.1 Spectroscopic instrument

DESI is a fourth-generation spectroscopic survey. It is a robotic fibre-fed spectroscopic instrument. It is installed on the 4-meter Mayall telescope at Kitt Peak National Observatory in Arizona, USA. It contains 5000 fibres, each able to observe a separate spectrum, allowing the instrument to gather 5000 spectra at a time, covering for one exposure around 3deg^2 part of the sky. This is a great improvement over its predecessor SDSS ([Dawson et al., 2016]) which had 1000 fibres positioned manually, therefore taking a lot more time than DESI. The DESI spectrograph covers a range of $3600 - 9800\text{\AA}$ where the light is divided into three wavelength channels: Blue (b), Red (r) and Near Infrared (Z). Finally, the light is captured by CCDs installed in a vacuum cryostat. Two different CCD models are used: STA4150 (4096×4096 pixels) and CCDs produced by LBL specifically for DESI (4114×4128 pixels). More details on the instrument can be found here [Collaboration et al., 2016a].

4.2 Tracers

DESI catalogues its observations under multiple classes each presenting a of matter tracer in the observable Universe ([Collaboration et al., 2016b]).

BGS

The first catalogue is the Bright Galaxy Survey (BGS). It is a flux limited galaxy catalogue that is divided into two parts based in the magntidue of the obsereved galaxy. BGS Bright contains galaxies with magnitudes in the r band < 19.5 and BGS faint with magnitudes $19.5 - 20.175$, these are present at a redshift range of $z < 0.6$. To identify if the obsereved object is a galaxy or a star it is cross references with the Gaia catalogue. If it is not present then it is a galaxy. There are over 10 million galaxies in the BGS catalogue. These also present the galaxies closest to us.

LRG

The second catalogue is the Luminous Red Galaxies (LRG) which contains old elliptical galaxies, identified due the $4000\text{\AA}$ break line in their spectra, in which the star formation rate has stopped. The LRG present galaxies in the redshift range $0.4 < z < 1.0$.

ELG

The third catalogue is the Emission Line Galaxies (ELG) which contains younger spiral galaxies with star formation. It is divided into two subcategories: ELG VLOP with galaxies at redshifts $0.6 < z < 1.1$ and ELG LOP with galaxies at $1.1 < z < 1.6$.

QSO

The fourth catalogue is the Quasar-Stellar Objects (QSO). The catalogue contains quasars, active galactic nuclei which are powered by the accretion of super massive black holes. The quasars measured at a redshift of $0.9 < z < 2.1$ are used as direct matter tracers, while quasars at $2.1 < z < 4.0$ will be used for their Lyman- α forest. This represents the furthest objects observed by DESI.

DESI at the time of writing this thesis has entered an extension phase of observation until 2028 where the sky coverage will go up to 17000deg^2 .

4.3 ZTF and DESI Synergies

ZTF and DESI present two of the current generation surveys that will tackle known tensions in the cosmological model. In addition to that, ZTF's sky coverage is in fact encompassed by DESI's coverage and the BGS sample is actually present at the same redshift range as ZTF. This creates synergies between the surveys that could be explored.

Firstly, DESI's spectroscopic instrument has more accurate measurements of redshifts than the SEDM, therefore redshifts provided by DESI to ZTF can improve the mapping of the expanding history of the Universe using supernovae. They will also provide more accurate measurements of growth rate of structures using ZTF since it heavily relies on comparing the observed redshift to the cosmological redshift due to the expansion (see Section 2.2).

Secondly, the BGS Bright sample contains galaxies in the same redshift range as the ZTF DR2. This highly favours conducting cross-survey analyses, which is the main goal of this PhD regarding the measurement of the growth rate of structures $f\sigma_8$. The DESI growth rate of structure measurements at low redshift is limited by the cosmic variance (the statistical uncertainty that arises due to only having access to one realization of the Universe) and galaxy bias. Supernova measurements can directly probe the underlying velocity field (see Section 2.2) independently of any galaxy bias. Therefore a combination of both DESI and ZTF is projected to yield more precise measurements of $f\sigma_8$ at a low redshift. This is particularly important since the measurement of the growth rate of structures is a direct test of GR and is most sensitive at low redshift.

CHAPTER

NaCl

Contents

1	NaCl framework	82
1.1	The idea behind the framework	82
1.2	Differences with the SALT models	82
1.3	Constraints	84
1.4	Regularization	85
1.5	Error models	86
1.6	NaCl output	87
2	Improving NaCl	90
2.1	Implementing a model for spectrophotometric data	90
2.2	Regularization	90
2.3	NaCl training recipe	92
3	NaCl on real data	94
3.1	NaCl training results	94

1 NaCl framework

The first part of my work for this thesis was the continuation of the development of a new framework designed to train spectrophotometric models of type Ia supernova called NaCl ("Nouveaux Algorithmes de Courbes de Lumière") which was initially started by G. Augarde during his PhD, defended in September of 2022 ([Augarde, 2022]).

1.1 The idea behind the framework

The framework is meant to be able to easily train a spectrophotometric model all while being able to accurately propagate all known sources of uncertainties. It is meant to be easy to use and manipulate in order to account for any new developments in how spectrophotometric models are parameterized. The idea is to then have a framework that is fast and capable of conducting model training easily every time one can do so on a cosmological dataset of type Ia supernova. We will see how this will allow us to overcome issues previous spectrophotometric models have encountered, such as calibration issues. The entirety of the framework has been coded in Python3.

1.2 Differences with the SALT models

This parameterization, like SALT2, is a classic principal component analysis, or PCA, with the addition of a color law. The main differences with SALT models is in the training procedure. NaCl training and lightcurve fitting is done simultaneously on the same dataset used for the cosmological analysis. This reduces the model systematics obtained from SALT models due to the change of datasets when conducting a cosmological analysis. It also means we can directly obtain a covariance matrix containing the correlations between supernova lightcurve parameters and also model parameters. This directly affects the distance modulus.

In NaCl the date of maximum emission t_{max} is fitted during the training of NaCl. This is a big difference from what is done for the SALT models where t_{max} is fitted beforehand. The reason why in NaCl we do that is to account for the possible changes of t_{max} that may arise due to the changes in the spectrophotometric model during the training.

Calibration uncertainties are modeled as nuisance parameters in NaCl that are fitted during the training, held by a covariance matrix allowing us to account for the correlations between filters. This also ensure we have an exact propagation of the calibration systematics in our training and prevents us from doing multiple trainings.

1.2.1 Practical implementation of the SALT2 model

In order to be able to determine the global model surfaces (M_0 and M_1), we will be describing them with the help of B-splines. This is a particular set of polynomials of order $m - 1$, called splines of degree m , that can form a basis. Their smoothness will allow our model surfaces to have complex forms. We use splines of degree 4 in NaCl. M_0 and M_1 are then defined as:

$$M_{0|1} = \sum_{k=1}^{n_p} \sum_{l=1}^{n_\lambda} \theta_{kl}^{[0|1]} \mathcal{B}_k(p) \mathcal{B}_l(\lambda) \quad (4.1)$$

where $\mathcal{B}$ are the splines, θ are the model parameters that we will determine. Similarly, the evaluation of the filters is also done on the same spline basis of the model. For the color law $CL(\lambda)$ it is implemented as a degree 5 polynomial.

Since we do not determine the magnitude of the supernova on the spectra, we add an extra polynomial that for each spectra in our dataset. The polynomial $SpecRecal(\lambda)$ can be then added to the spectroscopic element of the model such that:

$$\phi_{spec}(\lambda, p) = \frac{SpecRecal(\lambda)}{1+z} \mathcal{S}\left(\frac{\lambda}{1+z}, \frac{p}{1+z}\right) \quad (4.2)$$

This entails that there will be a degeneracy between the color law CL and the recalibration polynomials, therefore during the fitting of the spectra we do not determine CL (essentially it is not treated as a parameter that can influence the overall shape of the spectral features of the model).

The model parameters are then determined by minimizing a log-likelihood which takes the form of:

$$-2\ln(\mathcal{L}) = N\ln(2\pi) + \ln(|\mathbf{V}|) + \mathbf{R}^T \mathbf{V}^{-1} \mathbf{R} \quad (4.3)$$

where $\mathbf{R}$ are the model residuals and $\mathbf{V}$ the covariance matrix independent of the parameters. If $\mathbf{V}$ contains only measurement uncertainties, the problem then becomes minimizing the following quantity:

$$\chi^2 = \mathbf{R}^T \mathbf{V}^{-1} \mathbf{R} \quad (4.4)$$

which will be the case for us during this PhD.

The minimization of the log-likelihood is via the Fisher scoring algorithm. This method allows us to obtain the desired model parameters that best describe the data using a second order descent of χ^2 (slope and curvature information). If we denote $\mathbf{M}$ the model, β the model parameters, $\mathbf{J}$ the model Jacobian and $\mathbf{H}_\chi$ the Hessian (second order derivatives matrix) of the χ^2 then it is possible to show using the Fisher scoring algorithm that for a set of parameters β_i at an instance i to be updated at an instance $i+1$ with $\beta_{i+1} = \beta_i + \delta\beta$ we have:

$$\delta\beta = -(\mathbb{E}[\mathbf{H}_\chi])^{-1}(\beta_i) \nabla \chi^2(\beta_i) \quad (4.5)$$

where $\mathbb{E}$ is the expected value. It is possible to relate the χ^2 Hessian with the model Hessian in order to simplify this expression. Will be omitting this demonstration but we get $\mathbb{E}[\mathbf{H}_\chi] = 2\mathbf{J}^T \mathbf{V}^{-1} \mathbf{J}$ for an unbiased model and this relation is true at the minimum value of the log-likelihood. Additionally, the gradient of χ^2 can be written as $\nabla \chi^2 = -2\mathbf{J}^T \mathbf{V}^{-1} \mathbf{R}$ where $\mathbf{R}$ is the model residuals vector. Finally, we get the following expression for $\delta\beta$:

$$\delta\beta = \left(\mathbf{J}^T \mathbf{V}^{-1} \mathbf{J}\right)^{-1} \mathbf{J}^T \mathbf{V}^{-1} \mathbf{R} \quad (4.6)$$

This algorithm requires us to invert the Hessian matrix $\mathbf{H} = 2\mathbf{J}^T \mathbf{V}^{-1} \mathbf{J}$ at each step, which is always positive definite by construction, allowing us to use efficient factorisation methods such as the Cholesky decomposition. In practice, minor degeneracies are handled by augmenting the diagonal of the Hessian with a small damping term following the Levenberg-Marquardt approach, and by enforcing parameter constraints through Lagrange multipliers, both of which improve the numerical stability of the inversion.

1.3 Constraints

Since this model is built empirically, we can see that there will be a lot of degeneracies between the individual and model parameters. To break those degeneracies we will be adding a set of constraints.

When NaCl was first developed, the constraints used were the same as the ones for SALT2 and SALT3 ([Kenworthy et al., 2021]) models. Although those constraints break the degeneracies that exist between parameters, we've decided to include a new set of constraints that would address two things : the first that one of the constraints is non linear which made the model training take a longer time, the other was that some constraints depended on the supernovae used in the training sample. An example such a constraint was $\langle X_1 \rangle = 0$ which defined the average stretch of a supernovae. This is isn't an issue if we are comparing the stretch of supernovae that were fitted using the same model. However since NaCl will be used to train different models on different datasets, it'll be more difficult to directly compare the parameters of the supernovae.

During this PhD, the following sets of constraints were developed. I will be detailing what they do and why they were developed.

Model normalization:

The model normalization is degenerated with the value of X_0 . We impose that :

$$\int M_0(\lambda, p=0) \frac{\lambda}{hc} B(\lambda) d\lambda = 1 \quad (4.7)$$

$$\int M_1(\lambda, p=0) \frac{\lambda}{hc} B(\lambda) d\lambda = 0 \quad (4.8)$$

where $B(\lambda)$ is the restframe standard B -band.

Date of maximum emission:

There exists a degeneracy between the model and the date of maximum emission t_{max} . It's possible to shift the entire model in order to compensate for a change in the date t_{max} . For that we impose that at $p=0$, the model presents a maxima in the standard B -band :

$$\frac{d}{dp} \int M_0(\lambda, p=0) \frac{\lambda}{hc} B(\lambda) d\lambda \Big|_{p=0} = 0 \quad (4.9)$$

$$\frac{d}{dp} \int M_1(\lambda, p=0) \frac{\lambda}{hc} B(\lambda) d\lambda \Big|_{p=0} = 0 \quad (4.10)$$

Color Law:

We also need to account for the degeneracy that exists between the spectral shape of the model and CL . Tilting the M_0 and M_1 can be completely corrected for by changing CL . We impose then that $CL(B) = 0$ and $CL(V) = 1$ with B and V being the central wavelengths of the restframe standard B -band and V -band.

Average color:

Another degeneracy exists between the individual supernova color c and the M_0 . Changing the spectral features of the model can be completely compensated by the value of c . For that we need to define the "average" supernova color. We impose that the M_0 surface would have its $B - V$ color equal to 0 at peak luminosity and that c accounts for the relative variations in color with respect to that. The constraint now becomes :

$$\int [M_0(\lambda, p=0)B(\lambda) - M_0(\lambda, p=0)V(\lambda)] \frac{\lambda}{hc} d\lambda = 0 \quad (4.11)$$

Average stretch:

A similar degeneracy also exists with X_1 . A global change of the X_1 's can be absorbed by changing the shape of the M_0 surface along the phase axis. We have decided to fix the value of $X_1 = 1$ such that it would be equivalent to having the same X_1 value using SALT2. For that we fix the difference between the model evaluated for $X_1 = 0$ and $X_1 = 1$ at a phase of 15 :

$$\int [M_0(\lambda, p=15) + M_1(\lambda, p=15)] \frac{\lambda}{hc} B(\lambda) d\lambda - \alpha \int M_0(\lambda, p=15) \frac{\lambda}{hc} B(\lambda) d\lambda = 0 \quad (4.12)$$

with $\alpha = 10^{-0.4 \times 0.162}$ the flux difference we have found in SALT2.

Average width:

The last unwanted degree of freedom is regarding the width of the model. We can see that the definition of X_1 relies on a difference between the model with different stretch parameters but it doesn't define what the width of the model is. We can always make the model larger or smaller while maintaining the constraints on all of the other parameters. One way to bypass that issue is to constrain the model at a certain phase. We choose to fix the model at a phase $p = 15$, also known as Δm_{15} as :

$$\int M_0(\lambda, p=15) \frac{\lambda}{hc} d\lambda = 10^{-0.4} \quad (4.13)$$

In practice, we write down the constraints as a function $C_k(\beta) = \alpha_k$ where $C_k(\beta)$ is a function evaluating the k -th constraint of value α_k . This is then added as penalty that takes the form $\mu_{cons} \sum_k (C_k(\beta) - \alpha_k)^2$ where μ_{cons} is hyperparameter that determines the strength of the penalty. We can write this in a matrix $\mathbf{C}(\beta)$ and we obtain :

$$\chi_{cons}^2 = \mu_{cons} \mathbf{C}(\beta)^T \mathbf{C}(\beta) \quad (4.14)$$

1.4 Regularization

Because our training dataset contains a finite number of data points, it's impossible that the entirety of the global parameters would be constrained by data. To deal with that we add regularization to the model during the fit, which is a penalty added to the model in the areas where we would have extremely large variations due to poor constraining power of the data. This is specifically really important for NaCl since we are fitting the date of maximum emission there are data points that could go outside of the

model range.

The regularization penalty that was initially developed was expressed as :

$$\mu_{reg} \sum_{kl} \left(\theta_{k,l}^2 + (\theta_{k+1,l} - \theta_{k,l})^2 \right) \quad (4.15)$$

where θ_{kl} are the model parameters as seen in equation 4.1. The first sum is a penalty that makes the θ 's tend to zero when no data is available. The second term is a penalty of the phase derivatives of the surfaces that limits the sharp variations of the model observed when no data is available, or at poorly constrained regions of the model. Finally, μ_{reg} is a hyperparameter that determines the strength of the penalty. Writing this in a matrix form gives us :

$$\chi_{reg}^2 = \mu_{reg} \beta^T \mathbf{P} \beta \quad (4.16)$$

where $\mathbf{P}$ is what we call the regularization matrix, that allows us to obtain the expression in equation 4.15, and β our set of parameters.

1.5 Error models

Dealing with an empirical model means that we expect that our model would not perfectly describe the data. More specifically, the residuals of the model would not be explained solely by the measurement uncertainties on the data. This is a variability that exists between supernovae, unable to be captured by the SALT2 parameterization.

It's also important to note that all of the photometric measurements are affected by uncertainties of the photometric instruments. This can play a crucial role in our measurements of cosmological parameters as it has been shown in the most recent analyses to be one of the leading systematics ([Rubin et al., 2024], [Vincenzi et al., 2024]).

We can account for both the supernova variability and calibration uncertainties with the help of error models.

1.5.1 Error Snake

The first error model is the Error Snake, which is supposed to capture errors directly due to variability in the flux. If we call V the variance of the data points, then the error model is implemented as :

$$V = \sigma_{meas}^2 + \sigma_{snake}^2 \quad (4.17)$$

where σ_{meas} are the measurement uncertainties and σ_{snake} is the additional variance added that is our Error snake model.

The general expression of this term is :

$$\sigma_{snake} = G(\lambda, p) \times \phi(\lambda, p) \quad (4.18)$$

where ϕ is the concatenation of both the measured photometric and spectroscopic fluxes used in our training dataset. $G(\lambda, p)$ can take multiple forms depending on how we would like to model our Error snake.

The simplest way to model it would be as a constant. The expression becomes :

$$G(\lambda, p) = g \quad (4.19)$$

Physically this means that the variability of the supernovae in our dataset are independent of phase and wavelength.

Alternatively, we can imagine something more complex by modeling G as a surface like M_0 and M_1 that is to be determined during the training. The expression then becomes:

$$G(\lambda, p) = g(\lambda, p) \quad (4.20)$$

Here, the variability does depend on phase and wavelength.

Lastly, we thought of adding a bit more complexity in our Error snake since we know that it's possible that our dataset could be contaminated by very peculiar supernovae or by objects that are hard to distinguish from type Ia supernova such as type Ib and type Ic supernovae or . We decided to also include an extra term in G that takes up one value for each supernova in our dataset, we call it γ_{SN} . We expect that the γ_{SN} value of a peculiar supernova would be larger than the others. The expression then becomes :

$$G(\lambda, p) = \gamma_{SN} \times g(\lambda, p) \quad (4.21)$$

which is what we would like to use in the final analysis of the LEMAÎTRE dataset. However it is important to note that for the data challenges, specifically DC1 (see Chapter 4), we will be using the simplest form of the Error snake.

1.5.2 Color Scatter

COMMENT : Waiting for the final implementation

1.5.3 Calibration Uncertainties

The calibration uncertainties are taken into account directly in the model. One parameter η is added for each transmission band in our dataset held by a prior that is known beforehand. For the LEMAÎTRE project we have 12 bands, so we'll have 12 η parameters. The expression of the photometric model becomes :

$$\phi_{phot}(\lambda, p) = X_0(1 + \eta)(1 + z) \int S(\lambda, p) T(\lambda) \frac{\lambda}{hc} d\lambda \quad (4.22)$$

With an additional penalty to the model :

$$\chi_{calib}^2 = \eta^T \mathbf{V}_\eta^{-1} \eta \quad (4.23)$$

$\mathbf{V}_\eta$ being the η covariance matrix that is already known.

1.6 NaCl output

At the end of a training, we get a vector containing all of the model and individual supernova parameters and the full covariance matrix calculated from the Hessian of the model : $\mathbf{V} = 2\mathbf{H}^{-1}$. For the cosmological analysis, we will be marginalizing over all but the X_0 , X_1 and c parameters.




Fig. 4.1 Visual representation of the error snake on simulated lightcurves from DC1 (see section (reference)). In 4.1a we see how the model represents the data without an error snake model. In 4.1b we see a highlighted section for each lightcurves which are the variations modeled by the error snake.




Fig. 4.2 Example of the NaCl covariance matrix after an output, marginalized over all but the X_0 , X_1 and c parameters.

2 Improving NaCl

The foundational framework of NaCl was established by Guy Augarde ([Augarde, 2022]), whose work defined its structure and validated it on LSST-like simulations. However, several improvements were identified as necessary to handle realistic data. As part of my PhD, I contributed to developing these improvements, which are presented in the following sections.

2.1 Implementing a model for spectrophotometric data

Training a spectrophotometric model is done on two types of dataset. Classical models used in cosmological analyses like SALT2 or SALT3 rely on photometric and spectroscopic data obtained separately. Models that aim at refining the standardization scheme of supernovae rely on spectrophotometric data like the SNfactory data ([Aldering et al., 2002]). The SNIFS instrument used for SNfactory allows for us to obtain a series of well calibrated spectral sequences for each observed supernova making the need for separate photometric data obsolete. Being able to use such datasets in NaCl will then require us to add supplementary information in the model to account for that.

If we recall in Chapter 2, the SALT parameterization is expressed as:

$$\mathcal{S}(\lambda, p) = [M_0(\lambda, p) + X_1 M_1(\lambda, p)] 10^{0.4cCL(\lambda)} \quad (4.24)$$

used to describe the following spectroscopic and photometric fluxes:

$$\phi_{spec}(\lambda, p) = \frac{1}{1+z} \mathcal{S}\left(\frac{\lambda}{1+z}, \frac{p}{1+z}\right) \quad (4.25)$$

$$\phi_{phot}(\lambda, p) = X_0(1+z) \int \mathcal{S}\left(\lambda, \frac{p}{1+z}\right) \frac{\lambda}{hc} T((1+z)\lambda) d\lambda \quad (4.26)$$

For spectrophotometric data, the goal is to then describe one data-type that follows:

$$\phi_{specphot}(\lambda, p) = X_0 \frac{1}{1+z} \mathcal{S}\left(\frac{\lambda}{1+z}, \frac{p}{1+z}\right) \quad (4.27)$$

thus removing the need for a recalibration polynomial *SpecRecal* and reintroducing the color law *CL* which will be determined on the spectra. My work was to code this model in NaCl which led to the first use of NaCl on a real dataset where the results are presented in Section 3.


2.2 Regularization

An issue that we've had with the previously mentioned form of regularization is that it is very sensitive to the value of μ_{reg} . If it's too small the model continues to present very strong variations. If it's too strong the model becomes attenuated and this biases our parameters, which means the cosmological parameters calculated with the output of NaCl would be biased. We show the effects of the strength of the regularization on the model in Figure 4.3

To try and overcome this issue, I've developed during this PhD a regularization scheme that adapts the force of the regularization based on the number of points seen by each model parameter. This regularization scheme is inspired by [Lenz et al., 2022]. We can separate $\mathbf{P}$ matrix as follows:


$$\mathbf{P} = \mathbf{P}^{data} + \mathbf{P}^{smooth} \quad (4.28)$$

$$\mu_{reg} = 10^{-6}$$



(a)

$$\mu_{reg} = 1$$



(b)

Fig. 4.3 Effects of the strength of the regularization on the shape of the M_0 . In 4.3a we see how the model behaves with a $\mu_{reg} = 10^{-6}$. In 4.3b we see how the model behaves with a $\mu_{reg} = 1$.

with $\mathbf{P}^{data}$ and $\mathbf{P}^{smooth}$ the matrices determining the strength based on the number of points and to smoothen the model where there is no data as much as possible. In order to determine $\mathbf{P}^{data}$, we define $\mathcal{N}_i$ as the number of data points seen by each θ_i parameters (θ_{kl} written as a vector). We then introduce for each $\mathcal{N}_i$ a parameter μ_i which will act as a gate function of $\mathcal{N}_i$. This will determine the force of our regularization. The idea is that if $\mathcal{N}_i = 0$ we regularize model strongly, otherwise we don't.

COMMENT : This specific version of the regularization wasn't implemented in the final version of NaCl since we still had to test other things in between. I will be rewriting this part of the manuscript again.

μ_i is then defined as:

$$\mu_i^{data} = \max(\mu_{reg}^{data} - \mathcal{N}_i, 0) \quad (4.29)$$

We can use that to define a diagonal matrix $\mathbf{M}^{data}$ where $\mathbf{M}_{ii}^{data} = \mu_i^{data}$. It is then clear that the simplest way to define $\mathbf{P}^{data}$ will be to write it as:

$$\mathbf{P}^{data} = \mathbf{M}^{data} \times \mathbb{I} \quad (4.30)$$

We then define $\mathbf{P}^{smooth}$ as a penalty on the first order derivative of the model in phase, like in equation 4.15. We apply the same adaptive method mentioned previously and we get:

$$\mathbf{P}^{smooth} = \mathbf{M}^{smooth} \times \mathbb{P} \quad (4.31)$$

with $\mathbb{P}$ being :

$$\mathbb{P} = \begin{bmatrix} 2 & -1 & 0 & & & \dots \\ -1 & 1 & -1 & 0 & & \dots \\ 0 & & & \dots & & \\ & & \dots & & & 0 \\ & \dots & 0 & -1 & 1 & -1 \\ \dots & & & 0 & -1 & 2 \end{bmatrix} \quad (4.32)$$

with μ_{reg}^{smooth} would be the hyperparameter determining the strength of the regularization on the derivatives.

Since the hyperparameters are now included in the $\mathbf{P}$ matrix, the penalty now becomes :

$$\chi_{reg}^2 = \beta^T \mathbf{P} \beta \quad (4.33)$$

2.3 NaCl training recipe

During the tests of NaCl on simulations of LEMAITRE-like supernova, we have noticed that fitting all of the parameters can cause issues. If the model parameters are not close to the truth we found cases where the error model would dominate. This is not problematic for the minimizer as it sees this as a way to describe the data but physically none of the parameters are being fitted. Another issue encountered was the possibility of converging the minimizer at a local minima rather than a global minima. We've developed during the PhD a training recipe that allows for having good initial guesses of the supernova and model parameters. This helps accelerate the fit and makes it more stable. The current recipe is the following:

1. Lightcruve fit with the initial NaCl model. In this part we are only fitting X_0 , X_1 , c and t_{max} for all supernova lightcurves.

2. SpecRecal fit with the initial NaCl model. Here we are estimating the recalibration polynomial coefficients. This part of the fit is near instantaneous due to the linearity of the polynomial.
3. Initial model guess. In this part we aim to obtain relatively close initial guesses for M_0 , M_1 and CL to the truth. This also ensures that the model respects the constraints.
4. Initial error model guess. Here we have an initial guess for the error model for a fixed model.
5. Full training. All parameters are fitted.

The first two steps of the recipe do not take a lot of time to complete. The Initial model guess step can take some time to converge due to large number of parameters of the M_0 and M_1 surface, however the number of steps is limited to 15 in that step since the goal is to simply get an initial guess. The error model fit is also near instantaneous. Finally, freeing all of the parameters in step 5 takes the most amount of time.

3 NaCl on real data

There have been models that were empirically constructed in similar ways as the SALT models with SNfactory data, namely SNEMO ([Saunders et al., 2018]), SUGAR ([Léget et al., 2020]), Twins Embedding ([Boone et al., 2021]) and SuPAErnova ([Stein et al., 2022]) as presented earlier. These models usually incorporate different parameterizations than SALT2 in order to try and fully understand the spectrophotometric evolution of supernovae. One thing to note is that it is easy to implement different models in NaCl. So one could modify the current implementation of the SALT2 model and add extra parameters and test if it improves the description of supernovae.

For now, we present the results obtained when training NaCl on the SNfactory data with a SALT2 parameterization.

3.1 NaCl training results

3.1.1 Data coverage

Training a SALT2-like model with NaCl on the SNfactory was the first time NaCl was used on real data and was part of the PhD to test the framework. The SNfactory dataset used here contains 224 supernova and 3135 spectra. We can see in the coverage of the data on the model's $\lambda - p$ space in Figure 4.4 that we have a very rich dataset that covers a very large part of the model's domain of definition.




Fig. 4.4 Coverage of the SNfactory data in the $\lambda - p$ space.

3.1.2 Fitting the spectra


We present in Figure 4.5 examples of spectra that were fitted with NaCl. These are 2 spectra randomly picked out that belong to two different supernova and are taken at different phases. What we see is that the trained model seems to accurately describe the evolution of those spectra. The errorsnake model employed shows an intrinsic dispersion of the fluxes at an order of about 10%. We present in the next subsection the results of full dataset.

The entirety of the model training takes around 30 minutes on a Dell Precision 3581 laptop equipped with an Intel(R) Core(TM) i7-13700H, 14 cores, up to 5.0 GHz, 32 GB RAM and a NVIDIA RTX A1000 6GB GPU.


3.1.3 Training residuals

We would like to examine how well a SALT2 like parameterized model is able to describe the spectrophotometric evolution of supernovae. In order to do that, we will be looking at the weighted residuals of the model on the entirety of the data and check if there are any particular features that stand out. This is presented in Figure 4.6 where we have calculated the weighted residuals which represent by how many σ our model differs from the data, also known as pulls. We do that for each data point and stack them on top of each other if they have been observed at the same time. So for example if we have 2 spectra measured at phase of $p = -1$ with data points at $5000\text{\AA}$, it'll appear as one point on the Figure and its value will be the average of the 2 pulls. We then get a Figure that represents the accuracy of the model in the $\lambda - p$ space. This also means that each horizontal line represents on average how well a spectrum is modeled at that particular restframe phases, while the vertical lines are for specific restframe wavelengths.

We see in the Figure 4.6 that for most of the $\lambda - p$ space the residuals show good agreement between the model and the data. But what also stands out is that we can identify three distinct wavelength regions where the model is unable to capture the variability of the data. These three regions are found at $\sim 3500\text{\AA}$, $\sim 6000\text{\AA}$ and $\sim 8000\text{\AA}$, which correspond to UV, Si II and Ca II. These are known to have very diverse forms ([Wang et al., 2012], [Parrent et al., 2016]) and we can see here that they are not captured by a SALT2 parameterization. It is important to mention that the models mentioned in Chapter 2 that go beyond the SALT2 parameterization can capture those variabilities with the help of extra parameters.



(a)



(b)

Fig. 4.5 Two spectra with the train NaCl model. The blue points are the data, the red line is the trained model and the highlighted area is the error model.




Fig. 4.6 Weighted residuals at the end of the training done with NaCl on the SNfactory data. The color bar represents by how many σ our model differs from the data. We also use as examples 3 spectra at 3 different phases and show where they would be present on the plot.

CHAPTER 5

LEMAITRE Data Challenges

Contents

1	LEMAÎTRE Data Challenges	100
1.1	General description of the DCs	100
1.2	Georges analysis pipeline	100
1.3	DC1	103
1.4	DC2	103
1.5	DC3	103
1.6	DC4	103
1.7	DC5	104
2	pre-DC1	105
2.1	Data generation : simulation	105
2.2	Goals	106
3	Results : Potential biases & systematics	107
3.1	Model & data agreement	107
3.2	Parameter reconstruction and covariance matrix	107
3.3	Distance modulus and cosmological parameters reconstruction	108

1 LEMAÎTRE Data Challenges

In this section we will be detailing the LEMAÎTRE Data Challenges, DCs from now on. The goal of these challenges is to validate the new analysis pipeline that is part of the LEMAÎTRE project, which is an extremely important key test before running the full analysis chain on the real data.

1.1 General description of the DCs

The LEMAÎTRE analysis pipeline is brand new from the point where the lightcurves are extracted from the ZTF, SNLS and HSC surveys up until cosmological parameters are determined. The pipeline can be divided into 2 main parts : the lightcurve extraction and calibration pipeline, and the Georges pipeline (or the cosmological analysis pipeline). Determining whether this new pipeline can accurately predict cosmological parameters without any biases is crucial and must be investigated thoroughly, especially since many of the elements of the pipeline depend on each other.

We will be focused here on the validity of the cosmological analysis pipeline, Georges. For that, the pipeline will face a series of challenges, each with an increasing level of complexity representing a challenge we are expected to face in the final analysis on the LEMAÎTRE dataset. Those challenges are the DCs and they are constituted of realistic simulations of ZTF, SNLS and HSC lightcurves and spectra. We will firstly describe the different elements of the Georges pipeline and then the different elements of the DCs. Each DC is constituted of 100 realizations of the simulations, essentially containing the exact same supernova but by applying the noise on the data differently. This will allow us to perform a robust coverage test for each DC.

1.2 Georges analysis pipeline

The Georges pipeline consists of mainly four modules : PeTS, NaCl, EDRIS, cosmologix.

1.2.1 PeTS & miniPeTS

PeTS

The PeTS module aims at "cleaning" the training dataset we will ll be using in the cosmological analysis. The reason we need to do that is because some of the supernova lightcurves can contain issues such as few points, very noisy data or a mix of both that can introduce issues when trying to accurately model the supernova with NaCl.

In order to avoid that, PeTS applies a set of data quality cuts that will ensure that we select a subset on which we will be able to properly measure all of the supernova parameters.

The selection will be applied on supernovae that meets the following criteria:

- The date of maximum emission (t_{max}) needs to be properly defined
- The color and the stretch parameters need to be accurately measured
- The supernova must contain at least 5 points with an SNR superior to 3, present in 2 bands and present within 50 days of the initial guess of t_{max} .

In order to do that, PeTS initially fits all of the lightcurves with the SALT2.4 model using `sncosmo` ([Barbary et al., 2021]). **The `sncosmo` fit needs to converge in order to select a supernova in our final dataset.** That initial fit allows us to have an initial guess on the parameters c , X_1 and t_{max} .

The next thing is to create a log-likelihood profile while varying the value of t_{max} . The supernova is then selected if the following criteria are met on the parameter t_{max} :

- The uncertainty on t_{max} derived from the log-likelihood profile must be inferior to 1 day
- The log-likelihood profile must be symmetric. The condition chosen for that in PeTS is :

$$|3\sigma(t_{max})_{Left} - 3\sigma(t_{max})_{Right}| < 0.3$$

- The log likelihood profile must present only 1 minimum within 3σ

Lastly, the uncertainties on c and X_1 need to be inferior to 0.3 and 1.

If all of these conditions are met, the supernova will be present in the final dataset. For the DCs, we want to apply PeTS on only 1 realization since the quality cuts shouldn't vary between realizations. Running PeTS, however, presents a difficulty. Creating the log-likelihood profile for each supernova takes a lot of computational time. It would take us a really long to do so if we wanted to perform multiple tests on any DC. In order to avoid that and gain a bit of time, we introduced miniPeTS.

miniPeTS

With miniPeTS the goal is to achieve similar quality cuts to PeTS but much faster. This means that we don't expect to necessarily obtain the best quality dataset, but good enough to test the pipeline on the DCs. We apply similar cuts to the ZTF DR2 cuts (ref). We can summarize the quality cuts done with miniPeTS in the following list:

- The supernova needs to be measured in at least 2 bands
- The supernova needs to have at least 5 total measurements
- The supernova needs to have at least 2 measurement points before t_{max}
- The supernova needs to have at least 2 measurement points after t_{max}
- $-0.2 < c < 0.8$
- $|X_1| < 3$
- SNLS and HSC supernova need to be measured at $z > 0.2$
- $E(B - V)_{MW} < 0.25$

1.2.2 Data compression

Before we can start training a model with NaCl on the data, we decided to accelerate the training by compressing the data on the basis of the model.

Lightcurve compression

The way we compress the lightcurves data is by averaging the fluxes measured per night. We find that doing so reduces the size of lightcurves files by almost half.

Spectra compression

We have also noticed that the resolution of the spectra is much higher than the resolution of our model. The spatial dimension of our model in NaCl is defined by a B-spline basis of order 4. Projecting the spectra on that basis can be written as

$$\sum_i p_i \mathcal{B}_i(\lambda)$$

The idea is retain the p_i parameters alongside their uncertainties. The size of the spectral data is then reduced by a factor of 10.

1.2.3 NaCl

The next step in the Georges analysis pipeline is the model training with NaCl. The X_0 , X_1 and c along with the marginalized covariance matrix will be the input for EDRIS in order to estimate the supernova distances.

1.2.4 EDRIS

The observed distance modulus of type Ia supernova contains a natural dispersion with respect the Hubble law of around $\sim 0.4\text{mag}$. This can be reduced by standardizing the distance modulus. We commonly use the Tripp relation (ref) which reduces this dispersion to $\sim 0.15\text{mag}$. It is expressed as follows :

$$\mu = m_B - (M_B - \alpha X_1 + \beta c) \quad (5.1)$$

$$m_B = -2.5 \log_{10}(X_0) \quad (5.2)$$

with m_B being the observed magnitude, M_B the absolute magnitude and (α, β) are the standardization parameters also known as the brighter slower and brighter bluer parameters. This is due to the fact that type Ia supernova with a larger stretch parameter tend to be brighter as well as supernova that are bluer.

The remaining 0.15mag dispersion are due an intrinsic variability between supernovae that isn't modeled by the Tripp relation.

Determining the standardized parameters can be difficult. Since telescopes can only observe objects up to a certain magnitude we expect a cutoff in magnitude for each survey. Coupled with the intrinsic dispersion of supernova, this means that the population of supernova observed by a survey will be skewed, favoring redder and faster supernova. We call this **selection effects**. This makes a direct determination of (α, β) biased and will bias any cosmological analysis.

The goal of EDRIS is to accurately estimate unbiased standardized distances for complete and incomplete supernova Ia samples. EDRIS takes as input the parameters from NaCl alongside their covariance matrix, and will use that to infer standardization parameters (α, β) , M_B , selection effects and the intrinsic dispersion denoted σ_{int} .

1.2.5 Cosmologix

The final step of the pipeline is to infer the cosmological parameters from the distances and we do that with cosmologix ¹ (Betoule et al in review), a fast JAX-coded software designed to accurately infer cos-

¹<https://lemaitre.pages.in2p3.fr/cosmologix/home.html>

mological parameters. It is also possible to combine the supernova results with other probes such as BAO and CMB. This will be useful in the analysis of the real data.

1.3 DC1

The first data challenge will be a consistency test for the full chain. It's the first time we've tested the full pipeline together on the same dataset. This will also serve as a stepping stone for the analysis of the future DCs since they will be incrementing layers of complexity on top of what was done in DC1. We will be including simulated lightcurves and spectra from ZTF and SNLS as well as lightcurves from HSC.

The simulated data will include measurement uncertainties, an error snake and an intrinsic dispersion. We will not be including any selection effects so EDRIS will need to find the standardization parameters. The main challenge will be for NaCl to ensure that training the model with our current model constraints and regularization on a LEMAITRE like dataset does not bias our cosmological analysis.

The metric used to determine if we have passed the DC or not will be to check the coverage for the cosmological parameters after 100 realizations and ensuring it stays within 1σ .

1.4 DC2

The second data challenge serves as a test to the model selection effects in EDRIS. The simulations will be identical to DC1 with the addition of a selection function for each survey.

The metric used to determine if we have passed the DC or not will be the same as DC1.

1.5 DC3

In the third data challenge, calibration uncertainties and supernova outliers will be added the simulation. This serves as a main test to NaCl's error model to see how well it can identify outliers as well as determining the impact of calibration uncertainties on the reconstructed cosmology.

The metric used to determine if we have passed the DC or not will be the same as DC1 and DC2.

1.6 DC4


In the fourth data challenge, the simulations will include as many realistic effects as possible adding all sources of uncertainties and systematics that we know can affect our analysis. This includes realistic distributions of the stretch and color (ref madeleine), non type Ia contamination, astrophysical effects and color scatter effects. This will be a challenge for both NaCl and EDRIS.

The metric we will use here to determine if we have passed the DC or not will be the same as before. Since astrophysical effects aren't directly taken into account during the modeling of type Ia supernova, we will quantifying its effects on the reconstructed cosmology and see how we can minimize it.

1.7 DC5

In the fifth and final DC we will be placing ourselves in the same conditions of the realistic simulations by blinding the data. The party that will be blinding the data will not be the same as the one running the pipeline simulation. The goal is to obtain the correct cosmological parameters after unblinding the data.

We can summarize the DCs in the following diagram :



2 pre-DC1

Part of my PhD work, I took charge of the analysis of pre-DC1, a slightly different version of the DC1 where we use a SALT model that respects the NaCl constraints to generate the supernova lightcurves. This allows us to remain coherent in assessing whether the pipeline introduces internal biases at the level of the model training or not.

2.1 Data generation : simulation

All of the elements concerning the generation of the pre-DC1 supernovae can be summarized in the following list :

- All of the supernova data was generated with SkySurvey ².
- The cosmology that was used is Planck18 found in `astropy.cosmology` ($\Omega_m = 0.30966$) (ref).
- The color and stretch of the supernovae were drawn from a normal distribution centered around 0, with $\sigma(c) = 0.1$ and $\sigma(X_1) = 1$.
- There are no redshift errors added to any supernovae.
- A spectrum is generated for each supernova from ZTF and SNLS drawn from a phase when the SN was observed.
- Magnitude cuts were done before the application of any noise and intrinsic dispersion. For each survey the magnitude cuts are written in table 5.1. This ensures that there are no selection effects added to the simulations.
- The rate of the supernovae were chosen to be Frohmaier+19 (ref).
- The milky way dust extinction was drawn out from the Planck dust map (ref) and the $E(B-V)_{MW}$ values follow the CM89 dust law (ref).
- The addition of the intrinsic dispersion, σ_{int} , is applied on the observed magnitudes of each supernova.
- Lightcurve measurement uncertainties were drawn from `sncosmo` ³ as a function of the skynoise taken from the real observational logs.
- An error snake is also added to the lightcurves drawn from a normal distribution with a dispersion of 5% of the flux. The error snake term is the same for a given band and given night for a given supernova.

ZTF	18.3
SNLS	24.2
HSC	24.6

Tab. 5.1 List of the magnitude cuts on DC1 simulations

²<https://github.com/MickaelRigault/skysurvey>

³<https://sncosmo.readthedocs.io/en/stable/>

2.2 Goals

The goal behind pre-DC1 is to check the consistency of the Georges pipeline in reconstructing the input distances and cosmology. The final results of the analysis chain will be determined after running the Georges pipeline on 100 realizations of the pre-DC1 dataset.

We define the following set of goals to achieve at the end of pre-DC1 :

1. That the reconstructed supernova parameters X_0 , X_1 , c and t_{max} are unbiased
2. That the reconstructed distance moduli μ are unbiased
3. That the reconstructed parameter uncertainties extracted from the inverse of the Fisher matrix (σ_{Fisher}) are equivalent to the parameter uncertainties extracted from the empirical covariance matrix constructed after the 100 realizations (σ_{Emp})
4. That the reconstructed cosmological parameters Ω_m and w are unbiased in the context of a Λ CDM cosmology with a constant w .

3 Results : Potential biases & systematics

3.1 Model & data agreement

The results we present here are obtained on 1 realization of the pre-DC1 simulations. The average time it takes to train one pre-DC1 sample of supernova is X. This is done on the IN2P3 Computing Cluster⁴.

In Figures 5.1 and 5.2 we are plotting the photometric and spectroscopic residuals weighted by the measurement uncertainties. We see that after the model training, we are able to accurately describe the data with NaCl within the model's definition domain. This is also extended in Figure 5.3 when we look at the weighted residuals for each measurement obtained with a specific filter.




Fig. 5.1 Photometric residuals for 1 realization of pre-DC1

3.2 Parameter reconstruction and covariance matrix

Since the pre-DC1 dataset is generated with a model that respects the NaCl constraints, we expect to reconstruct the exact same parameters used in the generation. In Figure 5.4 we compare the generated and reconstructed parameters averaged over all 100 realizations of pre-DC1. We see that the reconstructed

⁴<https://cc.in2p3.fr/en/>




Fig. 5.2 Spectroscopic residuals for 1 realization of pre-DC1

parameters match the input parameters within the statistical uncertainties of the reconstruction.

We are also interested in comparing the constructed empirical covariance matrix and the Fisher covariance matrix from NaCl. In Figure (ref), we compare the output uncertainties (σ_{Fisher}) from 1 pre-DC1 realization with the uncertainties extracted from the empirical covariance matrix (σ_{Emp}) constructed from the run on the 100 pre-DC1 realizations. We see that for each block of parameters X_0 , X_1 and c , σ_{Fisher} is equivalent to σ_{Emp} and well within the statistical uncertainties of σ_{Emp} . This is also highlighted when we look at their ratios where they are centered around 1. We are now convinced that the NaCl can produce unbiased parameters along their uncertainties in a LEMAITRE-like analysis.

3.3 Distance modulus and cosmological parameters reconstruction

We use the EDRIS module to determine the distance modulus of the supernovae at the end of the lightcurve fit and model training with NaCl. It's important to note that we are not assessing the efficiency of EDRIS in reconstructing distances as this has been already established in D. Kuhn's PhD thesis⁵ where it was shown that EDRIS does not bias supernova distances in our setup. If we do find

⁵<https://theses.hal.science/tel-05451069>




Fig. 5.3 Photometric residuals per band for 1 realization of pre-DC1

biased distances, then it would come from NaCl.

The results in Figure 5.5 show the mean binned distance residuals from 100 realizations of pre-DC1 as well as a comparison between the empirical uncertainties on the binned distances and the Fisher uncertainties from EDRIS. We see that NaCl does not introduce biases on the supernova distances. We can now move on to determining the cosmological parameters with cosmologix.

In Figures 5.6 and ref(om and w coverage plots) we show the results from 1 realization of pre-DC1 and from the 100 realizations of pre-DC1. We are plotting the contours for all the Ω_m and w parameters obtained. What we see is that we recover unbiased results within 1σ ensuring that our pipeline works.




Fig. 5.4 Weighted residuals of each supernova parameter for 100 realization of pre-DC1




Fig. 5.5 Hubble diagram for 1 realization of pre-DC1




Fig. 5.6 Ω_m and w contours for 1 realization of pre-DC1

CHAPTER

Field level inference with peculiar velocities from ZTF SN Ia

Contents

1	The BORG forward model	114
1.1	The BORG algorithm	114
1.2	Utility of field level inferences	115
2	Self-consistent simulation analysis	123

1 The BORG forward model

Once we have obtained the distances of supernovae and know the rate of the cosmic expansion we can deduce z_{cosmo} and z_{pv} . The goal is to then be able to measure the growth rate of structures, $f\sigma_8$. There are multiple methods that we can use in order to make that measurement and for the work in this PhD, we decided to use a field level inference, BORG, since it can model non-linearities in the velocity field of the supernovae and allow for a cross-survey analysis being done with the same tool.

1.1 The BORG algorithm

BORG stands for *Bayesian Origin Reconstruction from Galaxies* ([Jasche and Wandelt, 2013]) - a Bayesian inference framework that uses a forward model to tackle the inverse problem of how structures are formed in at the large scales of the Universe. The way BORG can be applied on type Ia Supernova is summarized in Figure 6.1.




Fig. 6.1 Schematic summarizing what BORG intends to do when faced with type Ia Supernova data.

It is a field level approach designed to infer an initial density field δ_{IC} , which would be our initial conditions and forward model its evolution to obtain a final density field δ_f . The Bayesian logic behind BORG makes it that we are exploring a large scale structure posterior by inferring a Gaussian prior for the initial density field δ_{IC} . Modeling the evolving density field from δ_{IC} to δ_f requires a gravity model that can capture the non linear effects of structure formation. I will be explaining the details of the different parts of BORG in the next subsections. In a sense, we are generating, or simulating, a numerical "twin" of our Universe by sampling the posterior of the initial conditions using a Bayesian approach.

1.2 Utility of field level inferences

Here mentioned: knowledge at a deeper level than just parameter sampling, if that were the case a lot of other parameters can be just sampled. Non linearities. A digital twin can allow us to quantify matter clumping and distribution at the level of each halo. etc...

1.2.1 BORG's Bayesian inference chain

The logic behind BORG relies on Bayesian inference, as stated in the name of the algorithm, and this method is very well adapted for the problem we are trying to solve. The Bayesian framework works in the following way : Based on the Bayes theorem, the probability $\mathcal{P}$ of a specific realization of an event or data d being described by a set of parameters θ can be written as :

$$\mathcal{P}(\theta|d) = \frac{\mathcal{P}(d|\theta)\mathcal{P}(\theta)}{\mathcal{P}(d)} \quad (6.1)$$

where $\mathcal{P}(d|\theta)$ is the probability that the data is known given a model which is also called the likelihood, $\mathcal{P}(\theta)$ the probability of the model parameters in the absence of any data, also known as the prior, and $\mathcal{P}(d)$ the probability of the data before it is known, also called the evidence. The probability $\mathcal{P}(\theta|d)$ is called the posterior.

In the context of our problem, determining a prior on the evolved density field is a challenge due to the non-linear gravitational nature of structure formation ([Fis, 1995], [Jasche and Kitaura, 2010]). So even if one assumes a Gaussian or a log-normal for the δ_f they will be missing the description of fine filamentary features of the observed cosmic web ([Peebles, 1980]). Instead what is done in BORG is treat the problem as a Bayesian inference on the initial conditions (δ_{IC}). The biggest gain out of that, is that the prior is physically motivated from results obtained from Planck CMB data ([Ade et al., 2016], [Collaboration et al., 2019]) and inflation which is a Gaussian prior. Combining that with a gravity model to evolve the density field and by marginalizing over the final density fields, we can then constrain our initial conditions.

From [Jasche and Lavaux, 2019] we can obtain the final posterior as :

$$\mathcal{P}(\{\delta_{IC}^{(i)}\}|\{N^{(i)}\}) = \frac{\mathcal{P}(\{\delta_{IC}^{(i)}\})\mathcal{P}(\{N^{(i)}\}|\{G^{(i)}(\{\delta_{IC}^{(i)}\})\})}{\mathcal{P}(\{N^{(i)}\})} \quad (6.2)$$

i being the index of the voxel we are evaluating with $N^{(i)}$ the number of objects within that voxel and $G^{(i)}(\{\delta_{IC}^{(i)}\})$ is the gravity model or the structure formation model defined such that

$$G^{(i)}(\{\delta_{IC}^{(i)}\}) = \delta_f^{(i)}$$

Given the dimensionality of the problem at hand, directly integrating this equation is practically unfeasible due to the computational power required. The solution to that is to then explore the posterior distribution of initial conditions to find the best configuration set that matches our observations. This will require us to use a set of Markov Chain Monte Carlo (MCMC) sampling algorithms.

A Markov Chain is a set of sequences $\{\theta_0, \theta_1, \theta_2, \dots, \theta_n, \dots\}$ where the conditional distribution of θ_{n+1} given the previous elements only depends on θ_n . If this distribution does not depend on n we call it a stationary transition probability, which is what we are interested in. To guarantee that distribution, the following condition must be satisfied :

$$\mathcal{T}(\theta|\theta')\mathcal{P}(\theta') = \mathcal{T}(\theta'|\theta)\mathcal{P}(\theta) \quad (6.3)$$

where $\mathcal{T}$ is the transition probability distribution. A property of that condition is that the probability flux from $\theta \rightarrow \theta'$ is equal to $\theta' \rightarrow \theta$ with respect to the target distribution. This is also known as the "balance flow".

Metropolis-Hastings algorithm

A very common example of an MCMC algorithm is the Metropolis-Hastings algorithm ([Metropolis et al., 1953], [HASTINGS, 1970]). To summarize how it works : given a state θ_n , a new state is drawn from a proposal distribution which is often Gaussian q such that :

$$\theta' \sim q(\theta', \theta_n) \quad (6.4)$$

With a Gaussian distribution we get $\theta' = \theta_n + \epsilon$; $\epsilon \sim \mathcal{N}(0, \sigma)$ with ϵ being a random perturbation drawn from the normal distribution where σ controls the step size. The next step is to then compute an acceptance ratio :

$$\alpha = \min \left(1, \frac{\mathcal{P}(\theta'|d)q(\theta_n|\theta')}{\mathcal{P}(\theta_n|d)q(\theta'|\theta_n)} \right) \quad (6.5)$$

The state then becomes :

$$\theta_{n+1} = \begin{cases} \theta' & \text{With probability } \alpha \\ \theta_n & \text{With probability } 1 - \alpha \end{cases} \quad (6.6)$$

This guarantees that better proposals are more likely to be accepted while worse proposal have a probability proportional to how much worse they are. The problem BORG might have with this algorithm is the step size and the computation time since the complexity of the algorithm is $\mathcal{O}(D)$ where D is the dimension of the parameter space. We expect BORG to have around 10^7 parameters to fit so if the step size that is chosen is too small the chain will only make tiny steps and take a very long time to reach a converged state, while if the step size is too large most proposals will land in low-probability regions and get rejected, also making the computational time too long and inefficient.

HMC algorithm

The Hamiltonian Monte-Carlo algorithm (HMC, [Brooks et al., 2011]) is much more suitable for our problem and is currently what is implemented in BORG. The general idea of HMC is to use concepts from Hamiltonian classical mechanics on high-dimensional problems. It uses gradient information to make significant proposals with a high acceptance rate. Let us define Θ as the position vector containing the a set of parameters to solve, $\mathbf{p}$ the momentum variables and $\mathcal{H}$ the Hamiltonian, the Hamilton equations give us :

$$\frac{d\Theta}{dt} = \frac{\partial \mathcal{H}}{\partial \mathbf{p}} \quad (6.7)$$

$$\frac{d\mathbf{p}}{dt} = -\frac{\partial \mathcal{H}}{\partial \Theta} \quad (6.8)$$

with the Hamiltonian being constant in time. Let $\mathbf{M}$ be the mass matrix which is symmetric and positive definite, then the Hamiltonian is :

$$\mathcal{H} = \frac{1}{2} \mathbf{p}^T \mathbf{M}^{-1} \mathbf{p} + U(\Theta) \quad (6.9)$$

where $U(\Theta)$ is the potential energy and $\frac{1}{2} \mathbf{p}^T \mathbf{M}^{-1} \mathbf{p}$ the kinetic energy. It's important to note that the choice of the mass matrix can strongly impact the performance of the sampler.

The link between Hamiltonian dynamics and probabilities can be established with the concept of canonical distributions from statistical mechanics. The probability distribution over the states and momenta are written as :

$$\mathcal{P}(\Theta, \mathbf{p}) = \frac{1}{Z} \exp\left(\frac{-\mathcal{H}}{k_B T}\right) \quad (6.10)$$

where k_B is the Boltzmann constant, T the temperature of the system and Z is the partition function. We can impose $k_B T = 1$ for simplicity and absorbing the Z partition function into the normalization which leads us to:

$$-\ln \mathcal{P}(\Theta, \mathbf{p}) = \mathcal{H}(\Theta, \mathbf{p}) = \frac{1}{2} \mathbf{p}^T \mathbf{M}^{-1} \mathbf{p} + U(\Theta) \quad (6.11)$$

We can see that the Hamiltonian formalism naturally allows the parameters Θ to be displaced along trajectories of a constant $\mathcal{H}$, guided by the gradient of the potential energy $U(\Theta) = -\ln \mathcal{P}(\Theta, \mathbf{p})$ towards regions of higher posterior probability. This means that the joint distribution is :

$$\mathcal{P}(\Theta, \mathbf{p}) \propto \exp(-\mathcal{H}(\Theta, \mathbf{p})) \quad (6.12)$$

We can then the same acceptance condition as the Metropolis-Hastings algorithm where we have :

$$\alpha(\Theta', \mathbf{p}', \Theta, \mathbf{p}) = \min\left(1, \frac{\mathcal{P}(\Theta', \mathbf{p}')}{\mathcal{P}(\Theta, \mathbf{p})}\right) = \min(1, \exp(-\mathcal{H}(\Theta', \mathbf{p}') + \mathcal{H}(\Theta, \mathbf{p}))) \quad (6.13)$$

In exact Hamiltonian dynamics, $\mathcal{H}$ is conserved so the expected acceptance rate is unity but due to numerical errors we expect to see a lower acceptance rate. Lastly, we marginalize over the momenta to obtain the final posterior on our parameters, for us that will be the initial conditions.

$$\mathcal{P}(\Theta) = \int \mathcal{P}(\Theta, \mathbf{p}) d\mathbf{p} \quad (6.14)$$

1.2.2 Particle mass assignment

The initial density field δ_{IC} is drawn from a Gaussian prior and represented as a set of particles distributed on a grid. The space we are working in is assigned on a grid so it is important to be able to assign the particles to that grid. BORG addresses by doing a Cloud In Cell (CIC) mass assignment on the grid. This mass assignment works by assigning a "cloud" to each particle and distributing its mass to the neighbouring grid points according to the overlap between the cloud and these points.

The assignment is done in the following way : Each particle is treated as a uniform-density cube having a mass of unity. Each particle overlaps with up to 8 neighbouring grid cells. The fraction of the particle's mass assigned to each cell is proportional to the volume overlap between the particle cube and the grid cell. If we consider a 3D grid with a cell size of Δx and a particle at the position $X_p = (x_p, y_p, z_p)$ lives inside a cell whose lower-left corner is a grid point (i, j, k) where

$$i = \left\lfloor \frac{x_p}{\Delta x} \right\rfloor, \quad j = \left\lfloor \frac{y_p}{\Delta x} \right\rfloor, \quad k = \left\lfloor \frac{z_p}{\Delta x} \right\rfloor$$

We define the fractional distance as

$$d_x = \frac{x_p}{\Delta x} - i, \quad d_y = \frac{y_p}{\Delta x} - j, \quad d_z = \frac{z_p}{\Delta x} - k$$

So that $d_x, d_y, d_z \in [0, 1[$.

The particle's mass m_p is distributed to the 8 surrounding nodes $(i+a, j+b, k+c)$ with $a, b, c \in 0, 1$:

$$\rho_{i+a, j+b, k+c} = \frac{m_p}{\Delta x^3} \cdot T_x^{(a)} \cdot T_y^{(b)} \cdot T_z^{(c)} \quad (6.15)$$

with $T_{x,y,z}^0 = 1 - d_{x,y,z}$ and $T_{x,y,z}^1 = d_{x,y,z}$.

1.2.3 Gravity Model

Simply evolving the δ_{IC} field using the equations of motion would be too computationally expensive since the number of particles in our field can go up to 10^9 particle. Developments in the implementation of a gravity model that is accurate and fast have been a very important topic in the development of BORG. I will be presenting the main models that are used in BORG. Additionally, the movements of the particles in the density field directly gives us access to the underlying velocity field.

LPT

The first gravity model is the Lagrangian Perturbation Theory or LPT, it can also be known as the Zeldovich approximation ([Zel'dovich, 1970a], [Crocce and Scoccimarro, 2006]). The idea is to expand the density field perturbatively $\delta(x, t) = \delta^{(1)}(x, t) + \delta^{(2)}(x, t) + \dots$, which is the Eulerian expansion, as well as the displacement field $\Psi(q, t) = \Psi^{(1)}(q, t) + \Psi^{(2)}(q, t) + \dots$ which maps the initial positions q (Lagrangian coordinates) to the final positions x (Eulerian coordinates). In comoving coordinates, the equations of motion for dark matter particles are :

$$\ddot{x} + 2H\dot{x} = -\frac{1}{a^2}\nabla_x\phi \quad (6.16)$$

where $H = \dot{a}/a$ is the Hubble parameter, a the scale factor and ϕ is the gravitational potential satisfying the Poisson equation :

$$\nabla_x^2\phi = 4\pi G\bar{\rho}a^2\delta(x, t) \quad (6.17)$$

With the position being:

$$x(q, t) = q + \Psi(q, t) \quad (6.18)$$

This means we can rewrite the equations of motion as :

$$\ddot{\Psi} + 2H\dot{\Psi} = -\frac{1}{a^2}\nabla_x\phi \quad (6.19)$$

To ensure the mass conservation, the continuity equation in the Lagrangian formalism gives us :

$$1 + \delta(x, t) = \frac{1}{|\det(J)|} \quad (6.20)$$

With J being the Jacobian that maps q to x .

$$J_{ij} = \frac{\partial x_i}{\partial q_j} = \delta_{ij} + \frac{\partial \Psi_i}{\partial q_j} \quad (6.21)$$

At first order, we can expand the determinant $\det(J) \simeq 1 + \nabla_q \cdot \Psi^{(1)}$ giving the first term in the expansion of the density field as :

$$\delta^{(1)}(x, t) \simeq -\nabla_q \cdot \Psi^{(1)} \quad (6.22)$$

If we continue the demonstration, we can obtain what the first order approximation is for the position of the density field particles at a time t and its velocity as:

$$x(q, t) = q - D_1(t)\nabla_q\phi^{(1)}(q) \quad (6.23)$$

$$v(q, t) = -aHfD_1(t)\nabla_q\phi^{(1)}(q) \quad (6.24)$$

Where D_1 is the linear growth factor and $f = d\ln D_1/d\ln a$ the linear growth rate of structures. It's important to note that this approximation breaks down when dense structures start to form such as

halos, therefore this approximation is a good description of structure formation at large scales. Going to higher orders such as 2LPT (2nd order approximation) delays this breakdown but does not prevent it. Physically, it can be shown that this occurs around $k \approx 10^{-1} h\text{Mpc}^{-1}$ at $z = 0$

2LPT

We can also use the second order approximation of the Lagrangian Perturbation Theory (2LPT) as our gravity solver. That means the displacement field is written as $\Psi(q, t) = \Psi^{(1)}(q, t) + \Psi^{(2)}(q, t)$ and one can demonstrate that the second order term of the displacement is :

$$\Psi^{(2)}(q, t) = -D_2(t) \nabla_q \phi^{(2)}(q) \quad (6.25)$$

where $D_2(t)$ is the second order growth factor. We also get the following Poisson equation for the gravitational potential:

$$\nabla_q^2 \phi^{(2)}(q) = \sum_{i>j} \left(\frac{\partial^2 \phi^{(1)}}{\partial^2 x_i} \frac{\partial^2 \phi^{(1)}}{\partial^2 x_j} - \left[\frac{\partial^2 \phi^{(1)}}{\partial x_i x_j} \right]^2 \right) \quad (6.26)$$

The final position and velocity of the density field in 2LPT are :

$$x(q, t) = q - D_1(t) \nabla_q \phi^{(1)} - D_2(t) \nabla_q \phi^{(2)} \quad (6.27)$$

$$v(q, t) = -aH \left[f_1 D_1 \nabla_q \phi^{(1)} + f_2 D_2 \nabla_q \phi^{(2)} \right] \quad (6.28)$$

where f_1 is the linear growth factor $f_1 = d \ln D_1 / d \ln a$ and $f_2 = d \ln D_2 / d \ln a$.

PM

The goal of the Particle Mesh (PM) gravity model is to evolve the particles in the simulation after having done a mass assignment, like CIC. The equations of motion and the Poisson are adapted to be solved on the BORG grid chosen, they are given by Equation 6.18 and $\frac{d^2 \Psi(q, a)}{da^2} = -\nabla_x \phi(x, a)$. I will explaining the steps it takes to evolve the field from a time t_i to a time t_{i+1} without going into the derivation of the equations at each step.

- Mass assignment, like CIC, to obtain the density field δ .
- Transform to Fourier space $\delta(x) \rightarrow \tilde{\delta}(k)$.
- Solve the Poisson equation in Fourier space.
- Compute the force field, or the gravitational acceleration, in Fourier space $\tilde{g}(k)$ then transform it to cartesian space to get $g(x)$.
- Interpolate the forces back to particles.
- Time integration to get the next position and momentum of the particles.

The PM gravity model is accurate up to the resolution of the grid, which is the most accurate gravity model we can use in BORG at the current moment. It allows us to have very accurate density and velocity fields but getting there requires having to repeat the full PM cycle at least 100 times ([Tassev et al., 2013]) which will produce very long computational time. If we decide to reduce the steps in PM we would lose accuracy on all scales.

COLA

The last model I will be presenting is COmoving Lagrangian Acceleration or COLA ([Tassev et al., 2013]). The goal of COLA is to be able to solve the equations of motion and the Poisson equation for particles that undergo trajectories in LPT. This method allows us to trade accuracy at small scales to gain computational speed while maintaining accuracy at large scales. This is done by decomposing the displacement field into two components :

$$\Psi(q, a) = \Psi^{LPT}(q, a) + \Psi_{res}^{COLA}(q, a) \quad (6.29)$$

where $\Psi^{LPT}(q, a)$ is the LPT displacement field and $\Psi_{res}^{COLA}(q, a)$ the residual displacement of each particle as observed from a frame comoving with with an "LPT observer" whose trajectory is defined by Ψ^{LPT} . If we temporarily omitt the Hubble expansion, we get the following equations of motion :

$$\frac{d^2\Psi(q, a)}{da^2} = -\nabla_x\phi(x, a) \quad (6.30)$$

This can be rewritten in the comoving frame of the "LPT" observer as :

$$\frac{d^2\Psi_{res}^{COLA}(q, a)}{da^2} = -\nabla_x\phi(x, a) - \frac{d^2\Psi^{LPT}(q, a)}{da^2} \quad (6.31)$$

We can think of the LPT as an additional force acting on the particles due to the use of a non-inertial frame of reference. LPT also experiences lower typical accelerations compared to the natural cosmological frame. Numerically, this produces equivalent accuracy to solving the full equations of motion while using fewer steps ([Tassev et al., 2013]). The expected number of STEPS in COLA is around 10 unlike PM. A visual representation of what COLA does can be seen in Figure 6.2.




Fig. 6.2 Schematic representing how COLA works found in [Bartlett et al., 2025]. The orange line represents the motion of the particle from the LPT frame of reference and the blue lines are the residuals COLA calculates from the true position to correct for it.

If we denote $x(a) = x_{LPT}(a) + x_{COLA}(a)$ and the momentum $p = p_{LPT} + p_{COLA}$ the equations to solve then become :

$$\frac{dx}{da} = \mathcal{D}(a)p_{COLA} - \frac{dD_1}{d\Psi^{(1)}} + \frac{dD_2}{d\Psi^{(2)}} \quad (6.32)$$

$$\frac{dp_{COLA}}{da} = \mathcal{K}(a) \left[\nabla(\Delta^{-1}\delta) - \mathcal{V}[D_1](a)\Psi^{(1)} + \mathcal{V}[D_2](a)\Psi^{(2)} \right] \quad (6.33)$$

with $\mathcal{D}(a) = \frac{f(a)}{a^2}$, also called the "drift" prefactor, and $\mathcal{K}(a) = -\frac{3}{2}\Omega_m H_0^2 f(a)/a$, also called the "kick" prefactor. These terms come from the kick-drift-kick (KDK) algorithm that is used to update the particle positions. The kick phase boosts the momentum of the particles while the drift phase changes their positions. The KDK scheme goes as follows : At an instant n , positions and momenta at an instant $n+1$ are determined in the following way :

- Kick $p_{n+1/2}^{COLA} = p_n^{COLA} + F_n^{COLA} \cdot \frac{\Delta a}{2}$
- Drift $x_{n+1} = x_n + p_{n+1/2}^{COLA} \cdot \delta a$
- Kick $p_{n+1}^{COLA} = p_{n+1/2}^{COLA} + F_{n+1}^{COLA} \cdot \frac{\Delta a}{2}$

where at each step, the force F^{COLA} is computed using a PM solver from above.

We can look at a visual difference between the 2LPT gravity solver, COLA and what a "true" density field like which is a N-body simulation named GADGET ([Springel, 2005], [Springel et al., 2021]) in Figure 6.3. The 2LPT results are obtained in 3 timesteps, the COLA results in 10 timesteps and GADGET results are obtained in 200 times. As expected, the COLA results manage to capture a lot of the details found on small scales as opposed to the 2LPT results while requiring far fewer timesteps than a full N-body simulation.




Fig. 6.3 Evolution of the same initial density field with three gravity solvers 2LPT, COLA and GADGET

1.2.4 Likelihood components of BORG

Density field likelihood

- sn are matter tracers - poisson likelihood $\rightarrow$ number of sn in voxel

$$\mathcal{P}(N_k^g | \lambda_k) = \prod_k \frac{(\lambda_k)^{N_k^g} e^{-\lambda_k}}{N_k^g!} \quad (6.34)$$

1.2.5 Advantages over summary-statistics analyses

- Retains phase information and non-linearity - analysis through cross-correlations with observables - digital twin of our observable universe may lead to new insights on it

worth citing Richard's paper on H_0 to show far we can go with FLI and what does the velocity field adds to density field information

1.2.6 Numerical implementation

- C++, JAX, HPC

2 Self-consistent simulation analysis

- Simulation description - Different flavours and tests - Results

CHAPTER

BORG applications on ZTF simulations

Contents

1	ZTF peculiar velocity data challenge (DC PV)	126
2	Supernova forward-model analysis	127
2.1	Bias model	127
3	Results of the field level inference	128
4	Comparison with other methods for $f\sigma_8$	128

1 ZTF peculiar velocity data challenge (DC PV)

Assessing the capability of BORG in inferring the density and velocity fields of the Universe using ZTF supernovae requires that our analysis extends the study of self-consistent simulations. The goal is determine whether BORG is capable of inferring the fields when confronted with external data generated using a different method to BORG. This allows us to mimic what we would expect when applying the method on the real data.

This brings to the last part of my work during this PhD: applying BORG on simulations generated within the ZTF collaboration. These set simulations, named DC PV (Data Challenge Peculiar Velocity), are generated within the ZTF collaboration in order to test methods in inferring $f\sigma_8$. The DC PV simulations are done similarly to the DC simulations as described in Chapter 4 with the main differences lie in adding structure information to the simulations. For that, the supernovae are simulated with the help of the Uchuu N-body simulation ([Ishiyama et al., 2021]), a high fidelity N-body dark matter simulation in a $2^3\text{Gpc}.h^{-1}$ box with 2.1 trillion particles which can be seen in Figure (ref)¹. The full box is then subdivided into 27 subboxes where the supernovae are simulated up to a redshift of $z = 0.11$, giving 27 quasi-independent realizations of the DC PV. Each supernova must be simulated where a particle is and is assigned the particle's velocity. This allows us to add proper motion to supernovae that is driven by structure formation.

We opted to simplify a few aspects of the simulations in order to have an initial assessment of how well can a BORG analysis run on ZTF-like supernova. Firstly, we have chosen to use only the distance moduli and redshifts of supernovae as inputs in BORG, as if we were doing the analysis after having run the full LEMAITRE pipeline. One could assume that in more complicated analyses that the Tripp relation parameters (α , β) would be sampled in BORG given that non-homogenous Malmquist bias is accounted for. Secondly, we do not impose the ZTF DR2 cosmology quality cuts on photometry nor the ZTF footprint, so all of the supernovae that are generated can be used. For a more realistic set of simulations, we would have to account for the geometry of the survey as well as the Milky-Way dust. This means introducing an angular mask $M(\text{RA}, \text{Dec})$ in the likelihood's selection function. This method was applied in BORG analyses such as [McAlpine et al., 2025] (Manticore-Local) and [McAlpine et al., 2026] (Manticore-Deep). Finally, we draw a set of 1000 randomly selected supernovae up to a redshift of $z = 0.06$ to match the expected number of supernovae in the volume-limited sample from ZTF. We show in Figure (ref) an example of simulated supernovae on top of the Uchuu simulation.

¹N-body and hydrodynamical simulations are considered the highest fidelity simulations we can get to describe structure formation as they solving the equations of motion for each particle/fluid without any approximations or grid mesh

2 Supernova forward-model analysis

The methodology used to generate supernova in the self consistent simulations and DC PV are different as we have mentioned earlier. This means kamel

- Dont know certain parameters such as α and σ_v so we need to determine - Need to know if there are model inconsistencies or additional noise in ZTF or systematic biases

2.1 Bias model

Results from antoine and ZTF simulation tend to favour non r_2 evolution $\rightarrow$ how do we account for it?

3 Results of the field level inference

- results - covariance additions ? - full dataset with selection?

4 Comparison with other methods for $f\sigma_8$

can compare with results from Rafael

Titre :

Titre ici

Résumé (*en français*) :

—

Mots-clés :

Supernova, galaxie

Title:

Title here

Abstract (*in english*):

—

Keywords:

Supernova, galaxy

Bibliography

[Fis, 1995] (1995). Velocity and potential fields from all-sky galaxy redshift surveys. *Monthly Notices of the Royal Astronomical Society*.

[Adame et al., 2025] Adame, A., Aguilar, J., Ahlen, S., Alam, S., Alexander, D., Allende Prieto, C., Alvarez, M., Alves, O., Anand, A., Andrade, U., Armengaud, E., Avila, S., Aviles, A., Awan, H., Bahr-Kalus, B., Bailey, S., Baltay, C., Bault, A., Behera, J., BenZvi, S., Beutler, F., Bianchi, D., Blake, C., Blum, R., Bonici, M., Brieden, S., Brodzeller, A., Brooks, D., Buckley-Geer, E., Burtin, E., Calderon, R., Canning, R., Carnero Rosell, A., Cereskaite, R., Cervantes-Cota, J., Chabanier, S., Chaussidon, E., Chaves-Montero, J., Chebat, D., Chen, S., Chen, X., Claybaugh, T., Cole, S., Cuceu, A., Davis, T., Dawson, K., de la Macorra, A., de Mattia, A., Deiosso, N., Dey, A., Dey, B., Ding, Z., Doel, P., Edelstein, J., Eftekharzadeh, S., Eisenstein, D., Elbers, W., Elliott, A., Fagrelus, P., Fanning, K., Ferraro, S., Ereza, J., Findlay, N., Flaughner, B., Font-Ribera, A., Forero-Sánchez, D., Forero-Romero, J., Frenk, C., Garcia-Quintero, C., Garrison, L., Gaztañaga, E., Gil-Marín, H., Gontcho, S. A., Gonzalez-Morales, A., Gonzalez-Perez, V., Gordon, C., Green, D., Gruen, D., Gspaner, R., Gutierrez, G., Guy, J., Hadzhiyska, B., Hahn, C., Hanif, M., Herrera-Alcantar, H., Honscheid, K., Howlett, C., Huterer, D., Iršič, V., Ishak, M., Joyce, R., Juneau, S., Karaçaylı, N., Kehoe, R., Kent, S., Kirkby, D., Kong, H., Koposov, S., Kremin, A., Krolewski, A., Lahav, O., Lai, Y., Lan, T.-W., Landriau, M., Lang, D., Lasker, J., Le Goff, J., Le Guillou, L., Leauthaud, A., Levi, M., Li, T., Lodha, K., Magneville, C., Manera, M., Margala, D., Martini, P., Matthewson, W., Maus, M., McDonald, P., Medina-Varela, L., Meisner, A., Mena-Fernández, J., Miquel, R., Moon, J., Moore, S., Moustakas, J., Mudur, N., Mueller, E., Muñoz-Gutiérrez, A., Myers, A., Nadathur, S., Napolitano, L., Neveux, R., Newman, J., Nguyen, N., Nie, J., Niz, G., Noriega, H., Padmanabhan, N., Paillas, E., Palanque-Delabrouille, N., Pan, J., Penmetsa, S., Percival, W., Pieri, M., Pinon, M., Poppett, C., Porredon, A., Prada, F., Pérez-Fernández, A., Pérez-Ràfols, I., Rabinowitz, D., Raichoor, A., Ramírez-Pérez, C., Ramirez-Solano, S., Rashkovetskyi, M., Ravoux, C., Rezaie, M., Rich, J., Rocher, A., Rockosi, C., Roe, N., Rosado-Marin, A., Ross, A., Rossi, G., Ruggeri, R., Ruhlmann-Kleider, V., Samushia, L., Sanchez, E., Saulder, C., Schlafly, E., Schlegel, D., Schubnell, M., Seo, H., Shafieloo, A., Sharples, R., Silber, J., Slosar, A., Smith, A., Sprayberry, D., Tan, T., Tarlé, G., Taylor, P., Trusov, S., Vaisakh, R., Valcin, D., Valdes, F., Valogiannis, G., Vargas-Magaña, M., Verde, L., Walther, M., Wang, B., Wang, M., Weaver, B., Weaverdyck, N., Wechsler, R., Weinberg, D., White, M., Wilson, M., Yi, L., Yu, J., Yu, Y., Yuan, S., Yèche, C., Zaborowski, E., Zarrouk, P., Zhang, H., Zhao, C., Zhao, R., Zhou, R., Zhuang, T., and Zou, H. (2025). Desi 2024 vii: cosmological constraints from the full-shape modeling of clustering measurements. *Journal of Cosmology and Astroparticle Physics*, 2025(07):028.

[Ade et al., 2016] Ade, P. A. R., Aghanim, N., Arnaud, M., Ashdown, M., Aumont, J., Baccigalupi, C., Banday, A. J., Barreiro, R. B., Bartlett, J. G., Bartolo, N., Battaner, E., Battye, R., Benabed, K., Benoît, A., Benoit-Lévy, A., Bernard, J.-P., Bersanelli, M., Bielewicz, P., Bock, J. J., Bonaldi, A., Bonavera, L., Bond, J. R., Borrill, J., Bouchet, F. R., Boulanger, F., Bucher, M., Burigana, C., Butler, R. C., Calabrese, E., Cardoso, J.-F., Catalano, A., Challinor, A., Chamballu, A., Chary, R.-R., Chiang, H. C., Chluba, J., Christensen, P. R., Church, S., Clements, D. L., Colombi, S., Colombo, L. P. L., Combet, C., Coulais, A., Crill, B. P., Curto, A., Cuttaia, F., Danese, L., Davies, R. D., Davis, R. J., de Bernardis, P., de Rosa, A., de Zotti, G., Delabrouille, J., Désert, F.-X., Di Valentino, E., Dickinson, C., Diego, J. M., Dolag, K., Dole, H., Donzelli, S., Doré, O., Douspis, M., Ducout, A., Dunkley, J., Dupac, X., Efstathiou, G., Elsner, F., Enßlin, T. A., Eriksen, H. K., Farhang, M., Fergusson, J., Finelli, F., Forni, O., Frailis, M., Fraisse, A. A., Franceschi, E., Frejsel, A., Galeotta, S., Galli, S., Ganga, K., Gauthier, C., Gerbino, M., Ghosh, T., Giard, M., Giraud-Héraud, Y., Giusarma, E., Gjerløw, E., González-Nuevo, J., Górski, K. M., Gratton, S., Gregorio, A., Gruppuso, A., Gudmundsson, J. E., Hamann, J., Hansen, F. K., Hanson, D., Harrison, D. L., Helou, G., Henrot-Versillé,

- Hornstrup, A., Hovest, W., Huang, Z., Huffenberger, K. M., Hurier, G., Jaffe, A. H., Jaffe, T. R., Jones, W. C., Juvela, M., Keihänen, E., Keskitalo, R., Kisner, T. S., Kneissl, R., Knoche, J., Knox, L., Kunz, M., Kurki-Suonio, H., Lagache, G., Lähteenmäki, A., Lamarre, J.-M., Lasenby, A., Lattanzi, M., Lawrence, C. R., Leahy, J. P., Leonardi, R., Lesgourgues, J., Levrier, F., Lewis, A., Liguori, M., Lilje, P. B., Linden-Vørnle, M., López-Caniego, M., Lubin, P. M., Macías-Pérez, J. F., Maggio, G., Maino, D., Mandolesi, N., Mangilli, A., Marchini, A., Maris, M., Martin, P. G., Martinelli, M., Martínez-González, E., Masi, S., Matarrese, S., McGehee, P., Meinhold, P. R., Melchiorri, A., Melin, J.-B., Mendes, L., Mennella, A., Migliaccio, M., Millea, M., Mitra, S., Miville-Deschênes, M.-A., Moneti, A., Montier, L., Morgante, G., Mortlock, D., Moss, A., Munshi, D., Murphy, J. A., Naselsky, P., Nati, F., Natoli, P., Netterfield, C. B., Nørgaard-Nielsen, H. U., Noviello, F., Novikov, D., Novikov, I., Oxborrow, C. A., Paci, F., Pagano, L., Pajot, F., Paladini, R., Paoletti, D., Partridge, B., Pasian, F., Patanchon, G., Pearson, T. J., Perdureau, O., Perotto, L., Perrotta, F., Pettorino, V., Piacentini, F., Piat, M., Pierpaoli, E., Pietrobon, D., Plaszczynski, S., Pointecouteau, E., Polenta, G., Popa, L., Pratt, G. W., Prézeau, G., Prunet, S., Puget, J.-L., Rachen, J. P., Reach, W. T., Rebolo, R., Reinecke, M., Remazeilles, M., Renault, C., Renzi, A., Ristorcelli, I., Rocha, G., Rosset, C., Rossetti, M., Roudier, G., Rouillé d'Orfeuil, B., Rowan-Robinson, M., Rubiño-Martín, J. A., Rusholme, B., Said, N., Salvatelli, V., Salvati, L., Sandri, M., Santos, D., Savelainen, M., Savini, G., Scott, D., Seiffert, M. D., Serra, P., Shellard, E. P. S., Spencer, L. D., Spinelli, M., Stolyarov, V., Stompor, R., Sudiwala, R., Sunyaev, R., Sutton, D., Suur-Uski, A.-S., Sygnet, J.-F., Tauber, J. A., Terenzi, L., Toffolatti, L., Tomasi, M., Tristram, M., Trombetti, T., Tucci, M., Tuovinen, J., Türler, M., Umana, G., Valenziano, L., Valiviita, J., Van Tent, F., Vielva, P., Villa, F., Wade, L. A., Wandelt, B. D., Wehus, I. K., White, M., White, S. D. M., Wilkinson, A., Yvon, D., Zacchei, A., and Zonca, A. (2016). Planck2015 results: Xiii. cosmological parameters. *Astronomy and Astrophysics*, 594:A13.
- [Aghanim et al., 2020] Aghanim, N., Akrami, Y., Ashdown, M., Aumont, J., Baccigalupi, C., Ballardini, M., Banday, A. J., Barreiro, R. B., Bartolo, N., Basak, S., Battye, R., Benabed, K., Bernard, J.-P., Bersanelli, M., Bielewicz, P., Bock, J. J., Bond, J. R., Borrill, J., Bouchet, F. R., Boulanger, F., Bucher, M., Burigana, C., Butler, R. C., Calabrese, E., Cardoso, J.-F., Carron, J., Challinor, A., Chiang, H. C., Chluba, J., Colombo, L. P. L., Combet, C., Contreras, D., Crill, B. P., Cuttaia, F., de Bernardis, P., de Zotti, G., Delabrouille, J., Delouis, J.-M., Di Valentino, E., Diego, J. M., Doré, O., Douspis, M., Ducout, A., Dupac, X., Dusini, S., Efstathiou, G., Elsner, F., Enßlin, T. A., Eriksen, H. K., Fantaye, Y., Farhang, M., Fergusson, J., Fernandez-Cobos, R., Finelli, F., Forastieri, F., Frailis, M., Fraisse, A. A., Franceschi, E., Frolov, A., Galeotta, S., Galli, S., Ganga, K., Génova-Santos, R. T., Gerbino, M., Ghosh, T., González-Nuevo, J., Górski, K. M., Gratton, S., Gruppuso, A., Gudmundsson, J. E., Hamann, J., Handley, W., Hansen, F. K., Herranz, D., Hildebrandt, S. R., Hivon, E., Huang, Z., Jaffe, A. H., Jones, W. C., Karakci, A., Keihänen, E., Keskitalo, R., Kiiveri, K., Kim, J., Kisner, T. S., Knox, L., Krachmalnicoff, N., Kunz, M., Kurki-Suonio, H., Lagache, G., Lamarre, J.-M., Lasenby, A., Lattanzi, M., Lawrence, C. R., Le Jeune, M., Lemos, P., Lesgourgues, J., Levrier, F., Lewis, A., Liguori, M., Lilje, P. B., Lilley, M., Lindholm, V., López-Caniego, M., Lubin, P. M., Ma, Y.-Z., Macías-Pérez, J. F., Maggio, G., Maino, D., Mandolesi, N., Mangilli, A., Marcos-Caballero, A., Maris, M., Martin, P. G., Martinelli, M., Martínez-González, E., Matarrese, S., Mauri, N., McEwen, J. D., Meinhold, P. R., Melchiorri, A., Mennella, A., Migliaccio, M., Millea, M., Mitra, S., Miville-Deschênes, M.-A., Molinari, D., Montier, L., Morgante, G., Moss, A., Natoli, P., Nørgaard-Nielsen, H. U., Pagano, L., Paoletti, D., Partridge, B., Patanchon, G., Peiris, H. V., Perrotta, F., Pettorino, V., Piacentini, F., Polastri, L., Polenta, G., Puget, J.-L., Rachen, J. P., Reinecke, M., Remazeilles, M., Renzi, A., Rocha, G., Rosset, C., Roudier, G., Rubiño-Martín, J. A., Ruiz-Granados, B., Salvati, L., Sandri, M., Savelainen, M., Scott, D., Shellard, E. P. S., Sirignano, C., Sirri, G., Spencer, L. D., Sunyaev, R., Suur-Uski, A.-S., Tauber, J. A., Tavagnacco, D., Tenti, M., Toffolatti, L., Tomasi, M., Trombetti, T., Valenziano, L., Valiviita, J., Van Tent, B., Vibert, L., Vielva, P., Villa, F., Vittorio, N., Wandelt, B. D., Wehus, I. K., White, M., White, S. D. M., Zacchei, A., and Zonca, A. (2020). Planck 2018 results: Vi. cosmological parameters. *Astronomy and Astrophysics*, 641:A6.
- [Akrami et al., 2020] Akrami, Y., Ashdown, M., Aumont, J., Baccigalupi, C., Ballardini, M., Banday, A. J., Barreiro, R. B., Bartolo, N., Basak, S., Benabed, K., Bersanelli, M., Bielewicz, P., Bock, J. J., Bond, J. R., Borrill, J., Bouchet, F. R., Boulanger, F., Bucher, M., Burigana, C., Butler, R. C., Calabrese, E., Cardoso, J.-F., Casaponsa, B., Chiang, H. C., Colombo, L. P. L., Combet, C., Contreras, D., Crill, B. P., de Bernardis, P., de Zotti, G., Delabrouille, J., Delouis, J.-M., Di Valentino, E., Diego, J. M., Doré, O., Douspis, M., Ducout, A., Dupac, X., Efstathiou, G., Elsner, F., Enßlin, T. A., Eriksen, H. K., Fantaye, Y., Fernandez-Cobos, R., Finelli, F., Frailis, M., Fraisse, A. A., Franceschi, E., Frolov, A., Galeotta, S., Galli, S., Ganga, K., Génova-

- J. E., Hamann, J., Handley, W., Hansen, F. K., Herranz, D., Hivon, E., Huang, Z., Jaffe, A. H., Jones, W. C., Keihänen, E., Keskitalo, R., Kiiveri, K., Kim, J., Krachmalnicoff, N., Kunz, M., Kurki-Suonio, H., Lagache, G., Lamarre, J.-M., Lasenby, A., Lattanzi, M., Lawrence, C. R., Le Jeune, M., Levrier, F., Liguori, M., Lilje, P. B., Lindholm, V., López-Caniego, M., Ma, Y.-Z., Macías-Pérez, J. F., Maggio, G., Maino, D., Mandolesi, N., Mangilli, A., Marcos-Caballero, A., Maris, M., Martin, P. G., Martínez-González, E., Matarrese, S., Mauri, N., McEwen, J. D., Meinhold, P. R., Mennella, A., Migliaccio, M., Miville-Deschênes, M.-A., Molinari, D., Moneti, A., Montier, L., Morgante, G., Moss, A., Natoli, P., Pagano, L., Paoletti, D., Partridge, B., Perrotta, F., Pettorino, V., Piacentini, F., Polenta, G., Puget, J.-L., Rachen, J. P., Reinecke, M., Remazeilles, M., Renzi, A., Rocha, G., Rosset, C., Roudier, G., Rubiño-Martín, J. A., Ruiz-Granados, B., Salvati, L., Savelainen, M., Scott, D., Shellard, E. P. S., Sirignano, C., Sunyaev, R., Suur-Uski, A.-S., Tauber, J. A., Tavagnacco, D., Tenti, M., Toffolatti, L., Tomasi, M., Trombetti, T., Valenziano, L., Valiviita, J., Van Tent, B., Vielva, P., Villa, F., Vittorio, N., Wandelt, B. D., Wehus, I. K., Zacchei, A., Zibin, J. P., and Zonca, A. (2020). Planck2018 results: Vii. isotropy and statistics of the cmb. *Astronomy and Astrophysics*, 641:A7.
- [Alam et al., 2017] Alam, S., Ata, M., Bailey, S., Beutler, F., Bizyaev, D., Blazek, J. A., Bolton, A. S., Brownstein, J. R., Burden, A., Chuang, C.-H., Comparat, J., Cuesta, A. J., Dawson, K. S., Eisenstein, D. J., Escoffier, S., Gil-Marín, H., Grieb, J. N., Hand, N., Ho, S., Kinemuchi, K., Kirkby, D., Kitaura, F., Malanushenko, E., Malanushenko, V., Maraston, C., McBride, C. K., Nichol, R. C., Olmstead, M. D., Oravetz, D., Padmanabhan, N., Palanque-Delabrouille, N., Pan, K., Pellejero-Ibanez, M., Percival, W. J., Petitjean, P., Prada, F., Price-Whelan, A. M., Reid, B. A., Rodríguez-Torres, S. A., Roe, N. A., Ross, A. J., Ross, N. P., Rossi, G., Rubiño-Martín, J. A., Saito, S., Salazar-Albornoz, S., Samushia, L., Sánchez, A. G., Satpathy, S., Schlegel, D. J., Schneider, D. P., Scóccola, C. G., Seo, H.-J., Sheldon, E. S., Simmons, A., Slosar, A., Strauss, M. A., Swanson, M. E. C., Thomas, D., Tinker, J. L., Tojeiro, R., Magaña, M. V., Vazquez, J. A., Verde, L., Wake, D. A., Wang, Y., Weinberg, D. H., White, M., Wood-Vasey, W. M., Yèche, C., Zehavi, I., Zhai, Z., and Zhao, G.-B. (2017). The clustering of galaxies in the completed sdss-iii baryon oscillation spectroscopic survey: cosmological analysis of the dr12 galaxy sample. *Monthly Notices of the Royal Astronomical Society*, 470(3):2617–2652.
- [Albrecht and Steinhardt, 1982] Albrecht, A. and Steinhardt, P. J. (1982). Cosmology for grand unified theories with radiatively induced symmetry breaking. *Phys. Rev. Lett.*, 48:1220–1223.
- [Aldering et al., 2002] Aldering, G., Adam, G., Antilogus, P., Astier, P., Bacon, R., Bongard, S., Bonnaud, C., Copin, Y., Hardin, D., Henault, F., Howell, D. A., Lemonnier, J.-P., Levy, J.-M., Loken, S. C., Nugent, P. E., Pain, R., Pecontal, A., Pecontal, E., Perlmutter, S., Quimby, R. M., Schahmanecche, K., Smadja, G., and Wood-Vasey, W. M. (2002). Overview of the Nearby Supernova Factory. In Tyson, J. A. and Wolff, S., editors, *Survey and Other Telescope Technologies and Discoveries*, volume 4836, pages 61 – 72. International Society for Optics and Photonics, SPIE.
- [Andernach and Zwicky, 2017] Andernach, H. and Zwicky, F. (2017). English and spanish translation of zwicky’s (1933) the redshift of extragalactic nebulae. *arXiv e-prints*.
- [Andrew Howell et al., 2006] Andrew Howell, D., Sullivan, M., Nugent, P. E., Ellis, R. S., Conley, A. J., Le Borgne, D., Carlberg, R. G., Guy, J., Balam, D., Basa, S., Fouchez, D., Hook, I. M., Hsiao, E. Y., Neill, J. D., Pain, R., Perrett, K. M., and Pritchett, C. J. (2006). The type ia supernova snls-03d3bb from a super-chandrasekhar-mass white dwarf star. *Nature*, 443(7109):308–311.
- [Astier et al., 2006] Astier, P., Guy, J., Regnault, N., et al. (2006). The Supernova Legacy Survey: measurement of Ω_M , Ω_Λ and w from the first year data set. *Astronomy and Astrophysics*, 447:31.
- [Augarde, 2022] Augarde, G. (2022). *Caractérisation de l’énergie noire : développement d’une nouvelle génération de modélisation spectrophotométrique empirique des supernovæ de type Ia*. PhD thesis. Thèse de doctorat dirigée par Regnault, Nicolas Physique de l’Univers Sorbonne université 2022.
- [Barbary et al., 2021] Barbary, K., Bailey, S., Barentsen, G., Barclay, T., Biswas, R., Boone, K., Craig, M., Feindt, U., Friesen, B., Goldstein, D., Jha, S. W., Jones, D. O., Mondon, F., Papadogiannakis, S., Perrefort, D., Pierel, J., Rodney, S., Rose, B., Saunders, C., Sipőcz, B., Sofiatti, C., Thomas, R. C., van Santen, J.,

- [Bartelmann and Schneider, 2001] Bartelmann, M. and Schneider, P. (2001). Weak gravitational lensing. *Physics Reports*, 340(4–5):291–472.
- [Bartlett et al., 2025] Bartlett, D. J., Chiarenza, M., Doerer, L., and Leclercq, F. (2025). Comoving computer acceleration (coca): N-body simulations in an emulated frame of reference. *Astronomy and Astrophysics*, 694:A287.
- [Baumont, 2007] Baumont, S. (2007). *Analyse des spectres VLT pour l'expérience SNLS Qualification de transients cosmologiques*. Theses, Université Paris-Diderot - Paris VII.
- [Bessell, 1990] Bessell, M. S. (1990). UBVRI passbands. *Publications of the Astronomical Society of the Pacific*, 102:1181–1199.
- [Betoule et al., 2014] Betoule, M., Kessler, R., Guy, J., Mosher, J., Hardin, D., Biswas, R., Astier, P., El-Hage, P., Konig, M., Kuhlmann, S., Marriner, J., Pain, R., Regnault, N., Balland, C., Bassett, B. A., Brown, P. J., Campbell, H., Carlberg, R. G., Cellier-Holzem, F., Cinabro, D., Conley, A., D’Andrea, C. B., DePoy, D. L., Doi, M., Ellis, R. S., Fabbro, S., Filippenko, A. V., Foley, R. J., Frieman, J. A., Fouchez, D., Galbany, L., Goobar, A., Gupta, R. R., Hill, G. J., Hlozek, R., Hogan, C. J., Hook, I. M., Howell, D. A., Jha, S. W., Le Guillou, L., Leloudas, G., Lidman, C., Marshall, J. L., Möller, A., Mourão, A. M., Neveu, J., Nichol, R., Olmstead, M. D., Palanque-Delabrouille, N., Perlmutter, S., Prieto, J. L., Pritchett, C. J., Richmond, M., Riess, A. G., Ruhlmann-Kleider, V., Sako, M., Schahmanche, K., Schneider, D. P., Smith, M., Sollerman, J., Sullivan, M., Walton, N. A., and Wheeler, C. J. (2014). Improved cosmological constraints from a joint analysis of the sdss-ii and snls supernova samples. *Astronomy and Astrophysics*, 568:A22.
- [Blake and Glazebrook, 2003] Blake, C. and Glazebrook, K. (2003). Probing Dark Energy Using Baryonic Oscillations in the Galaxy Power Spectrum as a Cosmological Ruler. *The Astrophysical Journal*, 594:665–673.
- [Blinnikov et al., 2006] Blinnikov, S. I., Röpke, F. K., Sorokina, E. I., Gieseler, M., Reinecke, M., Travaglio, C., Hillebrandt, W., and Stritzinger, M. (2006). Theoretical light curves for deflagration models of type ia supernova. *Astronomy & Astrophysics*, 453(1):229–240.
- [Boone et al., 2021] Boone, K., Aldering, G., Antilogus, P., Aragon, C., Bailey, S., Baltay, C., Bongard, S., Buton, C., Copin, Y., Dixon, S., Fouchez, D., Gangler, E., Gupta, R., Hayden, B., Hillebrandt, W., Kim, A. G., Kowalski, M., Küsters, D., Léget, P.-F., Mondon, F., Nordin, J., Pain, R., Pecontal, E., Pereira, R., Perlmutter, S., Ponder, K. A., Rabinowitz, D., Rigault, M., Rubin, D., Runge, K., Saunders, C., Smadja, G., Suzuki, N., Tao, C., Taubenberger, S., Thomas, R. C., and Vincenzi, M. (2021). The twins embedding of type ia supernovae. ii. improving cosmological distance estimates. *The Astrophysical Journal*, 912(1):71.
- [Breuval et al., 2025a] Breuval, L., Anand, G. S., Anderson, R. I., Beaton, R., Bhardwaj, A., Casertano, S., Clementini, G., Cruz Reyes, M., De Somma, G., Groenewegen, M. A. T., Huang, C. D., Kervella, P., Khan, S., Macri, L. M., Marconi, M., Minniti, J. H., Riess, A. G., Ripepi, V., Romaniello, M., Scolnic, D., Trentin, E., Wielgórski, P., and Yuan, W. (2025a). Converging on the cepheid metallicity dependence: Implications of nonstandard gaia parallax recalibration on distance measures. *The Astrophysical Journal*, 994(1):111.
- [Breuval et al., 2025b] Breuval, L., Huang, C. D., and Riess, A. G. (2025b). The legacy of henrietta leavitt: A re-analysis of the first cepheid period–luminosity relation. *Publications of the Astronomical Society of the Pacific*, 137(4):044001.
- [Brooks et al., 2011] Brooks, S., Gelman, A., Jones, G., and Meng, X.-L. (2011). *Handbook of Markov Chain Monte Carlo*. Chapman and Hall/CRC.
- [Brout et al., 2022] Brout, D., Scolnic, D., Popovic, B., Riess, A. G., Carr, A., Zuntz, J., et al. (2022). The Pantheon+ Analysis: Cosmological Constraints. *Astrophysical Journal*, 938(2):110.
- [Cardelli et al., 1989] Cardelli, J. A., Clayton, G. C., and Mathis, J. S. (1989). The Relationship between Infrared, Optical, and Ultraviolet Extinction. *Astrophysical Journal*, 345:245.
- [Carreres, 2023] Carreres, B. (2023). *Détermination du taux de croissance des structures avec les supernovae de type Ia à partir du relevé photométrique ZTF*. PhD thesis. Thèse de doctorat dirigée par Fouchez, Dominique

- [Carreres et al., 2023] Carreres, B., Bautista, J. E., Feinstein, F., Fouchez, D., Racine, B., Smith, M., Ame-nouche, M., Aubert, M., Dhawan, S., Ginolin, M., Goobar, A., Gris, P., Lacroix, L., Nuss, E., Regnault, N., Rigault, M., Robert, E., Rosnet, P., Sommer, K., Dekany, R., Groom, S. L., Sravan, N., Masci, F. J., and Purdum, J. (2023). Growth-rate measurement with type-ia supernovae using ztf survey simulations. *Astronomy and Astrophysics*, 674:A197.
- [Carroll et al., 1992] Carroll, S. M., Press, W. H., and Turner, E. L. (1992). The cosmological constant. *Annual Review of Astronomy and Astrophysics*, 30:499–542.
- [Chandrasekhar, 1931] Chandrasekhar, S. (1931). The Maximum Mass of Ideal White Dwarfs. *Astrophysical Journal*, 74:81.
- [Chevallier and Polarski, 2001] Chevallier, M. and Polarski, D. (2001). Accelerating Universes with Scaling Dark Matter. *International Journal of Modern Physics D*, 10:213–224.
- [Chotard, 2011] Chotard, N. (2011). *Étude de la variabilité des Supernovae de type Ia observées par la collaboration Nearby Supernova Factory*. Theses, Université Claude Bernard - Lyon I.
- [Collaboration et al., 2016a] Collaboration, D., Aghamousa, A., Aguilar, J., Ahlen, S., Alam, S., Allen, L. E., Prieto, C. A., Annis, J., Bailey, S., Bolland, C., Ballester, O., Baltay, C., Beaufore, L., Bebek, C., Beers, T. C., Bell, E. F., Bernal, J. L., Besuner, R., Beutler, F., Blake, C., Bleuler, H., Blomqvist, M., Blum, R., Bolton, A. S., Briceno, C., Brooks, D., Brownstein, J. R., Buckley-Geer, E., Burden, A., Burtin, E., Busca, N. G., Cahn, R. N., Cai, Y.-C., Cardiel-Sas, L., Carlberg, R. G., Carton, P.-H., Casas, R., Castander, F. J., Cervantes-Cota, J. L., Claybaugh, T. M., Close, M., Coker, C. T., Cole, S., Comparat, J., Cooper, A. P., Cousinou, M. C., Crocce, M., Cuby, J.-G., Cunningham, D. P., Davis, T. M., Dawson, K. S., de la Macorra, A., Vicente, J. D., Delubac, T., Derwent, M., Dey, A., Dhungana, G., Ding, Z., Doel, P., Duan, Y. T., Ealet, A., Edelstein, J., Eftekharzadeh, S., Eisenstein, D. J., Elliott, A., Escoffier, S., Evatt, M., Fagrelus, P., Fan, X., Fanning, K., Farahi, A., Farihi, J., Favole, G., Feng, Y., Fernandez, E., Findlay, J. R., Finkbeiner, D. P., Fitzpatrick, M. J., Flaughner, B., Flender, S., Font-Ribera, A., Forero-Romero, J. E., Fosalba, P., Frenk, C. S., Fumagalli, M., Gaensicke, B. T., Gallo, G., Garcia-Bellido, J., Gaztanaga, E., Fusillo, N. P. G., Gerard, T., Gershkovich, I., Giannantonio, T., Gillet, D., de Rivera, G. G., Gonzalez-Perez, V., Gott, S., Graur, O., Gutierrez, G., Guy, J., Habib, S., Heetderks, H., Heetderks, I., Heitmann, K., Hellwing, W. A., Herrera, D. A., Ho, S., Holland, S., Honscheid, K., Huff, E., Hutchinson, T. A., Huterer, D., Hwang, H. S., Laguna, J. M. I., Ishikawa, Y., Jacobs, D., Jeffrey, N., Jelinsky, P., Jennings, E., Jiang, L., Jimenez, J., Johnson, J., Joyce, R., Jullo, E., Juneau, S., Kama, S., Karcher, A., Karkar, S., Kehoe, R., Kennamer, N., Kent, S., Kilbinger, M., Kim, A. G., Kirkby, D., Kisner, T., Kitanidis, E., Kneib, J.-P., Kuposov, S., Kovacs, E., Koyama, K., Kremin, A., Kron, R., Kronig, L., Kueter-Young, A., Lacey, C. G., Lafever, R., Lahav, O., Lambert, A., Lampton, M., Landriau, M., Lang, D., Lauer, T. R., Goff, J.-M. L., Guillou, L. L., Suu, A. L. V., Lee, J. H., Lee, S.-J., Leitner, D., Lesser, M., Levi, M. E., L’Huillier, B., Li, B., Liang, M., Lin, H., Linder, E., Loebman, S. R., Lukić, Z., Ma, J., MacCrann, N., Magneville, C., Makarem, L., Manera, M., Manser, C. J., Marshall, R., Martini, P., Massey, R., Matheson, T., McCauley, J., McDonald, P., McGreer, I. D., Meisner, A., Metcalfe, N., Miller, T. N., Miquel, R., Moustakas, J., Myers, A., Naik, M., Newman, J. A., Nichol, R. C., Nicola, A., da Costa, L. N., Nie, J., Niz, G., Norberg, P., Nord, B., Norman, D., Nugent, P., O’Brien, T., Oh, M., Olsen, K. A. G., Padilla, C., Padmanabhan, H., Padmanabhan, N., Palanque-Delabrouille, N., Palmese, A., Pappalardo, D., Pâris, I., Park, C., Patej, A., Peacock, J. A., Peiris, H. V., Peng, X., Percival, W. J., Perruchot, S., Pieri, M. M., Pogge, R., Pollack, J. E., Poppett, C., Prada, F., Prakash, A., Probst, R. G., Rabinowitz, D., Raichoor, A., Ree, C. H., Refregier, A., Regal, X., Reid, B., Reil, K., Rezaie, M., Rockosi, C. M., Roe, N., Ronayette, S., Roodman, A., Ross, A. J., Ross, N. P., Rossi, G., Roza, E., Ruhlmann-Kleider, V., Rykoff, E. S., Sabiu, C., Samushia, L., Sanchez, E., Sanchez, J., Schlegel, D. J., Schneider, M., Schubnell, M., Secroun, A., Seljak, U., Seo, H.-J., Serrano, S., Shafieloo, A., Shan, H., Sharples, R., Sholl, M. J., Shourt, W. V., Silber, J. H., Silva, D. R., Sirk, M. M., Slosar, A., Smith, A., Smoot, G. F., Som, D., Song, Y.-S., Sprayberry, D., Staten, R., Stefanik, A., Tarle, G., Tie, S. S., Tinker, J. L., Tojeiro, R., Valdes, F., Valenzuela, O., Valluri, M., Vargas-Magana, M., Verde, L., Walker, A. R., Wang, J., Wang, Y., Weaver, B. A., Weaverdyck, C., Wechsler, R. H., Weinberg, D. H., White, M., Yang, Q., Yeche, C., Zhang, T., Zhao, G.-B., Zheng, Y., Zhou, X., Zhou, Z., Zhu, Y., Zou, H., and Zu, Y. (2016a). The desi experiment part ii: Instrument design.
- [Collaboration et al., 2016b] Collaboration, D., Aghamousa, A., Aguilar, J., Ahlen, S., Alam, S., Allen, L. E.,

T. C., Bell, E. F., Bernal, J. L., Besuner, R., Beutler, F., Blake, C., Bleuler, H., Blomqvist, M., Blum, R., Bolton, A. S., Briceno, C., Brooks, D., Brownstein, J. R., Buckley-Geer, E., Burden, A., Burtin, E., Busca, N. G., Cahn, R. N., Cai, Y.-C., Cardiel-Sas, L., Carlberg, R. G., Carton, P.-H., Casas, R., Castander, F. J., Cervantes-Cota, J. L., Claybaugh, T. M., Close, M., Coker, C. T., Cole, S., Comparat, J., Cooper, A. P., Cousinou, M. C., Croce, M., Cuby, J.-G., Cunningham, D. P., Davis, T. M., Dawson, K. S., de la Macorra, A., Vicente, J. D., Delubac, T., Derwent, M., Dey, A., Dhungana, G., Ding, Z., Doel, P., Duan, Y. T., Ealet, A., Edelstein, J., Eftekhazadeh, S., Eisenstein, D. J., Elliott, A., Escoffier, S., Evatt, M., Fagrellius, P., Fan, X., Fanning, K., Farahi, A., Farihi, J., Favole, G., Feng, Y., Fernandez, E., Findlay, J. R., Finkbeiner, D. P., Fitzpatrick, M. J., Flaughner, B., Flender, S., Font-Ribera, A., Forero-Romero, J. E., Fosalba, P., Frenk, C. S., Fumagalli, M., Gaensicke, B. T., Gallo, G., Garcia-Bellido, J., Gaztanaga, E., Fusillo, N. P. G., Gerard, T., Gershkovich, I., Giannantonio, T., Gillet, D., de Rivera, G. G., Gonzalez-Perez, V., Gott, S., Graur, O., Gutierrez, G., Guy, J., Habib, S., Heetderks, H., Heetderks, I., Heitmann, K., Hellwing, W. A., Herrera, D. A., Ho, S., Holland, S., Honscheid, K., Huff, E., Hutchinson, T. A., Huterer, D., Hwang, H. S., Laguna, J. M. I., Ishikawa, Y., Jacobs, D., Jeffrey, N., Jelinsky, P., Jennings, E., Jiang, L., Jimenez, J., Johnson, J., Joyce, R., Jullo, E., Juneau, S., Kama, S., Karcher, A., Karkar, S., Kehoe, R., Kennamer, N., Kent, S., Kilbinger, M., Kim, A. G., Kirkby, D., Kisner, T., Kitanidis, E., Kneib, J.-P., Koposov, S., Kovacs, E., Koyama, K., Kremin, A., Kron, R., Kronig, L., Kueter-Young, A., Lacey, C. G., Lafever, R., Lahav, O., Lambert, A., Lampton, M., Landriau, M., Lang, D., Lauer, T. R., Goff, J.-M. L., Guillou, L. L., Suu, A. L. V., Lee, J. H., Lee, S.-J., Leitner, D., Lesser, M., Levi, M. E., L'Huillier, B., Li, B., Liang, M., Lin, H., Linder, E., Loebman, S. R., Lukić, Z., Ma, J., MacCrann, N., Magneville, C., Makarem, L., Manera, M., Manser, C. J., Marshall, R., Martini, P., Massey, R., Matheson, T., McCauley, J., McDonald, P., McGreer, I. D., Meisner, A., Metcalfe, N., Miller, T. N., Miquel, R., Moustakas, J., Myers, A., Naik, M., Newman, J. A., Nichol, R. C., Nicola, A., da Costa, L. N., Nie, J., Niz, G., Norberg, P., Nord, B., Norman, D., Nugent, P., O'Brien, T., Oh, M., Olsen, K. A. G., Padilla, C., Padmanabhan, H., Padmanabhan, N., Palanque-Delabrouille, N., Palmese, A., Pappalardo, D., Pâris, I., Park, C., Patej, A., Peacock, J. A., Peiris, H. V., Peng, X., Percival, W. J., Perruchot, S., Pieri, M. M., Pogge, R., Pollack, J. E., Poppett, C., Prada, F., Prakash, A., Probst, R. G., Rabinowitz, D., Raichoor, A., Ree, C. H., Refregier, A., Regal, X., Reid, B., Reil, K., Rezaie, M., Rockosi, C. M., Roe, N., Ronayette, S., Roodman, A., Ross, A. J., Ross, N. P., Rossi, G., Roza, E., Ruhlmann-Kleider, V., Rykoff, E. S., Sabiu, C., Samushia, L., Sanchez, E., Sanchez, J., Schlegel, D. J., Schneider, M., Schubnell, M., Secroun, A., Seljak, U., Seo, H.-J., Serrano, S., Shafieloo, A., Shan, H., Sharples, R., Sholl, M. J., Shourt, W. V., Silber, J. H., Silva, D. R., Sirk, M. M., Slosar, A., Smith, A., Smoot, G. F., Som, D., Song, Y.-S., Sprayberry, D., Staten, R., Stefanik, A., Tarle, G., Tie, S. S., Tinker, J. L., Tojeiro, R., Valdes, F., Valenzuela, O., Valluri, M., Vargas-Magana, M., Verde, L., Walker, A. R., Wang, J., Wang, Y., Weaver, B. A., Weaverdyck, C., Wechsler, R. H., Weinberg, D. H., White, M., Yang, Q., Yeche, C., Zhang, T., Zhao, G.-B., Zheng, Y., Zhou, X., Zhou, Z., Zhu, Y., Zou, H., and Zu, Y. (2016b). The desi experiment part i: Science, targeting, and survey design.

[Collaboration et al., 2020] Collaboration, P., Aghanim, N., Akrami, Y., Arroja, F., Ashdown, M., Aumont, J., Baccigalupi, C., Ballardini, M., Banday, A. J., Barreiro, R. B., Bartolo, N., Basak, S., Battye, R., Benabed, K., Bernard, J.-P., Bersanelli, M., Bielewicz, P., Bock, J. J., Bond, J. R., Borrill, J., Bouchet, F. R., Boulanger, F., Bucher, M., Burigana, C., Butler, R. C., Calabrese, E., Cardoso, J.-F., Carron, J., Casaponsa, B., Challinor, A., Chiang, H. C., Colombo, L. P. L., Combet, C., Contreras, D., Crill, B. P., Cuttaia, F., de Bernardis, P., de Zotti, G., Delabrouille, J., Delouis, J.-M., Désert, F.-X., Di Valentino, E., Dickinson, C., Diego, J. M., Donzelli, S., Doré, O., Douspis, M., Ducout, A., Dupac, X., Efstathiou, G., Elsner, F., Enßlin, T. A., Eriksen, H. K., Falgarone, E., Fantaye, Y., Fergusson, J., Fernandez-Cobos, R., Finelli, F., Forastieri, F., Frailis, M., Franceschi, E., Frolov, A., Galeotta, S., Galli, S., Ganga, K., Génova-Santos, R. T., Gerbino, M., Ghosh, T., González-Nuevo, J., Górski, K. M., Gratton, S., Gruppuso, A., Gudmundsson, J. E., Hamann, J., Handley, W., Hansen, F. K., Helou, G., Herranz, D., Hildebrandt, S. R., Hivon, E., Huang, Z., Jaffe, A. H., Jones, W. C., Karakci, A., Keihänen, E., Keskitalo, R., Kiiveri, K., Kim, J., Kisner, T. S., Knox, L., Krachmalnicoff, N., Kunz, M., Kurki-Suonio, H., Lagache, G., Lamarre, J.-M., Langer, M., Lasenby, A., Lattanzi, M., Lawrence, C. R., Le Jeune, M., Leahy, J. P., Lesgourgues, J., Levrier, F., Lewis, A., Liguori, M., Lilje, P. B., Lilley, M., Lindholm, V., López-Caniego, M., Lubin, P. M., Ma, Y.-Z., Macías-Pérez, J. F., Maggio, G., Maino, D., Mandolesi, N., Mangilli, A., Marcos-Caballero, A., Maris, M., Martin, P. G., Martinelli, M., Martínez-González, E., Matarrese, S., Mauri, N., McEwen, J. D., Meerburg, P. D., Meinhold, P. R., Melchiorri, A., Mennella, A., Migliaccio, M., Millea, M., Mitra, S., Miville-

- M., Natoli, P., Nørgaard-Nielsen, H. U., Oxborrow, C. A., Pagano, L., Paoletti, D., Partridge, B., Patanchon, G., Pearson, T. J., Peel, M., Peiris, H. V., Perrotta, F., Pettorino, V., Piacentini, F., Polastri, L., Polenta, G., Puget, J.-L., Rachen, J. P., Reinecke, M., Remazeilles, M., Renault, C., Renzi, A., Rocha, G., Rosset, C., Roudier, G., Rubiño-Martín, J. A., Ruiz-Granados, B., Salvati, L., Sandri, M., Savelainen, M., Scott, D., Shellard, E. P. S., Shiraishi, M., Sirignano, C., Sirri, G., Spencer, L. D., Sunyaev, R., Suur-Uski, A.-S., Tauber, J. A., Tavagnacco, D., Tenti, M., Terenzi, L., Toffolatti, L., Tomasi, M., Trombetti, T., Valiviita, J., Van Tent, B., Vibert, L., Vielva, P., Villa, F., Vittorio, N., Wandelt, B. D., Wehus, I. K., White, M., White, S. D. M., Zacchei, A., and Zonca, A. (2020). *planck* 2018 results: I. overview and the cosmological legacy of *planck*. *Astronomy and Astrophysics*, 641:A1.
- [Collaboration et al., 2019] Collaboration, P., Akrami, Y., Arroja, F., Ashdown, M., Aumont, J., Baccigalupi, C., Ballardini, M., Banday, A. J., Barreiro, R. B., Bartolo, N., Basak, S., Benabed, K., Bernard, J. P., Bersanelli, M., Bielewicz, P., Bond, J. R., Borrill, J., Bouchet, F. R., Bucher, M., Burigana, C., Butler, R. C., Calabrese, E., Cardoso, J. F., Casaponsa, B., Challinor, A., Chiang, H. C., Colombo, L. P. L., Combet, C., Crill, B. P., Cuttaia, F., de Bernardis, P., de Rosa, A., de Zotti, G., Delabrouille, J., Delouis, J. M., Valentino, E. D., Diego, J. M., Doré, O., Douspis, M., Ducout, A., Dupac, X., Dusini, S., Efstathiou, G., Elsner, F., Enßlin, T. A., Eriksen, H. K., Fantaye, Y., Fergusson, J., Fernandez-Cobos, R., Finelli, F., Frailis, M., Fraisse, A. A., Franceschi, E., Frolov, A., Galeotta, S., Ganga, K., Génova-Santos, R. T., Gerbino, M., González-Nuevo, J., Górski, K. M., Gratton, S., Gruppuso, A., Gudmundsson, J. E., Hamann, J., Handley, W., Hansen, F. K., Herranz, D., Hivon, E., Huang, Z., Jaffe, A. H., Jones, W. C., Jung, G., Keihänen, E., Keskitalo, R., Kiiveri, K., Kim, J., Krachmalnicoff, N., Kunz, M., Kurki-Suonio, H., Lamarre, J. M., Lasenby, A., Lattanzi, M., Lawrence, C. R., Jeune, M. L., Levrier, F., Lewis, A., Liguori, M., Lilje, P. B., Lindholm, V., López-Caniego, M., Ma, Y. Z., Macías-Pérez, J. F., Maggio, G., Maino, D., Mandolesi, N., Marcos-Caballero, A., Maris, M., Martin, P. G., Martínez-González, E., Matarrese, S., Mauri, N., McEwen, J. D., Meerburg, P. D., Meinhold, P. R., Melchiorri, A., Mennella, A., Migliaccio, M., Miville-Deschênes, M. A., Molinari, D., Moneti, A., Montier, L., Morgante, G., Moss, A., Münchmeyer, M., Natoli, P., Oppizzi, F., Pagano, L., Paoletti, D., Partridge, B., Patanchon, G., Perrotta, F., Pettorino, V., Piacentini, F., Polenta, G., Puget, J. L., Rachen, J. P., Racine, B., Reinecke, M., Remazeilles, M., Renzi, A., Rocha, G., Rubiño-Martín, J. A., Ruiz-Granados, B., Salvati, L., Savelainen, M., Scott, D., Shellard, E. P. S., Shiraishi, M., Sirignano, C., Sirri, G., Smith, K., Spencer, L. D., Stanco, L., Sunyaev, R., Suur-Uski, A. S., Tauber, J. A., Tavagnacco, D., Tenti, M., Toffolatti, L., Tomasi, M., Trombetti, T., Valiviita, J., Tent, B. V., Vielva, P., Villa, F., Vittorio, N., Wandelt, B. D., Wehus, I. K., Zacchei, A., and Zonca, A. (2019). Planck 2018 results. ix. constraints on primordial non-gaussianity.
- [Crocce and Scoccimarro, 2006] Crocce, M. and Scoccimarro, R. (2006). Renormalized cosmological perturbation theory. *Physical Review D*, 73(6).
- [Croft et al., 1998] Croft, R. A. C., Weinberg, D. H., Katz, N., and Hernquist, L. (1998). Recovery of the Power Spectrum of Mass Fluctuations from Observations of the Lyman-alpha Forest. *The Astrophysical Journal*, 495:44–62.
- [Dawson et al., 2016] Dawson, K. S., Kneib, J.-P., Percival, W. J., Alam, S., Albareti, F. D., Anderson, S. F., Armengaud, E., et al. (2016). The sdss-iv extended baryon oscillation spectroscopic survey: Overview and early data. *The Astronomical Journal*, 151(2):44.
- [de Rham et al., 2011] de Rham, C., Gabadadze, G., and Tolley, A. J. (2011). Resummation of Massive Gravity. *Physical Review Letters*, 106:231101.
- [Deffayet, 2001] Deffayet, C. (2001). Cosmology on a brane in Minkowski bulk. *Physics Letters B*, 502:199–208.
- [Dekany et al., 2020] Dekany, R., Smith, R. M., Riddle, R., Feeney, M., Porter, M., Hale, D., Zolkower, J., Belicki, J., Kaye, S., Henning, J., Walters, R., Cromer, J., Delacroix, A., Rodriguez, H., Reiley, D. J., Mao, P., Hover, D., Murphy, P., Burruss, R., Baker, J., Kowalski, M., Reif, K., Mueller, P., Bellm, E., Graham, M., and Kulkarni, S. R. (2020). The zwicky transient facility: Observing system. *Publications of the Astronomical Society of the Pacific*, 132(1009):038001.
- [Delubac et al., 2015] Delubac, T. et al. (2015). Baryon acoustic oscillations in the Ly α forest of BOSS DR11

- [DES Collaboration et al., 2024] DES Collaboration, Abbott, T. M. C., Acevedo, M., Agüena, M., et al. (2024). The Dark Energy Survey: Cosmology Results With ~ 1500 New High-redshift Type Ia Supernovae Using the Full 5-year Dataset. *The Astrophysical Journal Letters*.
- [DES Collaboration et al., 2022a] DES Collaboration, Abbott, T. M. C., et al. (2022a). Dark Energy Survey Year 3 results: Cosmological constraints from galaxy clustering and weak lensing. *Physical Review D*, 105:023520.
- [DES Collaboration et al., 2022b] DES Collaboration, Abbott, T. M. C., et al. (2022b). Dark Energy Survey Year 3 Results: Cosmological Constraints from Galaxy Clustering and Weak Lensing. *Physical Review D*, 105:023520.
- [DESI Collaboration et al., 2025] DESI Collaboration, Abdul-Karim, M., et al. (2025). DESI DR2 Results II: Measurements of Baryon Acoustic Oscillations and Cosmological Constraints. *Physical Review D*, 112:083515.
- [DESI Collaboration et al., 2016] DESI Collaboration, Aghamousa, A., et al. (2016). The DESI Experiment Part I: Science, Targeting, and Survey Design. *arXiv e-prints*.
- [Di Valentino et al., 2021] Di Valentino, E. et al. (2021). In the realm of the Hubble tension — a review of solutions. *Classical and Quantum Gravity*, 38:153001.
- [Dodelson, 2003] Dodelson, S. (2003). *Modern Cosmology*.
- [Dvali et al., 2000] Dvali, G., Gabadadze, G., and Porrati, M. (2000). 4D gravity on a brane in 5D Minkowski space. *Physics Letters B*, 485:208–214.
- [Einstein, 1916] Einstein, A. (1916). Die Grundlage der allgemeinen Relativitätstheorie. *Annalen der Physik*, 49:769–822.
- [Eisenstein et al., 2005] Eisenstein, D. J. et al. (2005). Detection of Baryon Acoustic Oscillations in the Correlation Function of a Large Sample of Red Galaxies. *The Astrophysical Journal*, 633:560–574.
- [Eisenstein and Hu, 1998] Eisenstein, D. J. and Hu, W. (1998). Baryonic Features in the Matter Transfer Function. *The Astrophysical Journal*, 496:605–614.
- [Elias et al., 1985] Elias, J. H., Matthews, K., Neugebauer, G., and Persson, S. E. (1985). Type I supernovae in the infrared and their use as distance indicators. *Astrophysical Journal*, 296:379–389.
- [Fang et al., 2008] Fang, W., Hu, W., and Lewis, A. (2008). Challenges to the DGP model from horizon-scale growth and geometry. *Physical Review D*, 78:103509.
- [Fierz and Pauli, 1939] Fierz, M. and Pauli, W. (1939). On relativistic wave equations for particles of arbitrary spin in an electromagnetic field. *Proceedings of the Royal Society of London A*, 173:211–232.
- [Fixsen, 2009] Fixsen, D. J. (2009). The Temperature of the Cosmic Microwave Background. *The Astrophysical Journal*, 707:916–920.
- [Freedman, 2021] Freedman, W. L. (2021). Measurements of the hubble constant: Tensions in perspective*. *The Astrophysical Journal*, 919(1):16.
- [Frohmaier et al., 2019] Frohmaier, C., Sullivan, M., Nugent, P. E., Smith, M., Dimitriadis, G., Bloom, J. S., Cenko, S. B., Kasliwal, M. M., Kulkarni, S. R., Maguire, K., Ofek, E. O., Poznanski, D., and Quimby, R. M. (2019). The volumetric rate of normal type Ia supernovae in the local Universe discovered by the Palomar Transient Factory. *Monthly Notices of the Royal Astronomical Society*, 486(2):2308–2320.
- [Ginolin et al., 2025a] Ginolin, M., Rigault, M., Copin, Y., Popovic, B., Dimitriadis, G., Goobar, A., Johansson, J., Maguire, K., Nordin, J., Smith, M., Aubert, M., Barjou-Delayre, C., Burgaz, U., Carreres, B., Dhawan, S., Deckers, M., Feinstein, F., Fouchez, D., Galbany, L., Ganot, C., de Jaeger, T., Kim, Y.-L., Kuhn, D., Lacroix, L., Müller-Bravo, T. E., Nugent, P., Racine, B., Rosnet, P., Rosselli, D., Ruppin, F., Sollerman, J., Terwel, J. H., Townsend, A., Dekany, R., Graham, M., Kasliwal, M., Groom, S. L., Purdum, J., Rusholme, B., and van der Walt, S. (2025a). Ztf sn ia dr2: Colour standardisation of type ia supernovae and its dependence on

- [Ginolin et al., 2025b] Ginolin, M., Rigault, M., Smith, M., Copin, Y., Ruppin, F., Dimitriadis, G., Goobar, A., Johansson, J., Maguire, K., Nordin, J., Amenouche, M., Aubert, M., Barjou-Delayre, C., Betoule, M., Burgaz, U., Carreres, B., Deckers, M., Dhawan, S., Feinstein, F., Fouchez, D., Galbany, L., Ganot, C., Harvey, L., de Jaeger, T., Kenworthy, W. D., Kim, Y.-L., Kowalski, M., Kuhn, D., Lacroix, L., Müller-Bravo, T. E., Nugent, P., Popovic, B., Racine, B., Rosnet, P., Rosselli, D., Sollerman, J., Terwel, J. H., Townsend, A., Brugger, J., Bellm, E. C., Kasliwal, M. M., Kulkarni, S., Laher, R. R., Masci, F. J., Riddle, R. L., and Sharma, Y. (2025b). Ztf sn ia dr2: Environmental dependencies of stretch and luminosity for a volume-limited sample of 1000 type ia supernovae. *Astronomy and Astrophysics*, 695:A140.
- [Gordon et al., 2007] Gordon, C., Land, K., and Slosar, A. (2007). Cosmological constraints from type ia supernovae peculiar velocity measurements. *Physical Review Letters*, 99(8).
- [Gunn and Peterson, 1965] Gunn, J. E. and Peterson, B. A. (1965). On the Density of Neutral Hydrogen in Intergalactic Space. *The Astrophysical Journal*, 142:1633–1641.
- [Guth, 1981] Guth, A. H. (1981). Inflationary universe: A possible solution to the horizon and flatness problems. *Phys. Rev. D*, 23:347–356.
- [Guy et al., 2007] Guy, J., Astier, P., Baumont, S., Hardin, D., Pain, R., Regnault, N., Basa, S., Carlberg, R. G., Conley, A., Fabbro, S., Fouchez, D., Hook, I. M., Howell, D. A., Perrett, K., Pritchett, C. J., Rich, J., Sullivan, M., Antilogus, P., Aubourg, E., Bazin, G., Bronder, J., Filiol, M., Palanque-Delabrouille, N., Ripoche, P., and Ruhlmann-Kleider, V. (2007). Salt2: using distant supernovae to improve the use of type ia supernovae as distance indicators. *Astronomy & Astrophysics*, 466(1):11–21.
- [Guy et al., 2010] Guy, J., Sullivan, M., Conley, A., et al. (2010). The Supernova Legacy Survey 3-year sample: Type Ia supernovae photometric distances and cosmological constraints. *Astronomy and Astrophysics*, 523:A7.
- [Guzzo et al., 2008] Guzzo, L. et al. (2008). A test of the nature of cosmic acceleration using galaxy redshift distortions. *Nature*, 451:541–544.
- [Hamuy et al., 1995] Hamuy, M., Phillips, M. M., Maza, J., Suntzeff, N. B., Schommer, R. A., and Aviles, R. (1995). A Hubble Diagram of Distant Type Ia Supernovae. *Astronomical Journal*, 109:1.
- [Hamuy et al., 1996] Hamuy, M., Phillips, M. M., Suntzeff, N. B., et al. (1996). The Hubble diagram of the Calan/Tololo Type Ia supernovae and the value of H_0 . *The Astronomical Journal*, 112:2398.
- [Harrison, 1970] Harrison, E. R. (1970). Fluctuations at the threshold of classical cosmology. *Phys. Rev. D*, 1:2726–2730.
- [HASTINGS, 1970] HASTINGS, W. K. (1970). Monte carlo sampling methods using markov chains and their applications. *Biometrika*, 57(1):97–109.
- [Heymans et al., 2021a] Heymans, C. et al. (2021a). KiDS-1000 Cosmology: Multi-probe weak gravitational lensing and spectroscopic galaxy clustering. *Astronomy & Astrophysics*, 646:A140.
- [Heymans et al., 2021b] Heymans, C., Tröster, T., Asgari, M., Blake, C., Hildebrandt, H., Joachimi, B., Kuijken, K., Lin, C.-A., Sánchez, A. G., van den Busch, J. L., Wright, A. H., Amon, A., Bilicki, M., de Jong, J., Crocce, M., Dvornik, A., Erben, T., Fortuna, M. C., Getman, F., Giblin, B., Glazebrook, K., Hoekstra, H., Joudaki, S., Kannawadi, A., Köhlinger, F., Lidman, C., Miller, L., Napolitano, N. R., Parkinson, D., Schneider, P., Shan, H., Valentijn, E. A., Verdoes Kleijn, G., and Wolf, C. (2021b). Kids-1000 cosmology: Multi-probe weak gravitational lensing and spectroscopic galaxy clustering constraints. *Astronomy and Astrophysics*, 646:A140.
- [Hillier, Desmond John and Dessart, Luc, 2019] Hillier, Desmond John and Dessart, Luc (2019). Photometric and spectroscopic diversity of type ii supernovae. *Astronomy and Astrophysics*, 631:A8.
- [Hubble, 1929] Hubble, E. (1929). A Relation between Distance and Radial Velocity among Extra-Galactic Nebulae. *Proceedings of the National Academy of Sciences*, 15(3):168–173.
- [Iben and Tutukov, 1984] Iben, Jr., I. and Tutukov, A. V. (1984). Supernovae of type I as end products of the evolution of binaries with components of moderate initial mass. *Astrophysical Journal, Suppl. Ser.*,

- [Ishiyama et al., 2021] Ishiyama, T., Prada, F., Klypin, A. A., Sinha, M., Metcalf, R. B., Jullo, E., Altieri, B., Cora, S. A., Croton, D., de la Torre, S., Millán-Calero, D. E., Oogi, T., Ruedas, J., and Vega-Martínez, C. A. (2021). The uchuu simulations: Data release 1 and dark matter halo concentrations. *Monthly Notices of the Royal Astronomical Society*, 506(3):4210–4231.
- [Jackson, 1972] Jackson, J. C. (1972). A critique of Rees’s theory of primordial gravitational radiation. *Monthly Notices of the Royal Astronomical Society*, 156:1P–5P.
- [Jasche and Kitaura, 2010] Jasche, J. and Kitaura, F. S. (2010). Fast hamiltonian sampling for large-scale structure inference. *Monthly Notices of the Royal Astronomical Society*, 407(1):29–42.
- [Jasche and Lavaux, 2019] Jasche, J. and Lavaux, G. (2019). Physical bayesian modelling of the non-linear matter distribution: New insights into the nearby universe. *Astronomy and Astrophysics*, 625:A64.
- [Jasche and Wandelt, 2013] Jasche, J. and Wandelt, B. D. (2013). Bayesian physical reconstruction of initial conditions from large-scale structure surveys. *Monthly Notices of the Royal Astronomical Society*, 432(2):894–913.
- [Kaiser, 1987] Kaiser, N. (1987). Clustering in real space and in redshift space. *Monthly Notices of the Royal Astronomical Society*, 227:1–21.
- [Kamionkowski et al., 1997] Kamionkowski, M., Kosowsky, A., and Stebbins, A. (1997). A Probe of Primordial Gravity Waves and Vorticity. *Physical Review Letters*, 78:2058–2061.
- [Kenworthy et al., 2021] Kenworthy, W. D., Jones, D. O., Dai, M., Kessler, R., Scolnic, D., Brout, D., Siebert, M. R., Pierel, J. D. R., Dettman, K. G., Dimitriadis, G., Foley, R. J., Jha, S. W., Pan, Y.-C., Riess, A., Rodney, S., and Rojas-Bravo, C. (2021). Salt3: An improved type ia supernova model for measuring cosmic distances. *The Astrophysical Journal*, 923(2):265.
- [Khokhlov et al., 1993] Khokhlov, A., Mueller, E., and Hoefflich, P. (1993). Light curves of type IA supernova models with different explosion mechanisms. *Astronomy and Astrophysics*, 270:223–248.
- [Knobel, 2013] Knobel, C. (2013). An introduction into the theory of cosmological structure formation.
- [Koyama, 2008] Koyama, K. (2008). The cosmological constant and dark energy in braneworlds. *General Relativity and Gravitation*, 40:421–451.
- [Lavaux and Hudson, 2011] Lavaux, G. and Hudson, M. J. (2011). The 2M++ galaxy redshift catalogue. *Monthly Notices of the Royal Astronomical Society*, 416(4):2840–2856.
- [Leavitt, 1908] Leavitt, H. S. (1908). 1777 variables in the Magellanic Clouds. *Annals of Harvard College Observatory*, 60:87–108.3.
- [Lemaître, 1927] Lemaître, G. (1927). Un univers homogène de masse constante et de rayon croissant rendant compte de la vitesse radiale des nébuleuses extra-galactiques. *Annales de la Société Scientifique de Bruxelles*, 47:49–59.
- [Lemaître, 1931] Lemaître, G. (1931). A Homogeneous Universe of Constant Mass and Increasing Radius accounting for the Radial Velocity of Extra-Galactic Nebulae. *Monthly Notices of the Royal Astronomical Society*, 91:483–490.
- [Lenz et al., 2022] Lenz, D., Yeh, R., Mahadevan, V., Grindeanu, I., and Peterka, T. (2022). Adaptive regularization of b-spline models for scientific data.
- [Li et al., 2011] Li, W., Leaman, J., Chornock, R., Filippenko, A. V., Poznanski, D., Ganeshalingam, M., Wang, X., Modjaz, M., Jha, S., Foley, R. J., and Smith, N. (2011). Nearby supernova rates from the lick observatory supernova search – ii. the observed luminosity functions and fractions of supernovae in a complete sample. *Monthly Notices of the Royal Astronomical Society*, 412(3):1441–1472.
- [Linder, 2003] Linder, E. V. (2003). Exploring the Expansion History of the Universe. *Physical Review Letters*, 90:091301.
- [Lordet et al., 2026] Lordet, A. G., Goobar, A., Jasche, J., McAlpine, S., Sollerman, J., Kim, Y.-L., Rigault, M., Ginolin, M., Burgaz, U., Bellm, E. C., Graham, M. J., Jaimes, J. C., Masci, F. J., Purdum, J., and

- [Léget et al., 2020] Léget, P.-F., Gangler, E., Mondon, F., Aldering, G., Antilogus, P., Aragon, C., Bailey, S., Baltay, C., Barbary, K., Bongard, S., Boone, K., Buton, C., Chotard, N., Copin, Y., Dixon, S., Fagrelus, P., Feindt, U., Fouchez, D., Hayden, B., Hillebrandt, W., Kim, A., Kowalski, M., Kuesters, D., Lombardo, S., Lin, Q., Nordin, J., Pain, R., Pecontal, E., Pereira, R., Perlmutter, S., Ponder, K. A., Pruzhinskaya, M. V., Rabinowitz, D., Rigault, M., Runge, K., Rubin, D., Saunders, C., Says, L.-P., Smadja, G., Sofiatti, C., Suzuki, N., Taubenberger, S., Tao, C., and Thomas, R. C. (2020). Sugar: An improved empirical model of type ia supernovae based on spectral features. *Astronomy & Astrophysics*, 636:A46.
- [McAlpine et al., 2025] McAlpine, S., Jasche, J., Ata, M., Lavaux, G., Stiskalek, R., Frenk, C. S., and Jenkins, A. (2025). The manticore project i: a digital twin of our cosmic neighbourhood from bayesian field-level analysis.
- [McAlpine et al., 2026] McAlpine, S., Jasche, J., Lavaux, G., Doerer, L., and Loureiro, A. (2026). The manticore project ii: Bayesian digital twins of cosmic structure across the sdss and boss volumes.
- [Metropolis et al., 1953] Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., and Teller, E. (1953). Equation of State Calculations by Fast Computing Machines. *jcp*, 21(6):1087–1092.
- [Mukhanov et al., 1992] Mukhanov, V., Feldman, H., and Brandenberger, R. (1992). Theory of cosmological perturbations. *Physics Reports*, 215(5):203–333.
- [Mukhanov, 1988] Mukhanov, V. F. (1988). Quantum theory of gauge invariant cosmological perturbations. *Soviet Physics JETP*, 67:1297–1302.
- [Nomoto, 1982] Nomoto, K. (1982). Accreting white dwarf models for type I supernovae. I - Presupernova evolution and triggering mechanisms. *Astrophysical Journal*, 253:798–810.
- [Oppenheimer and Volkoff, 1939] Oppenheimer, J. R. and Volkoff, G. M. (1939). On Massive Neutron Cores. *Physical Review*, 55(4):374–381.
- [Pakmor et al., 2024] Pakmor, R., Seitzzahl, I. R., Ruiter, A. J., Sim, S. A., Röpke, F. K., Taubenberger, S., Bieri, R., and Blondin, S. (2024). Type ia supernova explosion models are inherently multidimensional. *Astronomy & Astrophysics*, 686:A227.
- [Palanque-Delabrouille et al., 2015] Palanque-Delabrouille, N. et al. (2015). Constraint on neutrino masses from the Lyman-alpha forest power spectrum with BOSS DR11 Quasars. *Journal of Cosmology and Astroparticle Physics*, 2015:045.
- [Parrent et al., 2016] Parrent, J. T., Milisavljevic, D., Soderberg, A. M., and Parthasarathy, M. (2016). Line identifications of type i supernovae: On the detection of si ii for these hydrogen-poor events. *The Astrophysical Journal*, 820(1):75.
- [Peebles, 1980] Peebles, P. J. E. (1980). *The large-scale structure of the universe*.
- [Peebles and Yu, 1970] Peebles, P. J. E. and Yu, J. T. (1970). Primeval Adiabatic Perturbation in an Expanding Universe. *Astrophysical Journal*, 162:815.
- [Penzias and Wilson, 1965a] Penzias, A. A. and Wilson, R. W. (1965a). A Measurement of Excess Antenna Temperature at 4080 Mc/s. *Astrophysical Journal*, 142:419–421.
- [Penzias and Wilson, 1965b] Penzias, A. A. and Wilson, R. W. (1965b). Measurement of the Flux Density of CAS a at 4080 Mc/s. *Astrophysical Journal*, 142:1149.
- [Pereira et al., 2013] Pereira, R., Thomas, R. C., Aldering, G., Antilogus, P., Baltay, C., Benitez-Herrera, S., Bongard, S., Buton, C., Canto, A., Cellier-Holzem, F., Chen, J., Childress, M., Chotard, N., Copin, Y., Fakhouri, H. K., Fink, M., Fouchez, D., Gangler, E., Guy, J., Hillebrandt, W., Hsiao, E. Y., Kerschhaggl, M., Kowalski, M., Kromer, M., Nordin, J., Nugent, P., Paech, K., Pain, R., Pécontal, E., Perlmutter, S., Rabinowitz, D., Rigault, M., Runge, K., Saunders, C., Smadja, G., Tao, C., Taubenberger, S., Tilquin, A., and Wu, C. (2013). Spectrophotometric time series of sn 2011fe from the nearby supernova factory. *Astronomy*

- [Perley et al., 2020] Perley, D. A., Fremling, C., Sollerman, J., Miller, A. A., Dahiwal, A. S., Sharma, Y., Bellm, E. C., Biswas, R., Brink, T. G., Bruch, R. J., De, K., Dekany, R., Drake, A. J., Duev, D. A., Filippenko, A. V., Gal-Yam, A., Goobar, A., Graham, M. J., Graham, M. L., Ho, A. Y. Q., Irani, I., Kasliwal, M. M., Kim, Y.-L., Kulkarni, S. R., Mahabal, A., Masci, F. J., Modak, S., Neill, J. D., Nordin, J., Riddle, R. L., Soumagnac, M. T., Strotjohann, N. L., Schulze, S., Taggart, K., Tzanidakis, A., Walters, R. S., and Yan, L. (2020). The Zwicky Transient Facility Bright Transient Survey. II. A Public Statistical Sample for Exploring Supernova Demographics. *Astrophysical Journal*, 904(1):35.
- [Perlmutter et al., 1999] Perlmutter, S., Aldering, G., Goldhaber, G., Knop, R. A., Nugent, P., Castro, P. G., Deustua, S., Fabbro, S., Goobar, A., Groom, D. E., Hook, I. M., Kim, A. G., Kim, M. Y., Lee, J. C., Nunes, N. J., Pain, R., Pennypacker, C. R., Quimby, R., Lidman, C., Ellis, R. S., Irwin, M., McMahon, R. G., Ruiz-Lapuente, P., Walton, N., Schaefer, B., Boyle, B. J., Filippenko, A. V., Matheson, T., Fruchter, A. S., Panagia, N., Newberg, H. J. M., Couch, W. J., and Project, T. S. C. (1999). Measurements of Ω and Λ from 42 High-Redshift Supernovae. *Astrophysical Journal*, 517(2):565–586.
- [Phillips, 1993] Phillips, M. M. (1993). The Absolute Magnitudes of Type IA Supernovae. *Astrophysical Journal Letters*, 413:L105.
- [Planck Collaboration et al., 2020] Planck Collaboration, Akrami, Y., et al. (2020). Planck 2018 results. X. Constraints on inflation. *Astronomy & Astrophysics*, 641:A10.
- [Popovic et al., 2025] Popovic, B. et al. (2025). DES-Dovekie: Improved Cosmological Constraints from a Reanalysis of the DES SN5YR Dataset. *arXiv e-prints*.
- [Riess et al., 1998] Riess, A. G. et al. (1998). Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *The Astronomical Journal*, 116:1009–1038.
- [Riess et al., 1996] Riess, A. G., Press, W. H., and Kirshner, R. P. (1996). A precise distance indicator: Type ia supernova multicolor light-curve shapes. *The Astrophysical Journal*, 473(1):88–109.
- [Riess et al., 2022] Riess, A. G., Yuan, W., Macri, L. M., Scolnic, D., Brout, D., Casertano, S., et al. (2022). A Comprehensive Measurement of the Local Value of the Hubble Constant with $1 \text{ km s}^{-1} \text{ Mpc}^{-1}$ Uncertainty from the Hubble Space Telescope and the SH0ES Team. *The Astrophysical Journal Letters*, 934(1):L7.
- [Rigault et al., 2020] Rigault, M., Brinnel, V., Aldering, G., Antilogus, P., Aragon, C., Bailey, S., Baltay, C., Barbary, K., Bongard, S., Boone, K., Buton, C., Childress, M., Chotard, N., Copin, Y., Dixon, S., Fagrelus, P., Feindt, U., Fouchez, D., Gangler, E., Hayden, B., Hillebrandt, W., Howell, D. A., Kim, A., Kowalski, M., Kuesters, D., Leget, P.-F., Lombardo, S., Lin, Q., Nordin, J., Pain, R., Pecontal, E., Pereira, R., Perlmutter, S., Rabinowitz, D., Runge, K., Rubin, D., Saunders, C., Smadja, G., Sofiatti, C., Suzuki, N., Taubenberger, S., Tao, C., and Thomas, R. C. (2020). Strong dependence of Type Ia supernova standardization on the local specific star formation rate. *Astronomy and Astrophysics*, 644:A176.
- [Roman, M. et al., 2018] Roman, M., Hardin, D., Betoule, M., Astier, P., Balle, C., Ellis, R. S., Fabbro, S., Guy, J., Hook, I., Howell, D. A., Lidman, C., Mitra, A., Möller, A., Mourão, A. M., Neveu, J., Palanque-Delabrouille, N., Pritchet, C. J., Regnault, N., Ruhlmann-Kleider, V., Saunders, C., and Sullivan, M. (2018). Dependence of type ia supernova luminosities on their local environment. *Astronomy and Astrophysics*, 615:A68.
- [Röpke and Hillebrandt, 2004] Röpke, F. K. and Hillebrandt, W. (2004). The case against the progenitor’s carbon-to-oxygen ratio as a source of peak luminosity variations in type Ia supernovae. *Astronomy and Astrophysics*, 420:L1–L4.
- [Rubin et al., 2023] Rubin, D., Aldering, G., Betoule, M., Fruchter, A., Huang, X., Kim, A. G., et al. (2023). Union Through UNITY: Cosmology with 2,000 SNe Using a Unified Bayesian Framework. *Astrophysical Journal*. submitted to ApJ.
- [Rubin et al., 2024] Rubin, D., Aldering, G., Betoule, M., Fruchter, A., Huang, X., Kim, A. G., Lidman, C., Linder, E., Perlmutter, S., Ruiz-Lapuente, P., and Suzuki, N. (2024). Union through unity: Cosmology with 2,000 sne using a unified bayesian framework.
- [Röpke and Sim, 2018] Röpke, F. K. and Sim, S. A. (2018). Models for type ia supernovae and related astro-

- [Sarkar et al., 2009] Sarkar, P., Yadav, J., Pandey, B., and Bharadwaj, S. (2009). The scale of homogeneity of the galaxy distribution in sdss dr6. *Monthly Notices of the Royal Astronomical Society: Letters*, 399(1):L128–L131.
- [Sasaki, 1986] Sasaki, M. (1986). Large scale quantum fluctuations in the inflationary universe. *Progress of Theoretical Physics*, 76:1036–1046.
- [Saunders et al., 2018] Saunders, C., Aldering, G., Antilogus, P., Bailey, S., Baltay, C., Barbary, K., Baugh, D., Boone, K., Bongard, S., Buton, C., Chen, J., Chotard, N., Copin, Y., Dixon, S., Fagrelus, P., Fakhouri, H. K., Feindt, U., Fouchez, D., Gangler, E., Hayden, B., Hillebrandt, W., Kim, A. G., Kowalski, M., Küsters, D., Leget, P.-F., Lombardo, S., Nordin, J., Pain, R., Pecontal, E., Pereira, R., Perlmutter, S., Rabinowitz, D., Rigault, M., Rubin, D., Runge, K., Smadja, G., Sofiatti, C., Suzuki, N., Tao, C., Taubenberger, S., Thomas, R. C., and Vincenzi, M. (2018). Snemo: Improved empirical models for type ia supernovae. *The Astrophysical Journal*, 869(2):167.
- [Sotiriou and Faraoni, 2010] Sotiriou, T. P. and Faraoni, V. (2010). f(R) theories of gravity. *Reviews of Modern Physics*, 82:451–497.
- [Springel, 2005] Springel, V. (2005). The cosmological simulation code gadget-2. *Monthly Notices of the Royal Astronomical Society*, 364(4):1105–1134.
- [Springel et al., 2021] Springel, V., Pakmor, R., Zier, O., and Reinecke, M. (2021). Simulating cosmic structure formation with the <sc>gadget</sc>-4 code. *Monthly Notices of the Royal Astronomical Society*, 506(2):2871–2949.
- [Stein et al., 2022] Stein, G., Seljak, U., Böhm, V., Aldering, G., Antilogus, P., Aragon, C., Bailey, S., Baltay, C., Bongard, S., Boone, K., Buton, C., Copin, Y., Dixon, S., Fouchez, D., Gangler, E., Gupta, R., Hayden, B., Hillebrandt, W., Karmen, M., Kim, A. G., Kowalski, M., Küsters, D., Léget, P.-F., Mondon, F., Nordin, J., Pain, R., Pecontal, E., Pereira, R., Perlmutter, S., Ponder, K. A., Rabinowitz, D., Rigault, M., Rubin, D., Runge, K., Saunders, C., Smadja, G., Suzuki, N., Tao, C., Taubenberger, S., Thomas, R. C., and Vincenzi, M. (2022). A probabilistic autoencoder for type ia supernova spectral time series. *The Astrophysical Journal*, 935(1):5.
- [Sullivan et al., 2010] Sullivan, M., Conley, A., Howell, D. A., Neill, J. D., Astier, P., Balland, C., Basa, S., Carlberg, R. G., Fouchez, D., Guy, J., Hardin, D., Hook, I. M., Pain, R., Palanque-Delabrouille, N., Perrett, K. M., Pritchett, C. J., Regnault, N., Rich, J., Ruhlmann-Kleider, V., Baumont, S., Hsiao, E., Kronborg, T., Lidman, C., Perlmutter, S., and Walker, E. S. (2010). The dependence of Type Ia Supernovae luminosities on their host galaxies. *Monthly Notices of the Royal Astronomical Society*, 406(2):782–802.
- [Tanikawa et al., 2019] Tanikawa, A., Nomoto, K., Nakasato, N., and Maeda, K. (2019). Double-detonation models for type ia supernovae: Trigger of detonation in companion white dwarfs and signatures of companions’ stripped-off materials. *The Astrophysical Journal*, 885(2):103.
- [Tassev et al., 2013] Tassev, S., Zaldarriaga, M., and Eisenstein, D. J. (2013). Solving large scale structure in ten easy steps with cola. *Journal of Cosmology and Astroparticle Physics*, 2013(06):036–036.
- [Tripp and Branch, 1999] Tripp, R. and Branch, D. (1999). Determination of the hubble constant using a two-parameter luminosity correction for type ia supernovae. *The Astrophysical Journal*, 525(1):209.
- [Vincenzi et al., 2024] Vincenzi, M., Brout, D., Armstrong, P., Popovic, B., Taylor, G., Acevedo, M., Camilleri, R., Chen, R., Davis, T. M., Hinton, S. R., Kelsey, L., Kessler, R., Lee, J., Lidman, C., Möller, A., Qu, H., Sako, M., Sanchez, B., Scolnic, D., Smith, M., Sullivan, M., Wiseman, P., Asorey, J., Bassett, B. A., Carollo, D., Carr, A., Foley, R. J., Frohmaier, C., Galbany, L., Glazebrook, K., Graur, O., Kovacs, E., Kuehn, K., Malik, U., Nichol, R. C., Rose, B., Tucker, B. E., Toy, M., Tucker, D. L., Yuan, F., Abbott, T. M. C., Agüena, M., Alves, O., Andrade-Oliveira, F., Annis, J., Bacon, D., Bechtol, K., Bernstein, G. M., Brooks, D., Burke, D. L., Rosell, A. C., Carretero, J., Castander, F. J., Conselice, C., da Costa, L. N., Pereira, M. E. S., Desai, S., Diehl, H. T., Doel, P., Ferrero, I., Flaugher, B., Friedel, D., Frieman, J., García-Bellido, J., Gatti, M., Giannini, G., Gruen, D., Gruendl, R. A., Hollowood, D. L., Honscheid, K., Huterer, D., James, D. J., Kuropatkin, N., Lahav, O., Lee, S., Lin, H., Marshall, J. L., Mena-Fernández, J., Menanteau, F., Miquel, R., Palmese, A., Pieres, A., Malagón, A. A. P., Porredon, A., Romer, A. K., Roodman, A., Sanchez,

- A. R., and Weaverdyck, N. (2024). The dark energy survey supernova program: Cosmological analysis and systematic uncertainties.
- [Wands et al., 2000] Wands, D., Malik, K. A., Lyth, D. H., and Liddle, A. R. (2000). New approach to the evolution of cosmological perturbations on large scales. *Physical Review D*, 62:043527.
- [Wang et al., 2012] Wang, X., Wang, L., Filippenko, A. V., Baron, E., Kromer, M., Jack, D., Zhang, T., Aldering, G., Antilogus, P., Arnett, W. D., Baade, D., Barris, B. J., Benetti, S., Bouchet, P., Burrows, A. S., Canal, R., Cappellaro, E., Carlberg, R. G., di Carlo, E., Challis, P. J., Crotts, A. P. S., Danziger, J. I., Valle, M. D., Fink, M., Foley, R. J., Fransson, C., Gal-Yam, A., Garnavich, P. M., Gerardy, C. L., Goldhaber, G., Hamuy, M., Hillebrandt, W., Höflich, P., Holland, S. T., Holz, D. E., Hughes, J. P., Jeffery, D. J., Jha, S. W., Kasen, D., Khokhlov, A. M., Kirshner, R. P., Knop, R. A., Kozma, C., Krisciunas, K., Lee, B. C., Leibundgut, B., Lentz, E. J., Leonard, D. C., Lewin, W. H. G., Li, W., Livio, M., Lundqvist, P., Maoz, D., Matheson, T., Mazzali, P. A., Meikle, P., Miknaitis, G., Milne, P. A., Mochnacki, S. W., Nomoto, K., Nugent, P. E., Oran, E. S., Panagia, N., Perlmutter, S., Phillips, M. M., Pinto, P., Poznanski, D., Pritchett, C. J., Reinecke, M., Riess, A. G., Ruiz-Lapuente, P., Scalzo, R. A., Schlegel, E. M., Schmidt, B. P., Siegrist, J., Soderberg, A. M., Sollerman, J., Sonneborn, G., Spadafora, A., Spyromilio, J., Sramek, R. A., Starrfield, S. G., Strolger, L. G., Suntzeff, N. B., Thomas, R. C., Tonry, J. L., Tornambe, A., Truran, J. W., Turatto, M., Turner, M., Van Dyk, S. D., Weiler, K. W., Wheeler, J. C., Wood-Vasey, M., Woosley, S. E., and Yamaoka, H. (2012). Evidence for type ia supernova diversity from ultraviolet observations with the hubble space telescope. *The Astrophysical Journal*, 749(2):126.
- [Wheeler and Harkness, 1990] Wheeler, J. C. and Harkness, R. P. (1990). Type I supernovae. *Reports on Progress in Physics*, 53(12):1467–1557.
- [Wheeler and Levreault, 1985] Wheeler, J. C. and Levreault, R. (1985). The peculiar type I supernova in NGC 991. *Astrophysical Journal*, 294:L17–L20.
- [York et al., 2000] York, D. G. et al. (2000). The Sloan Digital Sky Survey: Technical Summary. *The Astrophysical Journal*, 120:1579–1587.
- [Zel’dovich, 1970a] Zel’dovich, Y. B. (1970a). Gravitational instability: An approximate theory for large density perturbations. *Astronomy and Astrophysics*, 5:84–89.
- [Zel’dovich, 1970b] Zel’dovich, Y. B. (1970b). The Hypothesis of Cosmological Magnetic Inhomogeneity. *Sovast*, 13:608.
- [Zwicky, 1933] Zwicky, F. (1933). Die rotverschiebung von extragalaktischen nebeln. *Helvetica Physica Acta*, 6:110–127.