
Urinary bladder

3- Shape of the bladder :


The empty bladder is pyramidal, having an apex, a base, a superior and two inferolateral surfaces; it also has a neck (inferior angle) and 2 postero-superior angles.



Apex: points anteriorly and lies behind the upper margin of the symphysis pubis. It is connected to the umbilicus by the median umbilical ligament (Obliterated urachus).

ANU MEDICAL SCHOOL PROGRAM


- The empty bladder has the shape of upside-down pyramid having:




Urinary bladder

Base (posterior surface):

In males, it is related to seminal vesicles, vasa deferentia, rectovesical pouch containing small intestine and pelvic colon.






ANU MEDICAL SCHOOL PROGRAM

Urinary bladder

Base (posterior surface):

In females, it is related to the cervix and vagina. The ureters open in the postero-superior angles.




ANU
MEDICAL
SCHOOL
PROGRAM

Urinary bladder

Neck:

Lies inferiorly and is connected with the urethra. It is held in position by the **puboprostatic ligaments (male);** these are called the **pubovesical ligaments (female).** The muscle layer of the bladder here forms the internal urethral sphincter.





ANU MEDICAL SCHOOL PROGRAM

Urinary bladder

Superior surface:

In males. It is related to coils of small intestine and sigmoid colon. In females, it is related to the uterovesical pouch and uterus.







ANU MEDICAL SCHOOL PROGRAM

Urinary bladder

Inferolateral surfaces:

Related to the retropubic fat, obturator internus muscle and levator ani muscles.





ANU MEDICAL SCHOOL PROGRAM