
Urinary bladder

Interior of the urinary bladder:

The mucous membrane of the greater part of the empty bladder is thrown into folds that disappear when the bladder is full.

Trigone of the bladder: is a triangular area on the inner surface of the base of the bladder. Here, the mucous membrane is smooth.

The superior angles of the trigone correspond to the openings of the ureters, and the inferior angle to the internal urethral orifice.




ALEXANDRIA NATIONAL UNIVERSITY
ANU MEDICAL SCHOOL PROGRAM

Urinary bladder

Arterial supply:


Superior and inferior vesical arteries, branches of the internal iliac arteries.



Urinary bladder

Venous drainage:


The veins form the vesical venous plexus that drains into the internal iliac vein.



ANU MEDICAL SCHOOL PROGRAM

Urinary bladder

Lymph Drainage: internal and external iliac nodes.




ANU MEDICAL SCHOOL PROGRAM

Urinary bladder

Nerve supply:

The **sympathetic** inhibits contraction of the muscle of the bladder and stimulates the internal urethra sphincter.

The **parasympathetic** stimulates the muscle of the bladder and inhibits the internal urethra sphincter.



Hall: Guyton and Hall Textbook of Medical Physiology, 12th Edition
Copyright © 2011 by Saunders, an imprint of Elsevier, Inc. All rights reserved.

ANU MEDICAL SCHOOL PROGRAM