
Genital Ducts in the Female

ALEXANDRIA NATIONAL UNIVERSITY
ANU MEDICAL SCHOOL PROGRAM


Genital Ducts in the Female

In the presence of estrogen and the absence of testosterone and antimullerian hormone, the paramesonephric ducts develop into the main genital ducts of the female




Paramesonephric duct in female


- It lies first lateral to the mesonephric duct, crosses it anteriorly to its medial side and then grows downwards where it joins its fellow of the opposite side to form the **uterovaginal canal**



Paramesonephric duct in female

- The caudal part of the uterovaginal canal projects into the posterior wall of the urogenital sinus causing a small swelling called Mullerian tubercle.






a
Indifferent gonad
Mesonephric duct
Paramesonephric duct
Müller's tubercle
b
Ovary
Involuting mesonephric duct
Septal resorption
Sinovaginal node
c
Suspensory ligament
Ovary
Ovarian ligament
Epoöphoron
Paroöphoron
Uterus
Hydatid
Involuting mesonephric duct
Round ligament
Gartner's duct cyst

Paramesonephric duct in female:

1. Cranial vertical portion forms the uterine tube.
2. Middle horizontal part that crosses the mesonephric duct, forms uterine horns.
3. A caudal vertical part forming the uterovaginal canal that gives rise to the body

and cervix of the uterus and the upper 3/5 of the vagina.





ALEXANDRIA NATIONAL UNIVERSITY
ANU MEDICAL SCHOOL PROGRAM