
Pelvic floor

- It separates the pelvic cavity above from the perineum below. It slopes inferiorly and medially.
- It is formed of levator ani and coccygeus muscles on both sides.
- Levator ani muscles form the anterior part.
- Coccygeus muscles form the posterior part.



ANU MEDICAL SCHOOL PROGRAM

Levator ani

- The anterior part of the pelvic floor.


- Origin : body of the pubic bone, tendinous arch (linear thickening of the obturator fascia) and ischial spine.

- Insertion : the muscle is divided into 3 parts :

1- puborectalis.


2- pubococcygeus.

3- ischiococcygeus.



Levator ani


- Puborectalis :
- It forms a U-shaped sling with its opposite side around the anorectal junction.





Levator ani


- pubococcygeus : is attached to the fibers of the opposite side in the midline around the urethra and vagina (**pubovaginalis**) in females or around the urethra below the prostate (**puboprostaticus**) in males, perineal body, anal sphincters, anococcygeal body and coccyx.

- Iliococcygeus : is attached to the anococcygeal body and tip of the coccyx.






Pelvic diaphragm
Pubic symphysis
Levator ani
Puborectalis
Pubococcygeus
Iliococcygeus
Coccygeus
Sacro-iliac joint
Urogenital hiatus
Perineal body
Ischial tuberosity
External anal sphincter
Anococcygeal ligament (body)
Greater sciatic foramen
Sacrospinous ligament
* Plane of (B)


Pre-rectal fibres
(of puborectalis)
Urethral hiatus
Puborectalis
Vaginal canal
Pubococcygeus
Iliococcygeus
Rectal hiatus
Coccygeus