
Development of Urinary Bladder

The primitive urogenital sinus is connected to the allantois which extends to the umbilicus. It also receives the 2 mesonephric ducts.

- It differentiates into the following portions:

A- Vescical Part: the upper part that forms most of the urinary bladder and is continuous with the allantois.




ANU MEDICAL SCHOOL PROGRAM

Development of Urinary Bladder


B- Pelvic part: the middle part. It gives rise to the prostatic and membranous urethra in males and the entire urethra in females.

C- Phalic part: the lower part that grows toward the genital tubercle. It will give rise to the penile urethra in males and vestibule of the vagina in females.




ANU MEDICAL SCHOOL PROGRAM

Development of Urinary Bladder



Development of Urinary Bladder




Development of Urinary Bladder

- **The trigone** develops from absorbed parts of the mesonephric ducts (mesoderm).


- **The muscle layer** develops from the surrounding splachnic mesoderm.

- At birth, the urinary bladder is an abdominal organ. It becomes a pelvic organ at puberty.



Development of Urinary Bladder

- Initially, the bladder is continuous with the **allantois** which constricts and becomes a thick fibrous cord, the **urachus**. In adults, the urachus is represented by the **median umbilical ligament**.



ANU MEDICAL SCHOOL PROGRAM