

Paramesonephric duct in female


- It lies first lateral to the mesonephric duct, crosses it anteriorly to its medial side and then grows downwards where it joins its fellow of the opposite side to form the **uterovaginal canal**



Paramesonephric duct in female

- The caudal part of the uterovaginal canal projects into the posterior wall of the urogenital sinus causing a small swelling called Mullerian tubercle.






a
Indifferent gonad
Mesonephric duct
Paramesonephric duct
Müller's tubercle
b
Ovary
Involuting mesonephric duct
Septal resorption
Sinovaginal node
c
Suspensory ligament
Ovary
Ovarian ligament
Epoöphoron
Paroöphoron
Uterus
Hydatid
Involuting mesonephric duct
Round ligament
Gartner's duct cyst

Paramesonephric duct in female:

1. Cranial vertical portion forms the uterine tube.
2. Middle horizontal part that crosses the mesonephric duct, forms uterine horns.
3. A caudal vertical part forming the uterovaginal canal that gives rise to the body


and cervix of the uterus and the upper 3/5 of the vagina.




Paramesonephric duct
Cranial
Middle
Caudal (uterovaginal canal)
Fallopian tube
Uterine horns
Body , Cervix , upper 3/5 of vagina

Mesonephric duct and tubules in female

- The mesonephric duct degenerates except for Gartner's duct.
- The mesonephric tubules degenerate except epoophron and paroophron (in the broad ligament).



ANU MEDICAL SCHOOL PROGRAM