

ECEN 227 - Introduction to Finite Automata and Discrete Mathematics

Dr. Mahmoud Nabil
mnmahmoud@ncat.edu

North Carolina A & T State University

January 27, 2020

Talk Overview

- 1 Introduction
- 2 Floor and Ceiling
- 3 Function Properties
- 4 Function Inverse
- 5 Composition of Functions

Outline


- 1 Introduction
- 2 Floor and Ceiling
- 3 Function Properties
- 4 Function Inverse
- 5 Composition of Functions

Introduction

Function

A function f that maps elements of a set X to elements of a set Y , is a subset of $X \times Y$ such that for every $x \in X$, there is **exactly one** $y \in Y$ for which $(x, y) \in f$

Arrow Diagram of Function




$$X = \{ w, x, y, z \}$$

$$A = \{ a, b, c, d \}$$

$$f = \{ (w, a), (x, a), (y, d), (z, c) \}$$

- $f: X \rightarrow Y$ means f is a function from X to Y .

Arrow Diagram of Function




$$X = \{ w, x, y, z \}$$

$$A = \{ a, b, c, d \}$$

$$f = \{ (w, a), (x, a), (y, d), (z, c) \}$$

- $f: X \rightarrow Y$ means f is a function from X to Y .
- The set X is called the **domain** of f .

Arrow Diagram of Function



$$X = \{ w, x, y, z \}$$

$$A = \{ a, b, c, d \}$$


$$f = \{ (w, a), (x, a), (y, d), (z, c) \}$$

- $f: X \rightarrow Y$ means f is a function from X to Y .
- The set X is called the **domain** of f .
- The set Y is the **target** of f .

Well defined function

Well defined function

f should map every element in the domain to **exactly one element in the target** to be well defined function.



$$X = \{ w, x, y, z \}$$

$$A = \{ a, b, c, d \}$$


$$f = \{ (w, a), (x, a), (y, d), (z, c) \}$$

(Well Defined Function)

Well defined function

Well defined function

f should map every element in the domain to **exactly one element in the target** to be well defined function.




(Not a Function)

Well defined function

Well defined function

f should map every element in the domain to **exactly one element in the target** to be well defined function.




f is no longer a function because $(y, b), (y, d) \in f$.

(Not a Function)

Range

Range

For function $f: X \rightarrow Y$, an element y is in the range of f if and only if there is an $x \in X$ such that $(x, y) \in f$.



$$X = \{w, x, y, z\}$$

$$A = \{a, b, c, d\}$$

$$f = \{(w, a), (x, a), (y, c), (z, c)\}$$

Range: $\{a, c, d\}$

Excercise on Function Range

Express the range of each function using roster notation.

- Let $A = \{2, 3, 4, 5\}$.
f: $A \rightarrow \mathbb{Z}$ such that $f(x) = 2x - 1$.

Exercise on Function Range

Express the range of each function using roster notation.

- Let $A = \{2, 3, 4, 5\}$.
f: $A \rightarrow \mathbb{Z}$ such that $f(x) = 2x - 1$.
 - $\{3, 5, 7, 9\}$
- Let $A = \{2, 3, 4, 5\}$.
f: $A \times A \rightarrow \mathbb{Z}$, where $f(x,y) = x+y$.

Exercise on Function Range

Express the range of each function using roster notation.

- Let $A = \{2, 3, 4, 5\}$.
 $f: A \rightarrow \mathbb{Z}$ such that $f(x) = 2x - 1$.
 - $\{3, 5, 7, 9\}$
- Let $A = \{2, 3, 4, 5\}$.
 $f: A \times A \rightarrow \mathbb{Z}$, where $f(x,y) = x+y$.
 - $\{4, 5, 6, 7, 8, 9, 10\}$

Function Equality

Two functions, f and g , are equal if

- f and g have the **same domain**.
- f and g have the **same target**.
- $f(x) = g(x)$ for every element x in the domain.

Excercise on Function Equality

Ex. Indicate if f and g are equal fuctions

- $f: \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x) = x^2$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = |x|^2$.

Exercise on Function Equality

Ex. Indicate if f and g are equal functions

- $f: \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x) = x^2$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = |x|^2$.
 - $f = g$
- $f: \mathbb{R} \rightarrow \mathbb{Z}$, where $f(x) = x^2$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = x^2$.

Exercise on Function Equality

Ex. Indicate if f and g are equal functions

- $f: \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x) = x^2$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = |x|^2$.
 - $f = g$
- $f: \mathbb{R} \rightarrow \mathbb{Z}$, where $f(x) = x^2$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = x^2$.
 - $f \neq g$ different domains
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x) = x^3$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = |x|^3$.

Exercise on Function Equality

Ex. Indicate if f and g are equal functions

- $f: \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x) = x^2$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = |x|^2$.
 - $f = g$
- $f: \mathbb{R} \rightarrow \mathbb{Z}$, where $f(x) = x^2$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = x^2$.
 - $f \neq g$ different domains
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x) = x^3$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = |x|^3$.
 - $f \neq g$ because, $f(-2) = -8$, and $g(-2) = 8$.
- $f: \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x,y) = |x + y|$
 $g: \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x,y) = |x| + |y|$.

Exercise on Function Equality

Ex. Indicate if f and g are equal functions

- $f: \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x) = x^2$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = |x|^2$.
 - $f = g$
- $f: \mathbb{R} \rightarrow \mathbb{Z}$, where $f(x) = x^2$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = x^2$.
 - $f \neq g$ different domains
- $f: \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x) = x^3$
 $g: \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x) = |x|^3$.
 - $f \neq g$ because, $f(-2) = -8$, and $g(-2) = 8$.
- $f: \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$, where $f(x,y) = |x + y|$
 $g: \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$, where $g(x,y) = |x| + |y|$.
 - $f \neq g$ because, $f(-2,2) = 0$, and $g(-2,2) = 4$.

Outline

- 1 Introduction
- 2 Floor and Ceiling**
- 3 Function Properties
- 4 Function Inverse
- 5 Composition of Functions

Excercise on Function Range

Express the range of each function using roster notation.

Floor function

The floor function maps a **real number** to the **nearest integer** in the downward direction.

$$\begin{aligned}\text{floor}: \mathbb{R} &\rightarrow \mathbb{Z} \\ \text{floor}(x) &= \lfloor x \rfloor\end{aligned}$$



Cieling function

The floor function maps a **real number** to the **nearest integer** in the upward direction.

$$\begin{aligned}\text{ceil}: \mathbb{R} &\rightarrow \mathbb{Z} \\ \text{ceil}(x) &= \lceil x \rceil\end{aligned}$$



Examples

To compute the floor function slide *down* to nearest integer:





Examples

To compute the ceiling function slide *up* to nearest integer:



Examples

The ceiling and floor of an integer are the same:



Outline

- 1 Introduction
- 2 Floor and Ceiling
- 3 Function Properties**
- 4 Function Inverse
- 5 Composition of Functions

Function Properties

One-to-one

A function $f: X \rightarrow Y$ is **one-to-one or injective** if $x_1 \neq x_2$ implies that $f(x_1) \neq f(x_2)$.

Function Properties

One-to-one

A function $f: X \rightarrow Y$ is **one-to-one or injective** if $x_1 \neq x_2$ implies that $f(x_1) \neq f(x_2)$.

Onto

A function $f: X \rightarrow Y$ is **onto or surjective** if the range of f is equal to the target Y .

Function Properties

One-to-one

A function $f: X \rightarrow Y$ is **one-to-one or injective** if $x_1 \neq x_2$ implies that $f(x_1) \neq f(x_2)$.

Onto

A function $f: X \rightarrow Y$ is **onto or surjective** if the range of f is equal to the target Y .

Bijjective

A function is **bijjective or (one-to-one correspondence)** if it is both one-to-one and onto.

In formal definations

One-to-one

Every element in the target is covered by **one or less elements** from the domain.

In formal definations

One-to-one

Every element in the target is covered by **one or less elements** from the domain.

Onto

Every element in the target is covered by **one or more elements** from the domain.

In formal definitions

One-to-one

Every element in the target is covered by **one or less elements** from the domain.

Onto

Every element in the target is covered by **one or more elements** from the domain.

Bijjective


Every element in the target is covered by **exactly one element** from the domain.

Function Properties Examples

$f: X \rightarrow A$

$X = \{ w, x, y, z \}$


$A = \{ a, b, c, d, e \}$



f is not one-to-one because
 $f(w) = f(z) = c$

f is not one-to-one because $f(w) = f(z) = c$.


Function Properties Examples

 $f: X \rightarrow A$ $X = \{w, x, y, z\}$ $A = \{a, b, c, d, e\}$ 

f is not onto because
there are no elements in X
that map to d or e

f is not onto because no elements in X map to d or e .

Function Properties Examples

 $f: X \rightarrow A$
 $X = \{w, x, y, z\}$
 $A = \{a, b, c, d, e\}$


Now f is one-to-one but not onto


Now f is one-to-one but not onto.

Function Properties Examples

$$f: X \rightarrow A$$

$$X = \{w, x, y, z\}$$


$$A = \{a, b, c, d\}$$



Now f is one-to-one and onto


Now f is one-to-one and onto. f is a bijection.

Relative sizes of the domain and target




onto

$$|D| \geq |T|$$



one-to-one

$$|D| \leq |T|$$



bijection

$$|D| \leq |T| \text{ and } |D| \geq |T|$$

$$|D| = |T|$$

Excercise 1

- Let f be a function whose domain is $\{0,1\}^3$ and whose target is $\{0,1\}^2$.

Ex.

- Is it possible that f is one-to-one?

Exercise 1

- Let f be a function whose domain is $\{0,1\}^3$ and whose target is $\{0,1\}^2$.

Ex.

- Is it possible that f is one-to-one?
 - No
- Is it possible that f is onto?

Exercise 1

- Let f be a function whose domain is $\{0,1\}^3$ and whose target is $\{0,1\}^2$.

Ex.

- Is it possible that f is one-to-one?
 - No
- Is it possible that f is onto?
 - Yes

Excercise 2

For each of the functions below, indicate whether the function is onto, one-to-one, neither or both.

Ex.

- $f : \{0,1\}^3 \rightarrow \{0,1\}^3$. The output of f is obtained by taking the input string and replacing the first bit by 1, regardless of whether the first bit is a 0 or 1. For example, $f(001) = 101$ and $f(110) = 110$

Exercise 2

For each of the functions below, indicate whether the function is onto, one-to-one, neither or both.

Ex.

- $f : \{0,1\}^3 \rightarrow \{0,1\}^3$. The output of f is obtained by taking the input string and replacing the first bit by 1, regardless of whether the first bit is a 0 or 1. For example, $f(001) = 101$ and $f(110) = 110$
 - Neither one-to-one nor onto.

Exercise 2

For each of the functions below, indicate whether the function is onto, one-to-one, neither or both.

Ex.

- $f : \{0,1\}^3 \rightarrow \{0,1\}^3$. The output of f is obtained by taking the input string and replacing the first bit by 1, regardless of whether the first bit is a 0 or 1. For example, $f(001) = 101$ and $f(110) = 110$
 - Neither one-to-one nor onto.
- $f : \{0,1\}^3 \rightarrow \{0,1\}^3$. The output of f is obtained by taking the input string and reversing the bits. For example $f(011) = 110$.

Exercise 2

For each of the functions below, indicate whether the function is onto, one-to-one, neither or both.

Ex.

- $f : \{0,1\}^3 \rightarrow \{0,1\}^3$. The output of f is obtained by taking the input string and replacing the first bit by 1, regardless of whether the first bit is a 0 or 1. For example, $f(001) = 101$ and $f(110) = 110$
 - Neither one-to-one nor onto.
- $f : \{0,1\}^3 \rightarrow \{0,1\}^3$. The output of f is obtained by taking the input string and reversing the bits. For example $f(011) = 110$.
 - One-to-one and onto.

Excercise 2

- For each of the functions below, indicate whether the function is onto, one-to-one, neither or both. If the function is not onto or not one-to-one, give an example showing why.

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$

Excercise 2

- For each of the functions below, indicate whether the function is onto, one-to-one, neither or both. If the function is not onto or not one-to-one, give an example showing why.

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$

Excercise 2

- For each of the functions below, indicate whether the function is onto, one-to-one, neither or both. If the function is not onto or not one-to-one, give an example showing why.

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$

Excercise 2

- For each of the functions below, indicate whether the function is onto, one-to-one, neither or both. If the function is not onto or not one-to-one, give an example showing why.

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$
 - Not onto.
 - One to one.

Excercise 2

- For each of the functions below, indicate whether the function is onto, one-to-one, neither or both. If the function is not onto or not one-to-one, give an example showing why.

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$
 - Not onto.
 - One to one.
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = x - 4$

Excercise 2

- For each of the functions below, indicate whether the function is onto, one-to-one, neither or both. If the function is not onto or not one-to-one, give an example showing why.

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$
 - Not onto.
 - One to one.
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = x - 4$
 - Onto.
 - One to one.
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = 5x - 4$

Excercise 2

- For each of the functions below, indicate whether the function is onto, one-to-one, neither or both. If the function is not onto or not one-to-one, give an example showing why.

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$
 - Not onto.
 - One to one.
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = x - 4$
 - Onto.
 - One to one.
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = 5x - 4$
 - One-to-one
 - Not onto.

Outline

- 1 Introduction
- 2 Floor and Ceiling
- 3 Function Properties
- 4 Function Inverse**
- 5 Composition of Functions

Function Inverse

- If a function $f: X \rightarrow Y$ is a **bijection**, then the inverse of f is obtained by exchanging the first and second entries in each pair in f .
- The inverse of f is denoted by f^{-1}

$$f^{-1} = \{ (y, x) : (x, y) \in f \}.$$

Example 1

$$X = \{1, 2, 3\}$$

$$Y = \{7, 8, 9\}$$

$$1 \bullet \xrightarrow{\text{blue}} \bullet 7$$

$$2 \bullet \xrightarrow{\text{blue}} \bullet 8$$

$$3 \bullet \xrightarrow{\text{blue}} \bullet 9$$

$$f: X \rightarrow Y$$

$$f = \{ (1, 7), (2, 9), (3, 9) \}$$

$$7 \bullet \xrightarrow{\text{red}} \bullet 1$$


$$\textcircled{8} \bullet \xrightarrow{\text{red}} \bullet 2$$

$$\textcircled{9} \bullet \xrightarrow{\text{red}} \bullet 3$$

$$f^{-1} = \{ (7, 1), (9, 2), (9, 3) \}$$

f^{-1} is not a function. f does not have an inverse.

Example 2



$$g: X \rightarrow Y$$

$$g = \{ (1, 9), (2, 7), (3, 8) \}$$

$$g^{-1}(7) = 2$$



$$g^{-1} = Y \rightarrow X$$

$$g^{-1} = \{ (7, 2), (8, 3), (9, 1) \}$$

g^{-1} is a function. g has an inverse defined by

$$g^{-1}(8) = 3$$

$$g^{-1}(9) = 1$$

Excercise 1

- For each of the following functions, indicate whether the function has a well-defined inverse. If the inverse is well-defined, give the input/output relationship of f^{-1} .

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$

Excercise 1

- For each of the following functions, indicate whether the function has a well-defined inverse. If the inverse is well-defined, give the input/output relationship of f^{-1} .

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
 - f^{-1} is not well defined
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$

Excercise 1

- For each of the following functions, indicate whether the function has a well-defined inverse. If the inverse is well-defined, give the input/output relationship of f^{-1} .

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
 - f^{-1} is not well defined
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
 - $f^{-1}(x) = \sqrt[3]{x}$
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$

Excercise 1

- For each of the following functions, indicate whether the function has a well-defined inverse. If the inverse is well-defined, give the input/output relationship of f^{-1} .

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
 - f^{-1} is not well defined
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
 - $f^{-1}(x) = \sqrt[3]{x}$
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$
 - Not onto.
 - One to one.
 - f^{-1} is not well defined

Excercise 1

- For each of the following functions, indicate whether the function has a well-defined inverse. If the inverse is well-defined, give the input/output relationship of f^{-1} .

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
 - f^{-1} is not well defined
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = x - 4$
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
 - $f^{-1}(x) = \sqrt[3]{x}$
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$
 - Not onto.
 - One to one.
 - f^{-1} is not well defined

Excercise 1

- For each of the following functions, indicate whether the function has a well-defined inverse. If the inverse is well-defined, give the input/output relationship of f^{-1} .

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
 - f^{-1} is not well defined
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
 - $f^{-1}(x) = \sqrt[3]{x}$
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$
 - Not onto.
 - One to one.
 - f^{-1} is not well defined
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = x - 4$
 - Onto.
 - One to one.
 - $f^{-1}(x) = x + 4$
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = 5x - 4$

Excercise 1

- For each of the following functions, indicate whether the function has a well-defined inverse. If the inverse is well-defined, give the input/output relationship of f^{-1} .

Ex.

- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^2$
 - Not onto.
 - Not one to one.
 - f^{-1} is not well defined
- $f: \mathbb{R} \rightarrow \mathbb{R}. f(x) = x^3$
 - One to one
 - Onto.
 - $f^{-1}(x) = \sqrt[3]{x}$
- $h: \mathbb{Z} \rightarrow \mathbb{Z}. h(x) = x^3$
 - Not onto.
 - One to one.
 - f^{-1} is not well defined
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = x - 4$
 - Onto.
 - One to one.
 - $f^{-1}(x) = x + 4$
- $f: \mathbb{Z} \rightarrow \mathbb{Z}. f(x) = 5x - 4$
 - One-to-one
 - Not onto.
 - f^{-1} is not well defined

Excercise 2

$f : \{0,1\}^3 \rightarrow \{0,1\}^3$. The output of is obtained by taking the input string and reversing the bits. For example, $f(011) = 110$

- Indicate whether f has a well-defined inverse and write f^{-1} if exists.

Excercise 2

$f : \{0,1\}^3 \rightarrow \{0,1\}^3$. The output of is obtained by taking the input string and reversing the bits. For example, $f(011) = 110$

- Indicate whether f has a well-defined inverse and write f^{-1} if exists.

Sol:

- f has a well-defined inverse.
- $f^{-1} = f$

Outline

- 1 Introduction
- 2 Floor and Ceiling
- 3 Function Properties
- 4 Function Inverse
- 5 Composition of Functions**

Composition of functions

Composition of functions

Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$.


The **composition of g with f** , denoted $g \circ f$, is the function $(g \circ f): X \rightarrow Z$, such that for all $x \in X$, $(g \circ f)(x) = g(f(x))$.

Example


$$X = \{1, 2, 3\}$$

$$Y = \{a, b, c\}$$


$$Z = \{7, 10, 12\}$$



$$f: X \rightarrow Y$$




$$g: Y \rightarrow Z$$




$$g \circ f: X \rightarrow Z$$

Excercise




- What is the domain of $g \circ f$?

Excercise




- What is the domain of $g \circ f$?
 - $X = \{v, w, x, y, z\}$

Exercise




- What is the domain of $g \circ f$?
 - $X = \{v, w, x, y, z\}$
- What is the target of $g \circ f$?

Exercise




- What is the domain of $g \circ f$?
 - $X = \{v, w, x, y, z\}$
- What is the target of $g \circ f$?
 - $S = \{1, 2, 3, 4, 5\}$

Exercise




- What is the domain of $g \circ f$?
 - $X = \{v, w, x, y, z\}$
- What is the target of $g \circ f$?
 - $S = \{1, 2, 3, 4, 5\}$
- Give the arrow diagram for $g \circ f$.


Exercise




- What is the domain of $g \circ f$?
 - $X = \{v, w, x, y, z\}$
- What is the target of $g \circ f$?
 - $S = \{1, 2, 3, 4, 5\}$
- Give the arrow diagram for $g \circ f$.



Exercise




- What is the domain of $g \circ f$?
 - $X = \{v, w, x, y, z\}$
- What is the target of $g \circ f$?
 - $S = \{1, 2, 3, 4, 5\}$
- Give the arrow diagram for $g \circ f$.




- What is the range of $g \circ f$?

Exercise



- What is the domain of $g \circ f$?
 - $X = \{v, w, x, y, z\}$
- What is the target of $g \circ f$?
 - $S = \{1, 2, 3, 4, 5\}$
- Give the arrow diagram for $g \circ f$.



- What is the range of $g \circ f$?
 - $\{1, 3, 4\}$

Notes I

- $f \circ g$ is not the same as $g \circ f$.

Ex.

$$\begin{aligned} f: R^+ &\rightarrow R^+, f(x) = x^3 \\ g: R^+ &\rightarrow R^+, g(x) = x + 2 \end{aligned}$$

- $(f \circ g)(x) = f(g(x)) = (x + 2)^3$
- $(g \circ f)(x) = g(f(x)) = x^3 + 2$

Notes II

- It is possible to compose more than two functions.
- Composition is associative.

$$f \circ g \circ h = (f \circ g) \circ h = f \circ (g \circ h) = f(g(h(x)))$$

Ex.

$$\begin{aligned} f: R^+ &\rightarrow R^+, f(x) = x^3 \\ g: R^+ &\rightarrow R^+, g(x) = x + 2 \\ h: R^+ &\rightarrow R^+, h(x) = x - 1 \end{aligned}$$

- $(f \circ g)(x) = f(g(x)) = (x + 2)^3$
- $(f \circ g \circ h)(x) = f(g(h(x))) = (x + 1)^3$

Identity Function

Identity Function




The identity function **always maps a set onto itself** and maps every element onto itself.

- The identity function on A , denoted $I_A : A \rightarrow A$, is defined as $I_A(a) = a$, for all $a \in A$.


Note That

Let $f: A \rightarrow B$ be a bijection. Then $f^{-1} \circ f = I_A$ and $f \circ f^{-1} = I_B$.

Example

 $X = \{1, 2, 3\}$
 $Y = \{r, s, t\}$


The identity function on X : I_X



The identity function on Y : I_Y

The composition of f with the inverse of f has domain Y and target Y and maps each element to itself and is therefore the identity function on Y .



Questions 

