# ECEN 227 - Introduction to Finite Automata and Discrete Mathematics

Dr. Mahmoud Nabil mnmahmoud@ncat.edu

North Carolina A & T State University

November 11, 2019

## Talk Overview

1 Introduction to binary relations

- Properties of binary relations
- 3 Directed graphs, paths, and cycles

## Outline

- 1 Introduction to binary relations
- Properties of binary relations
- 3 Directed graphs, paths, and cycles

## Relation

#### Relation

A binary relation between two sets A and B is a subset R of  $A \times B$ .

#### Ex.

- S is the set of students at a university and C is the set of classes offered by the university.
- The relation E between S and C indicates whether a student is enrolled in a given class.
- Usually we can denote this relation as sEc.

## Relations and Function


Recall that functions have more restrictions on the connection between the domain and the target as follows.

- Each element in the domain should point to one and only one element in the target.
- This is not the case in the relations

# Arrow diagram for a relation

```
People = { Sue, Harry, Sam }
Files = { FileA, FileB, FileC, FileD }
```

Relation A: pAf if person p has access to file f



```
A = { (Sue, FileB) , (Sue, FileC) , (Sue, FileD) , (Harry, FileA) , (Harry, FileD) }
```

# Matrix representation for a relation

```
People = { Sue, Harry, Sam }
Files = { File A, File B, File C, File D }
```


Relation A: pAf if person p has access to file f

```
A = { (Sue, File B) (Sue, File C) (Sue, File D) (Harry, File A) (Harry, File D) }
```

# Binary Relation on a Set

- We can have a binary relation between a set A and itself.
- In this case we call it a binary relation on the set A.
- The result is a subset of A x A.
- The set A is called the domain of the binary relation.

#### Ex.




```
A = { a, b, c, d, e }

R ⊆ A x A

R = { (a, b) (b, c) (e, c) (c, e) (d, a) (d, d) }
```

Draw the arrow diagram and the matrix representation for the following relation. Define the set  $A = \{r, o, t, p, c\}$  and  $B = \{discrete, math, proof, proposition\}$ . Define the relation  $R \subseteq A \times B$  such that (letter, word) is in the relation if that letter occurs somewhere in the word.


Draw the arrow diagram and the matrix representation for the following relation. Define the set  $A = \{r, o, t, p, c\}$  and  $B = \{discrete, math, proof, proposition\}$ . Define the relation  $R \subseteq A \times B$  such that (letter, word) is in the relation if that letter occurs somewhere in the word.





Draw the arrow diagram for the following relation. The domain of relation D is  $\{2, 3, 12, 16, 27, 48\}$ . For x, y in the domain, xDy if y is an integer multiple of x.

Draw the arrow diagram for the following relation. The domain of relation D is  $\{2, 3, 12, 16, 27, 48\}$ . For x, y in the domain, xDy if y is an integer multiple of x.



## Outline


Introduction to binary relations

- Properties of binary relations
- 3 Directed graphs, paths, and cycles

Binary relation R can be characterized by six properties. The properties are defined and illustrated using arrow diagrams.

- The relation R can be either reflexive or anti-reflexive or neither.
- The relation R can be either symmetric or anti-symmetric or neither.
- The relation R can be either transitive or not transitive.


## Reflexive Relation



A = { a, b, c, d, e }

Relation R is
reflexive
if for all x ∈ A
x R x
a R a, b R b, c R c,
d R d, and e R e


## Anti-Reflexive Relation



$$A = \{ a, b, c, d, e \}$$

Relation R is anti-reflexive if for all x ∈ A it is not true that x R x

# Symmetric Relation




$$A = \{ a, b, c, d, e \}$$

Relation R on A is symmetric if for all x, y ∈ A x R y ↔ y R x

 $x R y \leftrightarrow y R x$ 

# Anti-Symmetric Relation




 $A = \{ a, b, c, d, e \}$ 

Relation R is anti-symmetric if for all x, y ∈ A x R y and y R x → x = y

Note: there is no



## Transitive Relation




```
A = \{ a, b, c, d, e \}
```

Relation R on A is transitive if for all x, y,  $z \in A$ if x R y and y R z, then x R z

eRa and aRb  $\longrightarrow$  eRb eRb and bRc  $\longrightarrow$  eRc eRa and aRc  $\longrightarrow$  eRc aRb and bRc  $\longrightarrow$  aRc


Given the below relation indicate whether the relation is:

- reflexive, anti-reflexive, or neither
- symmetric, anti-symmetric, or neither
- transitive or not transitive



Given the below relation indicate whether the relation is:

- reflexive, anti-reflexive, or neither
- symmetric, anti-symmetric, or neither
- transitive or not transitive



#### Answer.

- neither reflexive nor anti-reflexive.
- neither symmetric nor anti-symmetric.
- not transitive.



Given the below relation indicate whether the relation is:

- reflexive, anti-reflexive, or neither
- symmetric, anti-symmetric, or neither
- transitive or not transitive

The domain of the relation L is the set of all real numbers. For  $x, y \in R$ , xLy if x < y.

Given the below relation indicate whether the relation is:

- reflexive, anti-reflexive, or neither
- symmetric, anti-symmetric, or neither
- transitive or not transitive

The domain of the relation L is the set of all real numbers. For x,  $y \in R$ , xLy if x < y.

#### Answer.

- anti-reflexive: For any real number x, it is always false that x < x.
- anti-symmetric: For any two real numbers x and y, it can not be true that x < y and y < x.</li>
- transitive: If x < y and y < z, then x < z.

Given the below relation indicate whether the relation is:

- reflexive, anti-reflexive, or neither
- symmetric, anti-symmetric, or neither
- transitive or not transitive

The domain of the relation L is the set of all real numbers. For  $x, y \in R$ , xLy if  $x \le y$ .

Given the below relation indicate whether the relation is:

- reflexive, anti-reflexive, or neither
- symmetric, anti-symmetric, or neither
- transitive or not transitive

The domain of the relation L is the set of all real numbers. For  $x, y \in R$ , xLy if  $x \le y$ .

#### Answer.

- reflexive: For any real number x, it is always true that  $x \le x$ .
- anti-symmetric: For any two real numbers x and y, if  $x \le y$  and  $y \le x$ , then x = y.
- transitive: If  $x \le y$  and  $y \le z$ , then  $x \le z$ .

## Outline

Introduction to binary relations

- Properties of binary relations
- 3 Directed graphs, paths, and cycles

# Graph

Graph is simply a relation over set. It is used widely in computer science topics.

#### Ex.

- Internet pages.
- Friends on facebook.
- Train/Bus stations.
- Communication network.
- etc.

# Directed Graph (digraph)

#### Digraph


A directed graph (or digraph, for short) consists of a pair (V, E). V is a set of vertices, and E, a set of directed edges.

#### Vertex


An individual element of V is called a vertex.

#### Edge

A connection between two vertices.



# Graph Example



#### In-degree

The in-degree of a vertex is the number of edges pointing into it.

#### Out-degree

The out-degree of a vertex is the number of edges pointing out of it.

# Walks and directed Graph

A walk in a directed graph G is a sequence of alternating vertices and edges that starts and ends with a vertex.

$$< v_0, v_1, v_2, ..., v_n >$$

#### Walk length

The length of a walk is I, the number of edges in the walk.

#### Open walk

An open walk is a walk in which the first and last vertices are not the same.

#### Closed walk

A closed walk is a walk in which the first and last vertices are the same.

# Example



#### Walk:

$$\langle a, b, c, b, d \rangle$$
 walk length = 4

The walk is open because the first and last vertices are not the same

#### Walk:



walk length = 5 Edge (c, b) occurs twice so (c, b) is counted twice

This walk closed because the first and last vertices are the same.

Closed walk of length 1: (d, d)

Closed walk of length 0: ( a )

#### Defination

#### Trail

A trail is an open walk in which no edge occurs more than once.

#### Path

A path is a trail in which no vertex occurs more than once.


#### Circuit

A circuit is a closed walk in which no edge occurs more than once.

## Cycle

A cycle is a circuit in which no vertex occurs more than once, except the first and last vertices which are the same.

## Example




#### Walk:

(a, b, c, b, d) No edge occurs more than once. So this open walk is a trail.

b is reached twice So this trail is not a path

\( \begin{align\*} \begin{align\*}

The circuit is a cycle because only the first and last vertices are repeated.



- What is the in-degree of vertex d?
- What is the out-degree of vertex c?
- What is the head of edge (b, c)?
- What is the tail of edge (g, f)?
- List all the self-loops in the graph.
- Is <a, g, f, c, d> a walk in the graph? Is it a trail? Is it a path?
- Is <a, g, f, d, b> a walk in the graph? Is it a trail? Is it a path?
- Is <c, g, f, e> a circuit in the graph? Is it a cycle?



Questions &

