

A Generalized Binomial Framework for Moment Transformations Across Arbitrary Origins

Abstract

Statistical moments are fundamental descriptors of data and probability distributions. Transforming moments across origins traditionally relies on fragmented, order-specific algebraic expansions. We establish a unified, generalized binomial framework for transforming moments of any order $r \in \mathbb{N}$ across arbitrary initial and target origins $a \rightarrow k$. Any origin shift $b = a - k$ admits a symbolic binomial expansion, $\mu'_r(k) = (a + b)^r$, where powers a^j map to the j -th moment $\mu'_j(a)$. This unifies raw-to-raw, raw-to-central, and central-to-raw transformations under a single operational principle. We contrast our $O(r^2)$ sample-size-independent transformation with parallel algorithms (e.g., Pébay, 2008) that address dataset merging across processors. Validation on theoretical (Binomial, Normal) and empirical distributions confirms numerical exactness. An R package `convmoment` provides `conv_moment()`, `conv_moment_all()`, `raw2central()`, and `central2raw()`. Benchmarks show a $> 250\times$ speedup over raw data recomputation for $n = 10^5$.

Keywords: moment transformation, binomial theorem, central moments, raw moments, statistical computing, pedagogical statistics

1 Introduction and Literature Review

1.1 Background and Practical Importance

Moments play a central role across mathematical statistics, econometrics, signal processing, and data science. Raw moments, defined relative to an arbitrary origin a as $\mu'_r(a) = \mathbb{E}[(X - a)^r]$, and central moments, defined relative to the mean μ as $\mu_r = \mathbb{E}[(X - \mu)^r]$, provide fundamental structural information regarding location, scale, skewness, and kurtosis.

In applied and theoretical statistics, transforming moments across different reference origins is essential in several key domains:

1. **Edgeworth and Gram-Charlier Series Expansions:** Approximating arbitrary or nearly Gaussian probability density functions requires converting higher-order central moments into standardized moments or cumulants (Hall 1992; Blinnikov and Moessner 1998).
2. **Generalized Method of Moments (GMM):** In econometrics and financial modeling, parameter estimation frequently involves matching empirical sample moments computed around arbitrary origins to theoretical population moments (Hansen 1982).
3. **Image Processing and Pattern Recognition:** In computer vision, image moment invariants (Hu 1962; Mukundan and Ramakrishnan 1998) rely on shifting spatial origins to geometric centroids to achieve translation and scale invariance.

1.2 Review of Prior Art and Related Work

Despite the widespread application of moments, existing literature presents moment transformations in a fragmented and computationally inefficient manner:

- **Classroom and Textbook Literature:** Standard statistical textbooks (Cramér 1946; Mood et al. 1974; Kendall and Stuart 1977; Johnson et al. 1994) present separate, hardcoded algebraic formulas for converting raw moments about zero to central moments ($\mu_2 = \mu'_2 - (\mu'_1)^2$, $\mu_3 = \mu'_3 - 3\mu'_2\mu'_1 + 2(\mu'_1)^3$, etc.). These treatments lack a unified operational framework and force students and practitioners to memorize order-specific formulas or consult printed tables.
- **Parallel and Online Moment Algorithms:** A distinct line of research focuses on online and parallel statistical updates (Chan et al. 1983; Terriberry 2007; Pébay 2008; Pébay et al. 2016). These algorithms address the problem of merging two separate data subsets A and B across parallel computing nodes or streaming buffers. While Pébay (2008) provides formulas for pairwise moment aggregation, those methods solve dataset partitioning across multiple processors rather than the transformation of a single pre-computed set of moments or theoretical distribution parameters across arbitrary reference points $a \rightarrow k$.
- **Symbolic Computational Methods:** Advanced computer algebra systems use symbolic cumulant expansions and umbral calculus (Andrews and Stafford 2000; Di Nardo et al. 2008). While mathematically elegant, these techniques require complex symbolic software engines and are rarely accessible for routine statistical computations or classroom instruction.

1.3 Contributions and Novelty

To bridge the gap between elementary textbook formulas and complex parallel algorithms, this paper introduces a **unified, generalized binomial framework** for arbitrary moment transformations. The primary contributions of this work are:

1. **A Single Unified Operator:** We prove that any origin shift from a to k is completely characterized by the shift parameter $b = a - k$, unifying raw-to-raw, raw-to-central, and central-to-raw conversions under a single equation.
2. **The Symbolic Binomial Mnemonic:** We introduce a symbolic operator substitution $(a + b)^r$ with $a^j \mapsto \mu'_j(a)$, enabling practitioners to construct conversion formulas of any order $r \in \mathbb{N}$ instantly using Pascal's triangle.
3. **Sample-Size Independence:** We demonstrate that transforming pre-computed moments requires $O(r^2)$ operations independent of sample size n , offering a $> 250\times$ speedup over recomputing sample moments directly from large datasets ($n \geq 10^5$).
4. **Open-Source R Package:** We provide an R implementation (`convmoment`) featuring verified functions (`conv_moment()`, `conv_moment_all()`, `raw2central()`, `central2raw()`).

2 Traditional Approach to Moment Transformations

2.1 Raw Moments Between Arbitrary Origins

Consider transforming raw moments from an initial origin a to a target origin k . Traditionally, this requires expressing $(x_i - k)$ as $[(x_i - a) + (a - k)]$, expanding the polynomial for each specific moment order, and distributing the summation operator over sample size n .

2.1.1 Numerical Example

Suppose the first three moments of a dataset about origin $a = 2$ are known to be:

$$\mu'_1(2) = -1, \quad \mu'_2(2) = 7, \quad \text{and} \quad \mu'_3(2) = 39.$$

We wish to find the corresponding first three moments about a new origin $k = 5$.

2.1.1.1 1. First Moment Transformation ($r = 1$)

$$\begin{aligned}\mu'_1(5) &= \frac{1}{n} \sum_{i=1}^n (x_i - 5) = \frac{1}{n} \sum_{i=1}^n [(x_i - 2) - 3] \\ &= \frac{1}{n} \sum_{i=1}^n (x_i - 2) - 3 = \mu'_1(2) - 3 = -1 - 3 = -4.\end{aligned}$$

2.1.1.2 2. Second Moment Transformation ($r = 2$)

$$\begin{aligned}\mu'_2(5) &= \frac{1}{n} \sum_{i=1}^n (x_i - 5)^2 = \frac{1}{n} \sum_{i=1}^n [(x_i - 2) - 3]^2 \\ &= \frac{1}{n} \sum_{i=1}^n [(x_i - 2)^2 - 6(x_i - 2) + 9] \\ &= \mu'_2(2) - 6\mu'_1(2) + 9 = 7 - 6(-1) + 9 = 22.\end{aligned}$$

2.1.1.3 3. Third Moment Transformation ($r = 3$)

$$\begin{aligned}\mu'_3(5) &= \frac{1}{n} \sum_{i=1}^n (x_i - 5)^3 = \frac{1}{n} \sum_{i=1}^n [(x_i - 2) - 3]^3 \\ &= \frac{1}{n} \sum_{i=1}^n [(x_i - 2)^3 - 9(x_i - 2)^2 + 27(x_i - 2) - 27] \\ &= \mu'_3(2) - 9\mu'_2(2) + 27\mu'_1(2) - 27 \\ &= 39 - 9(7) + 27(-1) - 27 = -78.\end{aligned}$$

While straightforward for low orders, deriving transformations for order $r \geq 4$ using this approach requires expanding higher-degree polynomials manually, increasing the likelihood of algebraic errors.

2.2 Traditional Raw-to-Central Transformations

When converting raw moments about an arbitrary origin a to central moments μ_r (where target origin $k = \bar{x} = \mu'_1(a) + a$), standard textbooks prescribe fixed formulas:

$$\mu_1 = 0$$

$$\mu_2 = \mu'_2(a) - [\mu'_1(a)]^2$$

$$\mu_3 = \mu'_3(a) - 3\mu'_2(a)\mu'_1(a) + 2[\mu'_1(a)]^3$$

$$\mu_4 = \mu'_4(a) - 4\mu'_3(a)\mu'_1(a) + 6\mu'_2(a)[\mu'_1(a)]^2 - 3[\mu'_1(a)]^4$$

$$\mu_5 = \mu'_5(a) - 5\mu'_4(a)\mu'_1(a) + 10\mu'_3(a)[\mu'_1(a)]^2 - 10\mu'_2(a)[\mu'_1(a)]^3 + 4[\mu'_1(a)]^5.$$

These formulas lack a structural pattern that students or practitioners can easily reconstruct without explicit memorization.

3 The Proposed Generalized Binomial Method

3.1 Theoretical Derivation

Let a denote the initial origin and k denote the target origin. Define the **origin shift** b as:

$$b = a - k.$$

For any random variable X , the deviation from target origin k can be decomposed as:

$$X - k = (X - a) + (a - k) = (X - a) + b.$$

Raising both sides to the power of $r \in \mathbb{N}$ and applying the binomial theorem yields:

$$(X - k)^r = \sum_{j=0}^r \binom{r}{j} (X - a)^j b^{r-j}.$$

Taking the expectation $\mathbb{E}[\cdot]$ on both sides:

$$\mathbb{E}[(X - k)^r] = \sum_{j=0}^r \binom{r}{j} \mathbb{E}[(X - a)^j] b^{r-j}.$$

Since $\mathbb{E}[(X - a)^j] = \mu'_j(a)$ and defining $\mu'_0(a) \equiv 1$, we obtain the **Generalized Binomial Moment Transformation Theorem**:

Theorem 1 (Generalized Binomial Transformation).

For any initial origin a , target origin k , and origin shift $b = a - k$, the r^{th} moment about k is given by:

$$\mu'_r(k) = \sum_{j=0}^r \binom{r}{j} \mu'_j(a) b^{r-j}.$$

3.2 The Symbolic Binomial Operator and Pascal's Triangle

A powerful mnemonic representation of Theorem 1 is obtained by introducing a symbolic operator substitution:

$$\mu'_r(k) \equiv (a + b)^r \quad \text{with the substitution } a^j \mapsto \mu'_j(a) \text{ and } a^0 \equiv 1.$$

Expanding $(a + b)^r$ using standard binomial coefficients directly yields the required conversion formula:

The numerical coefficients are directly read from **Pascal's Triangle**:

$$\begin{array}{ccccccc} & & & & 1 & & & \\ & & & & & & & \\ & & & 1 & & 1 & & \\ & & 1 & & 2 & & 1 & \\ & 1 & & 3 & & 3 & & 1 \\ & & 1 & & 4 & & 6 & & 4 & & 1 \\ & 1 & & 5 & & 10 & & 10 & & 5 & & 1 \end{array}$$

Moment Order (r)	Symbolic Binomial Form	Explicit Moment Formula ($\mu'_r(k)$)
1	$(a + b)^1$	$\mu'_1(a) + b$
2	$(a + b)^2$	$\mu'_2(a) + 2\mu'_1(a)b + b^2$
3	$(a + b)^3$	$\mu'_3(a) + 3\mu'_2(a)b + 3\mu'_1(a)b^2 + b^3$
4	$(a + b)^4$	$\mu'_4(a) + 4\mu'_3(a)b + 6\mu'_2(a)b^2 + 4\mu'_1(a)b^3 + b^4$
5	$(a + b)^5$	$\mu'_5(a) + 5\mu'_4(a)b + 10\mu'_3(a)b^2 + 10\mu'_2(a)b^3 + 5\mu'_1(a)b^4 + b^5$

Table 1: Symbolic binomial expansion for moment orders 1–5.

3.3 Matrix Formulation

For computational implementations, Theorem 1 can be expressed concisely as a matrix-vector product. Let $\boldsymbol{\mu}'(a) = [1, \mu'_1(a), \mu'_2(a), \dots, \mu'_K(a)]^T$ be the column vector of moments up to order K . Then:

$$\boldsymbol{\mu}'(k) = \mathbf{C}_b \boldsymbol{\mu}'(a),$$

where $\mathbf{C}_b$ is a lower-triangular transformation matrix of dimension $(K + 1) \times (K + 1)$ with entries:

$$[\mathbf{C}_b]_{r,j} = \begin{cases} \binom{r}{j} b^{r-j} & \text{for } 0 \leq j \leq r \leq K, \\ 0 & \text{otherwise.} \end{cases}$$

3.4 Unified Transformation Paradigm

The generalized origin shift $b = a - k$ encapsulates all common moment conversions:

1. **Raw-to-Raw Conversion:** $b = a - k$, where a and k are arbitrary initial and final origins.
2. **Raw-to-Central Conversion:** Initial origin is a (often $a = 0$). The target origin is the mean $k = \bar{x} = \mu'_1(a) + a$. Hence, $b = a - (\mu'_1(a) + a) = -\mu'_1(a)$.

3. **Central-to-Raw Conversion:** Initial origin is the mean $a = \bar{x}$. The target origin is $k = 0$. Hence, $b = \bar{x} - 0 = \mu'_1(0)$.

3.5 Re-Evaluating the Numerical Example

Returning to our previous numerical example where $a = 2, k = 5, \mu'_1(2) = -1, \mu'_2(2) = 7, \mu'_3(2) = 39$:

$$b = a - k = 2 - 5 = -3.$$

Applying the symbolic expansion:

1. **First Moment:** $\mu'_1(5) = \mu'_1(2) + b = -1 + (-3) = -4$.
2. **Second Moment:** $\mu'_2(5) = \mu'_2(2) + 2\mu'_1(2)b + b^2 = 7 + 2(-1)(-3) + (-3)^2 = 7 + 6 + 9 = 22$.
3. **Third Moment:** $\mu'_3(5) = \mu'_3(2) + 3\mu'_2(2)b + 3\mu'_1(2)b^2 + b^3 = 39 + 3(7)(-3) + 3(-1)(-3)^2 + (-3)^3 = 39 - 63 - 27 - 27 = -78$.

The calculations are completed in seconds without manual algebraic expansion.

4 Moment Transformations in Named Probability Distributions

The generalized binomial method applies equally to theoretical probability distributions.

4.1 Discrete Distribution: Binomial Distribution $X \sim \text{Bin}(n, p)$

4.1.1 Traditional MGF Approach

The moment generating function of $X \sim \text{Bin}(n, p)$ is $M_X(t) = (1 - p + pe^t)^n$. Differentiating $M_X(t)$ at $t = 0$ yields:

$$\mu'_1(0) = np, \quad \text{and} \quad \mu'_2(0) = np(1 - p) + n^2p^2.$$

To evaluate central moments using MGFs, one must differentiate $M_{X-\mu}(t) = e^{-npt}(1 - p + pe^t)^n$, requiring nested applications of the product rule:

$$\left. \frac{d}{dt} M_{X-\mu}(t) \right|_{t=0} = (-np)(1) + (1)(np) = 0.$$

4.1.2 Generalized Binomial Approach

Using our framework:

- Initial origin $a = 0$.
- Target origin $k = \mu = np$.
- Origin shift $b = a - k = 0 - np = -np$.

1. **First Central Moment (μ_1):** $\mu_1 = \mu'_1(0) + b = np + (-np) = 0$.

2. **Second Central Moment (μ_2 - Variance):**

$$\begin{aligned} \mu_2 &= \mu'_2(0) + 2\mu'_1(0)b + b^2 \\ &= [np(1 - p) + n^2p^2] + 2(np)(-np) + (-np)^2 \\ &= np - np^2 + n^2p^2 - 2n^2p^2 + n^2p^2 = np(1 - p). \end{aligned}$$

3. **Third Central Moment (μ_3):** Given $\mu'_3(0) = np - 3np^2 + 3n^2p^2 + 2np^3 - 3n^2p^3 + n^3p^3$:

$$\begin{aligned}
\mu_3 &= \mu'_3(0) + 3\mu'_2(0)b + 3\mu'_1(0)b^2 + b^3 \\
&= (np - 3np^2 + 3n^2p^2 + 2np^3 - 3n^2p^3 + n^3p^3) \\
&\quad + 3(np - np^2 + n^2p^2)(-np) + 3(np)(-np)^2 + (-np)^3 \\
&= np(1-p)(1-2p).
\end{aligned}$$

4.2 Continuous Distribution: Normal Distribution $X \sim \mathcal{N}(\mu, \sigma^2)$

The raw moments about zero and central moments of the Normal distribution are:

Order (r)	Raw Moment Notation	Raw Moment Value ($\mu'_r(0)$)	Central Moment Notation	Central Moment Value
1	μ'_1	μ	μ_1	0
2	μ'_2	$\mu^2 + \sigma^2$	μ_2	σ^2
3	μ'_3	$\mu^3 + 3\mu\sigma^2$	μ_3	0
4	μ'_4	$\mu^4 + 6\mu^2\sigma^2 + 3\sigma^4$	μ_4	$3\sigma^4$

Table 2: Raw and central moments of the Normal distribution.

4.2.1 Raw-to-Central Derivation ($b = 0 - \mu = -\mu$)

1. **First Central Moment:** $\mu_1 = \mu'_1 + b = \mu - \mu = 0$.

2. **Second Central Moment:**

$$\mu_2 = \mu'_2 + 2\mu'_1b + b^2 = (\mu^2 + \sigma^2) + 2\mu(-\mu) + (-\mu)^2 = \sigma^2.$$

3. Third Central Moment:

$$\begin{aligned}\mu_3 &= \mu'_3 + 3\mu'_2b + 3\mu'_1b^2 + b^3 \\ &= (\mu^3 + 3\mu\sigma^2) + 3(\mu^2 + \sigma^2)(-\mu) + 3\mu(-\mu)^2 + (-\mu)^3 \\ &= \mu^3 + 3\mu\sigma^2 - 3\mu^3 - 3\mu\sigma^2 + 3\mu^3 - \mu^3 = 0.\end{aligned}$$

4. Fourth Central Moment:

$$\begin{aligned}\mu_4 &= \mu'_4 + 4\mu'_3b + 6\mu'_2b^2 + 4\mu'_1b^3 + b^4 \\ &= (\mu^4 + 6\mu^2\sigma^2 + 3\sigma^4) + 4(\mu^3 + 3\mu\sigma^2)(-\mu) + 6(\mu^2 + \sigma^2)(-\mu)^2 + 4\mu(-\mu)^3 + (-\mu)^4 \\ &= (\mu^4 + 6\mu^2\sigma^2 + 3\sigma^4) - (4\mu^4 + 12\mu^2\sigma^2) + (6\mu^4 + 6\mu^2\sigma^2) - 4\mu^4 + \mu^4 \\ &= 3\sigma^4.\end{aligned}$$

4.2.2 Central-to-Raw Derivation ($b = \mu - 0 = \mu$)

Converting central moments back to raw moments about zero ($a = \mu, k = 0, b = \mu$):

$$\begin{aligned}\mu'_1(0) &= \mu_1 + b = 0 + \mu = \mu, \\ \mu'_2(0) &= \mu_2 + 2\mu_1b + b^2 = \sigma^2 + 0 + \mu^2 = \mu^2 + \sigma^2, \\ \mu'_3(0) &= \mu_3 + 3\mu_2b + 3\mu_1b^2 + b^3 = 0 + 3\sigma^2\mu + 0 + \mu^3 = \mu^3 + 3\mu\sigma^2, \\ \mu'_4(0) &= \mu_4 + 4\mu_3b + 6\mu_2b^2 + 4\mu_1b^3 + b^4 = 3\sigma^4 + 0 + 6\sigma^2\mu^2 + 0 + \mu^4 = \mu^4 + 6\mu^2\sigma^2 + 3\sigma^4.\end{aligned}$$

5 Computational Implementation in R and Benchmarking

5.1 R Package Implementation (`convmoment`)

We implement the generalized binomial moment conversion framework in R (Wickham 2015). The function `conv_moment()` computes the converted moment of order r , while `conv_moment_all()` converts a vector of moments of orders $1 \dots K$ across any initial origin a and target origin k .

```
conv_moment <- function(x, a, k, r) {  
  b <- a - k  
  mu_from <- c(1, x[1:r]) # mu_0 = 1, followed by moments 1..r  
  coeff <- choose(r, 0:r)  
  power <- r:0  
  sum(mu_from * coeff * (b^power))  
}
```

```
conv_moment_all <- function(x, a, k) {  
  K <- length(x)  
  sapply(1:K, function(r) conv_moment(x, a, k, r))  
}
```

```
raw2central <- function(raw_moments, origin = 0) {  
  mean_val <- raw_moments[1] + origin  
  conv_moment_all(raw_moments, a = origin, k = mean_val)
```

```
}
```

```
central2raw <- function(central_moments, mean_val) {  
  conv_moment_all(central_moments, a = mean_val, k = 0)  
}
```

5.2 Empirical Validation on Sample Data

We validate the package functions using sample data $X = (20, 25, 29, 32, 40)$.

1. Compute direct raw moments about 0 (orders 1 to 5).
2. Compute direct central moments about mean.
3. Convert raw to central using `conv_moment_all`.
4. Convert central back to raw about 0.

The absolute differences between direct sample moments and converted moments are on the order of 10^{-14} to 10^{-10} , confirming numerical exactness.

5.3 Computational Performance Benchmarking

A major practical advantage of the generalized binomial transformation is **sample-size independence**. When working with large datasets ($n \gg 10^4$), recomputing moments directly from raw data following an origin shift requires $O(nr)$ operations across all n observations. In contrast, transforming pre-computed moments using `conv_moment_all()` requires only $O(r^2)$ operations, completely independent of sample size n .

We benchmark 1,000 origin-shift operations on a simulated dataset of size $n = 100,000$:

- Pre-computed moments about origin a .
- Benchmark 1: Recompute moments from raw sample data ($O(nr)$).
- Benchmark 2: Generalized Binomial Transformation ($O(r^2)$).

Transforming pre-computed moments using `conv_moment_all()` achieves a $> 250\times$ **speedup** compared to recomputing moments over raw sample data.

Furthermore, while hardcoded scalar expressions for low orders (e.g. $r \leq 5$) are static macros, the generalized binomial algorithm scales dynamically to arbitrary high-order moments ($r = 10, 20, 50+$) without requiring manual algebraic expansion.

A demonstration for 10th order moment conversion confirms accuracy to machine precision.

6 Discussion

6.1 Unification and Pedagogical Advantages

The central contribution of this paper is the unification of moment transformation theory into a single operational framework. By expressing any origin shift $a \rightarrow k$ via the single parameter $b = a - k$, the proposed binomial expansion $\mu'_r(k) = (a + b)^r$ eliminates the fragmentation present in traditional statistical textbooks (Cramér 1946; Mood et al. 1974; Kendall and Stuart 1977).

From a pedagogical perspective, students and researchers no longer need to consult printed lookup tables or memorize order-specific algebraic formulas for raw-to-central (μ_r) or central-to-raw (μ'_r) conversions. Instead, any higher-order moment transformation can be constructed on demand using Pascal’s triangle coefficients.

6.2 Comparison with Parallel and Online Algorithms

It is instructive to contrast our framework with parallel and online moment update algorithms (Chan et al. 1983; Terriberry 2007; Pébay 2008; Pébay et al. 2016). Parallel algorithms are designed for distributed computing environments, where a dataset X is partitioned into subsets A and B , and moment updating rules are applied to merge statistics across processors without passing raw data.

In contrast, our framework addresses the complementary problem of transforming moments across reference origins for a **single sample or theoretical distribution**. While parallel algorithms modify sample size ($n_A + n_B$) and combine local means, our binomial transformation operates on fixed pre-computed moments, achieving an $O(r^2)$ computational complexity that is entirely independent of sample size n .

6.3 Numerical Stability Considerations and Limitations

When converting uncentered raw moments of very high order ($r > 20$) for empirical datasets with large location offsets, binomial expansions can be susceptible to floating-point catastrophic cancellation due to alternating sum signs (Higham 2002). For ultra-high-order empirical transformations ($r \geq 30$), combining the binomial transformation with arbitrary-precision arithmetic libraries or log-space moment representations (Di Nardo et al. 2008) provides numerical safeguards.

7 Conclusion and Future Work

In this paper, we have presented a generalized binomial framework for transforming statistical moments of any order $r \in \mathbb{N}$ across arbitrary origins $a \rightarrow k$. By introducing the origin

shift parameter $b = a - k$ and the symbolic operator representation $\mu'_r(k) \equiv (a + b)^r$, we established a unified method that encompasses raw-to-raw, raw-to-central, and central-to-raw conversions under a single mathematical principle.

We validated the framework theoretically across discrete (Binomial) and continuous (Normal) probability distributions and presented an open-source R package implementation (`convmoment`). Computational benchmarks demonstrate that transforming pre-computed moments using our algorithm achieves a $> 250\times$ speedup over raw data recomputation on large datasets ($n = 100,000$) while remaining numerically exact.

Future extensions of this work include generalizing the binomial shift framework to **multi-variate product moments** (Bishop et al. 1975), joint cross-moments, and matrix-valued spatial moment invariants in higher-dimensional computer vision applications. Diagnostic visualization via moment ratio diagrams (Pearson 2020) offers another natural direction for exploring transformed moments.

8 Disclosure Statement

The author has no conflicts of interest to declare.

9 Data Availability Statement

The R package `convmoment` and all replication code are publicly available at <https://github.com/mahmudstat/convmoment>. The package vignette is available at <https://www.thinkermahmud.com/convmoment/index.html>.

10 Supplementary Material

The following supplementary materials are available online:

1. **R Package `convmoment`:** Complete source code, documentation, and vignette demonstrating all package functions. Available at <https://github.com/mahmudstat/convmoment>.
2. **Replication Code:** Full R script reproducing all numerical examples, distribution validations (Binomial, Normal), empirical validation, and computational benchmarks presented in the paper.
3. **Vignette:** A comprehensive tutorial vignette showing basic usage, theoretical distribution examples, and matrix formulation. Available at <https://www.thinkermahmud.com/convmoment/index.html>.

References

- Andrews, D. F. and Stafford, J. E. (2000). *Symbolic Computation for Statistical Inference*. Oxford University Press, Oxford.
- Bishop, Y. M. M., Fienberg, S. E., and Holland, P. W. (1975). *Discrete Multivariate Analyses: Theory and Practice*. MIT Press, Cambridge, MA.
- Blinnikov, S. and Moessner, R. (1998). Expansions for nearly gaussian distributions. *Astronomy and Astrophysics Supplement Series*, 130(1):193–205.
- Chan, T. F., Golub, G. H., and LeVeque, R. J. (1983). Algorithms for computing the sample variance: Analysis and recommendations. *The American Statistician*, 37(3):242–247.

- Cramér, H. (1946). *Mathematical Methods of Statistics*. Princeton University Press, Princeton, NJ.
- Di Nardo, E., Guarino, G., and Senato, D. (2008). A unified framework for tenable statistical cumulants via symbolic computations. *Journal of Symbolic Computation*, 43(5):355–373.
- Hall, P. (1992). *The Bootstrap and Edgeworth Expansion*. Springer-Verlag, New York.
- Hansen, L. P. (1982). Large sample properties of generalized method of moments estimators. *Econometrica*, 50(4):1029–1054.
- Higham, N. J. (2002). *Accuracy and Stability of Numerical Algorithms*. SIAM, Philadelphia, 2nd edition.
- Hu, M.-K. (1962). Visual pattern recognition by moment invariants. *IRE Transactions on Information Theory*, 8(2):179–187.
- Johnson, N. L., Kotz, S., and Balakrishnan, N. (1994). *Continuous Univariate Distributions, Volume 1*. John Wiley & Sons, New York, 2nd edition.
- Kendall, M. G. and Stuart, A. (1977). *The Advanced Theory of Statistics: Vol. 1. Distribution Theory*. Charles Griffin & Company, London, 4th edition.
- Mood, A. M., Graybill, F. A., and Boes, D. C. (1974). *Introduction to the Theory of Statistics*. McGraw-Hill, New York, 3rd edition.
- Mukundan, R. and Ramakrishnan, K. R. (1998). *Moment Functions in Image Processing: Theory and Applications*. World Scientific, Singapore.
- Pearson, R. K. (2020). Exploring data with the moment ratio diagram. *The American Statistician*, 74(4):400–410.

- Pébay, P. (2008). Formulas for robust, one-pass parallel computation of covariances and arbitrary-order statistical moments. Technical Report SAND2008-6212, Sandia National Laboratories, Albuquerque, NM.
- Pébay, P., Terriberly, T. B., Kolla, H., and Bennett, J. (2016). Numerically stable, one-pass, parallel computation of arbitrary-order statistical moments. *Computational Statistics*, 31(4):1271–1301.
- Terriberly, T. B. (2007). Computing higher-order moments online. Technical report, Xiph.org Foundation.
- Wickham, H. (2015). *R Packages: Organize, Test, Document, and Share Your Code*. O’Reilly Media, Sebastopol, CA.