

# Statistics MCQ Question Bank

First Paper

Abdullah Al Mahmud

Last updated: November 18, 2023



[www.statmania.info](http://www.statmania.info)

# 1 Basic Concept of Statistics

1. **Who is known as the Father of modern statistics?**  
(a) P.C. Mahalanobis      (b) Kazi Motaher Hos-sain      (c) Karl Pearson      (d) R.A. Fisher
2. **If  $\sum_{i=1}^{20} x_i^2 = 20$  and  $\sum_{i=1}^{20} x_i = 30$ , what is the value of  $\sum_{i=1}^{20} x_i^2 + \sum_{i=1}^{20} x_i + 100$ ?**  
(a) 130      (b) 200      (c) 150      (d) 2130
3. **A subset of a population is called—**  
(a) Constant      (b) Variable      (c) Sample      (d) Scale
4. **How many measurement scales are there?**  
(a) 2      (b) 3      (c) 4      (d) 5
5. **Which of the following is a continuous variable?**  
(a) Number of goals      (b) Natural number  
(c) Summation of Fibonacci series      (d) Success rate
6. **In which scale of measurement, zero is regarded as true zero?**  
(a) Nominal scale      (b) Interval scale      (c) Ratio scale      (d) Ordinal scale
7. **Which is a discrete variable?**  
(a) Weight      (b) Amount of rainfall      (c) Distance      (d) Grade in a subject
8. **If  $x_1 = 2, x_2 = -3, x_3 = 7$ , and  $x_4 = 12$ ,  $\sum_{i=1}^4 x_i^2 = ?$**   
(a) 26      (b) 106      (c) 206      (d) 216
9. **If  $x_1 = 2, x_2 = 3, x_3 = 4, x_4 = 6$ , and  $x_5 = 5$ ,  $\sum_{i=1}^4 x_i^2 = ?$**   
(a) 80      (b) 87      (c) 90      (d) 105
10. **Capital and profit belong to a variable which is—**  
i. Bivariate  
ii. Quantitative  
iii. Qualitative  
**Which one is correct?**  
(a) i and ii      (b) i and iii      (c) ii and iii      (d) i, ii and iii
11. **Which one falls in the category of interval scale?**  
(a) Temperature      (b) Speed      (c) Distance      (d) Film rating
12. **In which scale of measurement, zero is regarded as true zero?**  
(a) Nominal scale      (b) Interval scale      (c) Ratio scale      (d) Ordinal scale
13. **Which is a discrete variable?**  
(a) Weight      (b) Amount of rainfall      (c) Distance      (d) Grade in a subject

14. Which one is product of square?  
 (a)  $\prod x_i^2$  (b)  $(\prod x_i)^2$  (c)  $\sum x_i^2 \times \sum x$  (d)  $\sum x_i^2$
15. For which variable, determining number of terms is not possible?  
 (a) Discrete variable (b) Continuous variable (c) Quantitative variable (d) Qualitative variable
- Answer the next three question based on the following information.**  
**A farmer collects growth (in cm) of 10 plants in a month and finds that  $\sum x_i = 7$  and  $\sum x_i^2 = 15$**
16. What is the value of  $\sum(x_i + 4)$ ?  
 (a) 23 (b) 47 (c) 22 (d) 11
17. If  $x_1 = 2, x_2 = 3, x_3 = 5, x_4 = 7$  and  $y_1 = 3, y_2 = 4, y_3 = 5, y_4 = 8$ ;  $\sum_{i=1}^4 x_i y_i = ?$   
 (a) 14 (b) 201 (c) 93 (d) 109
18. From the following table,  $\sum_{i=1}^4 x_i y_i = ?$
- |   |    |    |   |    |
|---|----|----|---|----|
| X | 1  | 5  | 3 | 2  |
| Y | 20 | 12 | 3 | 14 |
- (a) 14 (b) 201 (c) 99 (d) 109
19. What is the value of  $\sum(x_i - 4)^2$ ?  
 (a) 23 (b) 135 (c) 484 (d) 119
20. If the square of summation is subtracted the sum of square, the value is -  
 (a) -8 (b) 34 (c) 8 (d) -34
21. Which one is not an example of ratio scale?  
 (a) Room no. (b) Income (c) Number of accidents (d) Weight
22. Which one is discrete?  
 (a) Weight (b) Amount of rainfall  
 (c) Temperature (d) No. of member in a family
23. Which type of scale of measurement are religion and blood group?  
 (a) Interval (b) Ratio (c) Nominal (d) Ordinal

## 2 Collection, Organization, and Presentation of Data

24. How many sources of data are there?  
 (a) 5 (b) 4 (c) 3 (d) 2
25. What is the raw material of research?  
 (a) Data (b) Theory (c) Graph (d) Mean
26. Data obtained through direct observation is called—  
 (a) Primary data (b) Secondary data (c) Original Data (d) Informal data

27. **Who invented Stem and Leaf plot?**  
 (a) Karl Pearson (b) R.A. Fisher (c) David Cox (d) John Tukey
28. **Which rule is suggested by H.G. Sturges for determining number of class (k)?**  
 (a)  $K = 1 + 3.322 \log N$  (b)  $K = 1 + 3.222 \log N$  (c)  $K = 1 - 3.222 \log N$  (d)  $K = 1 + 2.332 \log N$
29. **To show runs per over in a cricket match, which diagram can be used?**  
 (a) Histogram (b) Bar Diagram (c) Ogive (d) Frequency polygon

### 3 Measures of Central Tendency

#### 3.1 General Questions

30. **If a value is zero, which measure is not usable?**  
 (a) Arithmetic Mean (b) Harmonic Mean (c) Geometric Mean (d) Mode
31. **How many measure of central tendency are there?**  
 (a) 2 (b) 3 (c) 4 (d) 5
32. **Which measure of central tendency is suitable for qualitative variable?**  
 (a) Arithmetic Mean (b) Harmonic Mean (c) Quadratic Mean (d) Mode
33. **In presence of negative values, which measure is not usable?**  
 (a) Arithmetic Mean (b) Geometric Mean (c) Quadratic Mean (d) Harmonic Mean
34. **Inappropriate for algebraic analysis—**  
 i. Median  
 ii. Mode  
 iii. Geometric Mean  
 Which one is true?  
 (a) i (b) ii (c) i & ii (d) ii & iii

**Answer the next two questions based on the following information**

Accident	4	6	7	8	9
Frequency	2	0	4	4	1

35. **Fifth Decile is —**  
 (a) 0 (b) 8 (c) 7 (d) 6
36. **Which of the following is mode?**  
 (a) 4 (b) 8 (c) 0 (d) 7
37. **Which measure gives a value from within the values?**  
 (a) Arithmetic Mean (b) Geometric Mean (c) Median (d) Mode
38. **Which one is not a proper measure of central tendency?**  
 (a) 2nd Quartile (b) Third Decile (c) 3rd Quintile (d) 110th Percentile

39. Which one is smallest?

- (a)  $\sum_{i=1}^n (X_i - \text{Median})^2$  (b)  $\sum_{i=1}^n (X_i - \bar{X})^2$  (c)  $\sum_{i=1}^n (X_i - \sigma)^2$  (d)  $\sum_{i=1}^n (X_i - \text{Mode})^2$

40. Which measure is not used in determining skewness?

- (a) Arithmetic Mean (b) Geometric Mean (c) Median (d) Mode

41. When is the relationship  $AM = HM = GM$  true?

- (a) All values are equal (b) The values form a geometric progression  
(c) The values form an arithmetic progression (d) All values are distinct

42. In the presence of outlier(s), which measure of central tendency is suitable?

- (a) Arithmetic mean (b) Median (c) Quadratic mean (d) Power mean

43. If a rate is defined as  $R = \frac{c}{d}$ , where c is constant, then which measure is perfect?

- (a) Weighted arithmetic mean (b) Harmonic mean  
(c) Quadratic mean (d) Weighted geometric mean

44. Which measure might have more than one value?

- (a) Arithmetic mean (b) Geometric mean (c) Quadratic mean (d) Mode

45. Which relationship is correct?

- (a)  $AM \times GM = HM^2$  (b)  $AM \times HM = GM^2$  (c)  $AM \times HM = GM^3$  (d)  $AM \div GM = HM^2$

46. With negative observations, which cannot be used

- i. Arithmetic Mean  
ii. Geometric Mean  
iii. Harmonic Mean

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

### 3.2 Arithmetic Mean

47. Arithmetic Mean of first 25 natural numbers is –

- (a) 12 (b) 13 (c) 14 (d) 26

48. Arithmetic Mean of two numbers is 25. If a number is 40, what is the other number?

- (a) 40 (b) 50 (c) 25 (d) 10

49. Number of students in two classes are 50 and 55 and their combined arithmetic mean (AM) of marks is 82. If AM of the first class is 75, what is the AM of the other class?

- (a) 88.36 (b) 88.40 (c) 84.55 (d) 78.33

50. The summation of deviation of each value from their arithmetic mean is –

- (a) 0 (b) 1 (c) 2 (d) 4

51. For grouped data, which formula is correct for Arithmetic Mean?

- (a)  $\bar{x} = \frac{\sum f_i x_i}{\sum f_i}$  (b)  $\bar{x} = \frac{\sum x_i}{N}$  (c)  $\bar{x} = \frac{\sum f_i x_i}{n}$  (d)  $\bar{x} = \frac{\sum f_i}{N}$

52. Arithmetic mean of the series 2, 12, 22, ..., 92 is—  
 (a) 45 (b) 46 (c) 47 (d) 55
53. What is the arithmetic mean of first  $n$  odd natural numbers?  
 (a)  $\frac{n+1}{n}$  (b)  $n$  (c)  $n+1$  (d)  $\frac{n+1}{2}$
54. What is the arithmetic mean of first  $n$  even natural numbers?  
 (a)  $\frac{n+1}{2}$  (b)  $n+1$  (c)  $n$  (d)  $\frac{n-1}{2}$
55. The arithmetic mean of first  $n$  natural numbers—  
 (a)  $\frac{n}{2}$  (b)  $\frac{n+1}{2}$  (c)  $\frac{n^2}{2}$  (d)  $\frac{n^2-1}{2}$
56. Arithmetic means of three groups having equal no. of items are 30, 32, and 34. What is the combined mean?  
 (a) 30.33 (b) 32.67 (c) 32.00 (d) 33.00

### 3.3 Geometric Mean

57. Which data set is suitable for Geometric Mean?  
 (a) 1, -1, 2, 4, 6, 7 (b) 1, 2, 4, 8, 16, 32 (c) 0, 1, 2, 3, 4, 6 (d) 1, 1, 2, 3, 4, 4, 5

### 3.4 Mode

58. Which of the following may be used to determine mode?  
 (a) Histogram (b) Frequency Curve (c) Ogive (d) Frequency Polygon

### 3.5 Median

59. Median can be determined from the—  
 (a) Histogram (b) Frequency curve (c) Ogive (d) Pie Chart

Answer the next two (2) questions based on the following information

Class	$\leq 20$	20-25	25-50	50-60	60-70	$\geq 70$
Frequency	5	10	10	7	5	3
Cumulative Frequency	5	15	25	32	37	40

60. How many values are between 20 and 70?  
 (a) 20 (b) 32 (c) 35 (d) 37
61. Which one is the median class?  
 (a) 20-25 (b) 25-50 (c) 50-60 (d) 60-70

### 3.6 Partition Values

Answer the next two questions as per the following information.

42 44 59 64 70 72 74 91 94 are 9 values.

62. What is the 50th percentile?

- (a) 64 (b) 70 (c) 72 (d) 71

63. Below which value lie 70 percent values?

- (a) 42 (b) 44 (c) 59 (d) 74

64. Above which value lie 30% observations?

- (a) 3rd Quartile (b) Median (c) 30th Percentile (d) 70th percentile

## 4 Measures of Dispersion

65. Which of the following is the best measure of dispersion?

- (a) Range (b) Mean deviation  
(c) Standard deviation (d) Coefficient of variation

66. What is the minimum possible value of standard deviation?

- (a)  $\infty$  (b) -1 (c) 0 (d) 1

67. For two values, range is found to be 8. What are the values of mean deviation and standard deviation

- (a) (2,4) (b) (4,4) (c) (4,8) (d) (8,8)

68. What is the standard deviation of first 10 natural numbers?

- (a) 2.87 (b) 3.02 (c) 0 (d) 2.78

69. Which measure is unit-free?

- (a) Range (b) Mean deviation  
(c) Standard deviation (d) Coefficient of variation

## 5 Moments, Skewness, and Kurtosis

### 5.1 Moments

70. Which quantity uniquely characterizes a distribution?

- (a) Median (b) Quantile (c) Moments (d) Trend

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

71. Which can be used to measure dispersion?

- (a)  $\mu'_2$  (b)  $\mu_1$  (c)  $\mu_2$  (d)  $\mu'_1$

72. The formula of coefficient of variance (CV) is –

- (a)  $\frac{\mu_2}{n} \times 100$  (b)  $\frac{\mu_2}{\mu_1} \times 100$  (c)  $\frac{\mu_2}{\bar{x}} \times 100$  (d)  $\frac{\mu_3}{\sigma} \times 100$

73. **First moment around zero is –**  
 (a) 0 (b) 1 (c) -1 (d) Arithmetic Mean
74. **Which might have a negative value?**  
 (a)  $\mu_4$  (b)  $\mu_3$  (c)  $\mu'_2$  (d)  $\mu_2$
75. **2nd Central Moment is –**  
 (a)  $\mu_2 - \mu'_1$  (b)  $\mu_2 + \mu'_1$  (c)  $\mu_2 - \mu'^2_1$  (d)  $\mu'_2 - \mu'^2_1$
76. **First central moment is equal to –**  
 (a) 1 (b) 0 (c) -1 (d)  $\bar{x} - a$
77. **First moment around a is equal to –**  
 (a) 1 (b) 0 (c) -1 (d)  $\bar{x} - a$
78. **The first raw moment about 3 is -5. What is the value of arithmetic mean?**  
 (a) 2 (b) -2 (c) 0 (d) 8
79. **Moments can be–**  
 i. positive  
 ii. not negative  
 iii. positive or negative  
**Which one is correct?**  
 (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

## 5.2 Skewness

80. **Characteristics of a skewed distributon are –**  
 i.  $Mean \neq Median \neq Mode$   
 ii. Differences of upper and lower quartiles from median are unequal  
 iii. Frequency curve is asymmetric
81. **In a distribution,  $\mu_2 = 25$ ,  $\mu_3 = 20$ , and  $\mu_4 = 2200$ ; the distribution is –**  
 (a) Negatively skewed (b) leptokurtic (c) Platykurtic (d) Symmetric
82. **For a data,  $Q_3 = 41.6$ ,  $Q_1 = 17.2$ ,  $Median = 29$ , &  $AM = 30$ ; What is Coefficient of skewness?**  
 (a) 24.4 (b) 1 (c) 0.03 (d) 29.45
83. **In case of positive skewness, which one is correct?**  
 (a)  $Mean > Median > Mode$  (b)  $Mean < Median < Mode$   
 (c)  $Mean = Median = Mode$  (d)  $Mean > Median < Mode$
84. **For a symmetrical distribution,  $\beta_1 =$**   
 (a) 1 (b) -1 (c) 0 (d) 3
85.  **$\sqrt{\beta_1} = -0.23$  implies–**  
 (a) Left Skew (b) Symmetry (c) Right Skew (d) Mesokurtic
86. **First 3 moments about 2 are 1, 2 and 8, respectively. What is the arithmetic mena?**  
 (a) 1 (b) 2 (c) 3 (d) 4



87. What is the second central moments of first 10 natural numbers?  
 (a) 9.90 (b) 9.09 (c) 8.25 (d) 5.67
88. Frequencies of higher values are smaller in – distribution  
 (a) Positively skewed (b) Negatively skewed (c) Symmetric (d) Mesokurtic

### 5.3 Kurtosis

89. How many types of kurtosis are there?  
 (a) 2 (b) 3 (c) 4 (d) 5
90. The standard deviation of a mesokurtik distribution is 2. What is the value of the 4th central moment?  
 (a) 4 (b) 8 (c) 16 (d) 48
91.  $\beta_2 = \sqrt{9}$  implies data are—  
 (a) Leptokurtic (b) Platykurtic (c) Mesokurtic (d) Symmetric
92. For a mesokurtik distribution,  $\beta_2 = --$   
 (a) 0 (b) -3 (c) 3 (d) 1

### 5.4 Misc

93. Which is not used in constructing Box & Whisker Plot?  
 (a) Mode (b)  $X_L$  (c)  $Q_1 \& Q_3$  (d)  $Q_1, Q_2 \& Q_3$
94. In a symmatric distribution—  
 i. Arithmetic Mean = Mode = Median  
 ii.  $Q_2 - Q_1 = Q_3 - Q_2$   
 iii.  $Q_1 - X_L = X_H - Q_3$   
 Which one is true?  
 (a) i & ii (b) ii & iii (c) i & iii (d) i, ii & iii
95. Which is not included in five number summary?  
 (a) Arithmetic Mean (b)  $X_H$  (c)  $Q_2$  (d)  $Q_3$

## 6 Correlation and Regression

## 7 Time Series

96. Time Series has how many components?  
 (a) 2 (b) 3 (c) 4 (d) 5
97. Which component involves period more than one (01) year?  
 (a) Seasonal Variation (b) Cyclic Variation (c) Irregular Variation (d) Random Variation
98. Which one is not a component of Time Series  
 (a) Seasonal Variation (b) Cyclic Variation (c) General Trend (d) Regular Variation

99. A company is constantly getting greater revenue than previous year; this is—  
 (a) Seasonal Variation (b) General Trend (c) Irregular Variation (d) Cyclic Variation

100. Which is not a method of finding general trend?  
 (a) Graphical Method (b) Moving Average (c) Semi-Average (d) Moving Median

Answer the next two questions based on the following table:

Year	2007	2008	2009	2010	2011	2012
Sales	5	35	34	40	42	204

101. In Semi-Average method, what is the 2nd average?  
 (a) 74 (b) 24.67 (c) 95.33 (d) 28
102. What is the last value of 3-yearly moving average?  
 (a) 93.55 (b) 95.53 (c) 95.33 (d) 59.33
103. Which component of time series is affected by economic changes due to war?  
 (a) Trend (b) Seasonal Variation (c) Irregular Variation (d) Cyclic Variation
104. Demand for warm clothes is higher in winter season and less in summer. Which component of time series deals with this change?  
 (a) Trend (b) Seasonal Variation (c) Irregular Variation (d) Cyclic Variation
105. Death rates of a country for 7 years are given below:

Year	2009	2010	2011	2012	2013	2014	2015
Rate	5	7	6	8	7	12	13

In semi-average method, which year will be excluded?

- (a) 2012 (b) 2013 (c) 2015 (d) 2009
106. Which component of time series represents a natural disaster?  
 (a) Seasonal Variation (b) General Trend (c) Irregular Variation (d) Cyclic Variation
107. How many models of time series are there to combine the components?  
 (a) 2 (b) 3 (c) 4 (d) 5

## 8 Published Statistics in Bangladesh

108. Limitations of published statistics in Bangladesh are –

- i. Wrong data collection method
- ii. Insufficient data
- iii. Lack of proper training

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii
109. How many sources of published statistics are there in Bangladesh?  
 (a) 2 (b) 3 (c) 4 (d) 6

110. **Bangladesh Bureau of Statistics collect –**

- (a) Official statistics      (b) Non-official statistics(c) Semi-official statistics(d) None of the above

111. **Which statistics are published by an NGO?**

- (a) Official statistics      (b) Non-official statistics(c) Semi-official statistics(d) None of the above

112. **The primary source of official statistics in Bangladesh is –**

- (a) WHO                      (b) BBS                      (c) CPD                      (d) UNDP

113. **In Bangladesh, a census is usually done every – years**

- (a) 20                      (b) 15                      (c) 10                      (d) 12

Answer Key:

1. (d) R.A. Fisher
2. (c) 150
3. (c) Sample
4. (c) 4
5. (d) Success rate
6. (c) Ratio scale
7. (d) Grade in a subject
8. (c) 206
9. (c) 90
10. (a) i and ii
11. (a) Temperature
12. (c) Ratio scale
13. (d) Grade in a subject
14. (a)  $\prod x_i^2$
15. (b) Continuous variable
16. (b) 47
17. (c) 93
18. (c) 99
19. (d) 119
20. (d) -34
21. (a) Room no.
22. (d) No. of member in a family
23. (c) Nominal
24. (d) 2
25. (a) Data
26. (a) Primary data
27. (d) John Tukey
28. (a)  $K = 1 + 3.322 \log N$
29. (b) Bar Diagram
30. (c) Geometrtic Mean
31. (d) 5
32. (d) Mode
33. (b) Geometric Mean
34. (c) i & ii
35. (c) 7
36. (b) 8
37. (d) Mode
38. (d) 110th Percentile
39. (a)  $\sum_{i=1}^n (X_i - Median)^2$
40. (b) Geometric Mean
41. (a) All values are equal
42. (b) Median
43. (b) Harmonic mean
44. (d) Mode
45. (b)  $AM \times HM = GM^2$
46. (c) ii and iii
47. (b) 13
48. (d) 10
49. (a) 88.36
50. (a) 0
51. (a)  $\bar{x} = \frac{\sum f_i x_i}{\sum f_i}$
52. (c) 47
53. (b) n
54. (b)  $n + 1$
55. (b)  $\frac{n+1}{2}$
56. (c) 32.00
57. (b) 1, 2, 4, 8, 16, 32
58. (a) Histogram
59. (c) Ogive
60. (b) 32
61. (b) 25-50
62. (b) 70
63. (d) 74
64. (d) 70th percentile
65. (c) Standard deviation
66. (c) 0
67. (a) (2,4)
68. (a) 2.87
69. (d) Coefficient of variation
70. (c) Moments
70. (d) i, ii and iii
71. (c)  $\mu_2$
72. (c)  $\frac{\mu_2}{\bar{x}} \times 100$
73. (d) Arithmetic Mean
74. (b)  $\mu_3$
75. (d)  $\mu'_2 - \mu'^2_1$
76. (b) 0
77. (d)  $\bar{x} - a$
78. (b) -2
79. (b) i and iii
81. (b) leptokurtic
82. (d) 29.45
83. (a)  $Mean > Median > Mode$
84. (c) 0
85. (a) Left Skew
86. (c) 3
87. (c) 8.25
88. (a) Positively skewed
89. (b) 3
90. (d) 48
91. (c) Mesokurtic
92. (c) 3
93. (a) Mode
94. (d) i, ii & iii
95. (a) Arithmetic Mean
96. (c) 4
97. (b) Cyclic Variation
98. (d) Regular Variation
99. (b) General Trend
100. (d) Moving Median
101. (c) 95.33
102. (c) 95.33
103. (c) Irregular Variation
104. (b) Seasonal Variation
105. (b) 2013
106. (c) Irregular Variation
107. (a) 2
108. (d) i, ii and iii
109. (b) 3
110. (a) Official statistics
111. (c) Semi-official statistics
112. (b) BBS
113. (c) 10