

Statistics MCQ Question Bank

First Paper

Abdullah Al Mahmud

Last updated: April 28, 2026



www.statmania.info

Contents

1	Basic Concept of Statistics	2
1.1	Situation Set	9
2	Collection, Organization, and Presentation of Data	10
2.1	Situation Set	10
3	Measures of Central Tendency	13
3.1	General Questions	13
3.2	Arithmetic Mean	15
3.3	Harmonic Mean	17
3.4	Geometric Mean	17
3.5	Mode	18
3.6	Median	18
3.7	Partition Values	18
3.8	Situation Set	18
3.9	Multiple Completion	20
4	Measures of Dispersion	21
4.1	Situation Set	22
4.2	Multiple Completion	22
5	Moments, Skewness, and Kurtosis	23
5.1	Moments	23
5.2	Skewness	25
5.2.1	Situation Set	27
5.2.2	Multiple Completion	28
5.3	Kurtosis	28
5.4	Misc	29
5.5	Box and Whisker Plot	29
5.6	Five Number Summary	30
6	Correlation and Regression	30
6.1	Correlation	30
6.2	Situation Set	31
7	Time Series	31
7.1	Situation Set	36
7.2	Multiple Completion	38
8	Published Statistics in Bangladesh	38

1 Basic Concept of Statistics

1. **Who is known as the Father of modern statistics?**
(a) P.C. Mahalanobis (b) Kazi Motaher Hos-sain (c) Karl Pearson (d) R.A. Fisher
2. **Which is not a function of statistics?**
(a) Data collection (b) Data organization (c) Analysis (d) Database creation
3. **Which one is an example of an infinite population?**
(a) Students of Dhaka University (b) Cadets of SCC
(c) Minor planets in the solar system (d) Red blood cells in a person's body
4. **Which of the following is an example of an infinite population?**
(a) Employees of a multinational company (b) Trees in a national park
(c) Stars in the Milky Way (d) Passengers on a flight
5. **Which one represents an infinite population?**
(a) Books in a library (b) Fish in the Pacific Ocean
(c) Members of a sports club (d) Mobile phones in a city
6. **A researcher collected data on age and income of the people in a city. The variables are –**
i. bi-variate
ii. quantitative
iii. qualitative
Which one is correct?
(a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii
7. **Which of the following is correct?**
(a) $\sum_{i=1}^{20} cx_i = nc \sum_{i=1}^{20} x_i$ (b) $\sum_{i=1}^{20} cx_i = nc \sum_{i=1}^{20} x_i$ (c) $\sum_{i=1}^{20} cx_i = c \sum_{i=1}^{20} x_i$ (d) $\sum_{i=1}^{20} cx_i = c^2 \sum_{i=1}^{20} x_i$
8. **Which cannot be performed using Univariate data?**
(a) Central tendency (b) Dispersion (c) Skewness (d) Regression
9. **Which of the following cannot be analyzed using univariate data?**
(a) Mean (b) Variance (c) Correlation (d) Range
10. **Which statistical method requires bivariate or multivariate data?**
(a) Standard deviation (b) Histogram (c) Regression analysis (d) Median
11. **Which of the following is an example of an infinite population?**
(a) Patients in a hospital (b) Water molecules in the ocean
(c) Cars on a highway (d) Students in a university
12. **Which of the following is an example of a finite population?**
(a) Books in a school library (b) Stars in the universe
(c) Grains of sand on a beach (d) Atoms in the atmosphere

13. Which one represents an infinite population?
- (a) Trees in a forest (b) Grains of sand on a beach
(c) Books in a bookstore (d) Houses in a neighborhood
14. Cities ranked according to habitability level show – measurement scale
- (a) Nominal (b) Ratio (c) Interval (d) Ordinal
15. Classifying students based on their grades (A, B, C, etc.) represents which measurement scale?
- (a) Nominal (b) Ordinal (c) Interval (d) Ratio
16. Temperature measured in Celsius or Fahrenheit follows which type of measurement scale?
- (a) Nominal (b) Ordinal (c) Interval (d) Ratio
17. A survey categorizing people by their favorite color is an example of which measurement scale?
- (a) Nominal (b) Ordinal (c) Interval (d) Ratio
18. Which is not an example of shift of scale?
- (a) $y_i = \frac{x_i}{a}$ (b) $y_i = cx_i$ (c) $y_i = x_i - 2$ (d) $y_i = \frac{cx_i}{d}$
19. If $\sum_{i=1}^{20} x_i^2 = 20$ and $\sum_{i=1}^{20} x_i = 30$, what is the value of $\sum_{i=1}^{20} x_i^2 + \sum_{i=1}^{20} x_i + 100$?
- (a) 130 (b) 200 (c) 150 (d) 2130
20. If $\sum_{i=1}^{15} y_i^2 = 50$ and $\sum_{i=1}^{15} y_i = 25$, what is the value of $\sum_{i=1}^{15} y_i^2 - \sum_{i=1}^{15} y_i + 75$?
- (a) 100 (b) 50 (c) 125 (d) 45
21. Given $\sum_{i=1}^{10} a_i^2 = 40$ and $\sum_{i=1}^{10} a_i = 20$, find the value of $2\sum_{i=1}^{10} a_i^2 - 3\sum_{i=1}^{10} a_i + 60$.
- (a) 70 (b) 100 (c) 80 (d) 50
22. If $\sum_{i=1}^{25} z_i^2 = 75$ and $\sum_{i=1}^{25} z_i = 50$, compute $\sum_{i=1}^{25} z_i^2 + 2\sum_{i=1}^{25} z_i - 125$.
- (a) 50 (b) 75 (c) 100 (d) 25
23. A subset of a population is called–
- (a) Constant (b) Variable (c) Sample (d) Scale
24. What is $\sum_{i=1}^n bx_i$ equal to?
- (a) $b \sum_{i=1}^n nx_i$ (b) $b \sum_{i=1}^n x_i$ (c) $\sum_{i=1}^n nx_i$ (d) $bn \sum_{i=1}^n x_i$
25. How many measurement scales are there?
- (a) 2 (b) 3 (c) 4 (d) 5

26. **Which of the following is a continuous variable?**
 (a) Number of goals (b) Natural number
 (c) Summation of Fibonacci series (d) Success rate
27. **In which scale of measurement, zero is regarded as true zero?**
 (a) Nominal scale (b) Interval scale (c) Ratio scale (d) Ordinal scale
28. **Which measurement scale does height belong to?**
 (a) Nominal (b) Ordinal (c) Interval (d) Ratio
29. **Which is a discrete variable?**
 (a) Weight (b) Amount of rainfall (c) Distance (d) Grade in a subject
30. **Which is a discrete variable?**
 (a) Height of a building (b) Number of cars in a parking lot
 (c) Amount of milk in a container (d) Time taken to complete a task
31. **Which is a discrete variable?**
 (a) Speed of a car (b) Number of students in a class
 (c) Volume of water in a tank (d) Temperature of a room
32. **Which is a discrete variable?**
 (a) Blood pressure (b) Number of books on a shelf
 (c) Length of a river (d) Amount of sugar in a cup
33. **Which is a discrete variable?**
 (a) Shoes sizes available in a store (b) Distance between two cities
 (c) Volume of a gas (d) Weight of a parcel
34. **Which is a discrete variable?**
 (a) Grades on a multiple-choice test (A, B, C, D) (b) Temperature during the day
 (c) Height of a person (d) Time spent on an activity
35. **Which is a discrete variable?**
 (a) Outcomes of rolling a die (b) Speed of a train
 (c) Rainfall in a region (d) Age of a tree
36. **Which is a discrete variable?**
 (a) Counts of people in a room (b) Temperature recorded every hour
 (c) Weight of an animal (d) Height of a plant
37. **Which is a discrete variable?**
 (a) Number of languages spoken by a person (b) Time taken to complete a race
 (c) Length of a road (d) Volume of water in a tank
38. **Which is a discrete variable?**
 (a) Length of a rope (b) Weight of books in a library
 (c) Distance (d) No. of particles in atoms

39. If $x_1 = 2, x_2 = -3, x_3 = 7$, and $x_4 = 12$, $\sum_{i=1}^4 x_i^2 = ?$
- (a) 26 (b) 106 (c) 206 (d) 216
40. If $x_1 = 5, x_2 = -4, x_3 = 9$, and $x_4 = 0$, what is $\sum_{i=1}^4 x_i^2$?
- (a) 82 (b) 97 (c) 107 (d) 122
41. If $x_1 = 3, x_2 = 2, x_3 = -6$, and $x_4 = 4$, what is $\sum_{i=1}^4 x_i^2$?
- (a) 45 (b) 65 (c) 85 (d) 89
42. If $x_1 = 4, x_2 = 1, x_3 = -2$, and $x_4 = 3$, find $\sum_{i=1}^4 (x_i^2 + 3)$?
- (a) 40 (b) 50 (c) 42 (d) 56
43. If $y_1 = 5, y_2 = 2, y_3 = -1$, and $y_4 = 4$, $\sum_{i=1}^4 (y_i^2 + 2) = ?$
- (a) 50 (b) 40 (c) 54 (d) 60
44. Given $z_1 = 3, z_2 = 0, z_3 = -3$, and $z_4 = 2$, determine $\sum_{i=1}^4 (z_i^2 + 5)$.
- (a) 30 (b) 40 (c) 35 (d) 45
45. If $x_1 = 4, x_2 = -2, x_3 = 1$, and $x_4 = 5$, calculate $\sum_{i=1}^4 (2x_i^2 - x_i)$?
- (a) 38 (b) 42 (c) 46 (d) 84
46. If $x_1 = 3, x_2 = 1, x_3 = 0$, and $x_4 = 2$, find $\sum_{i=1}^4 x_i^2 - \sum_{i=1}^4 x_i$?
- (a) 7 (b) 9 (c) 8 (d) 13
47. If $x_1 = 5, x_2 = 4, x_3 = -3$, and $x_4 = 2$, find $\sum_{i=1}^4 (x_i^2 + x_i)$?
- (a) 58 (b) 62 (c) 66 (d) 72
48. If $x_1 = 2, x_2 = 3, x_3 = -1$, and $x_4 = 0$, calculate $\sum_{i=1}^4 (x_i^2 - 2)$?
- (a) 0 (b) 6 (c) 8 (d) 10
49. If $x_1 = 2, x_2 = 3, x_3 = 4, x_4 = 6$, and $x_5 = 5$, $\sum_{i=1}^4 x_i^2 = ?$
- (a) 80 (b) 87 (c) 90 (d) 105

50. If $f_i = 3, 5, 7$ and $x_i = 2, 4, 7$; what is the value of $\sum_{i=1}^3 f_i x_i^2$?
- (a) 450 (b) 350 (c) 345 (d) 435
51. If $f_i = 2, 4, 6$ and $x_i = 3, 5, 7$, what is the value of $\sum_{i=1}^3 f_i x_i^3$?
- (a) 950 (b) 1125 (c) 2612 (d) 1330
52. Given $f_i = 1, 3, 5$ and $x_i = 2, 4, 6$, find the value of $\sum_{i=1}^3 f_i x_i^4$.
- (a) 1356 (b) 1536 (c) 1650 (d) 7264
53. If $f_i = 3, 5, 7$ and $x_i = 2, 4, 6$, compute $\sum_{i=1}^3 f_i x_i^2$.
- (a) 260 (b) 280 (c) 344 (d) 320
54. Find the value of $\sum_{i=1}^{12} f_i (x_i - 7)^2$ where $\sum_{i=1}^{12} f_i x_i^2 = 400$, $\sum_{i=1}^{12} f_i x_i = 40$, $\sum_{i=1}^{12} f_i = 10$
- (a) 320 (b) 330 (c) 250 (d) 430
55. If $x_1 = 3$, $x_2 = -1$, $x_3 = 2$, and $x_4 = 0$, find $\sum_{i=1}^4 (x_i^3 + 2x_i)$?
- (a) 12 (b) 18 (c) 24 (d) 28
56. If $x_1 = 4$, $x_2 = 1$, $x_3 = -2$, and $x_4 = 3$, calculate $\sum_{i=1}^4 (x_i^2 + 4x_i - 1)$?
- (a) 16 (b) 24 (c) 34 (d) 50
57. If $x_1 = 1$, $x_2 = 2$, $x_3 = -3$, and $x_4 = 4$, find $\sum_{i=1}^4 (3x_i^3 - x_i^2)$?
- (a) 108 (b) 114 (c) -8 (d) 201
58. If $x_1 = 5$, $x_2 = 0$, $x_3 = -1$, and $x_4 = 2$, determine $\sum_{i=1}^4 (x_i^3 + x_i^2 + 3)$?
- (a) 173 (b) 174 (c) 164 (d) 172
59. Capital and profit belong to a variable which is—
- Bivariate
 - Quantitative
 - Qualitative
- Which one is correct?
- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii
60. Which one falls in the category of interval scale?
- (a) Temperature (b) Speed (c) Distance (d) Film rating

61. Which one falls in the category of nominal scale?
 (a) Height (b) Temperature (c) Gender (d) Age
62. Which of the following is an example of an ordinal scale?
 (a) Temperature (b) IQ Score (c) Educational Level (d) Weight
63. Which of the following is not example of a ratio scale?
 (a) Temperature (b) Time (c) Blood Pressure (d) Speed
64. In which scale of measurement, zero is regarded as true zero?
 (a) Nominal scale (b) Interval scale (c) Ratio scale (d) Ordinal scale
65. Which is a discrete variable?
 (a) Weight (b) Amount of rainfall (c) Distance (d) Grade in a subject
66. Which one is product of square?
 (a) $\prod x_i^2$ (b) $(\prod x_i)^2$ (c) $\sum x_i^2 \times \sum x$ (d) $\sum x_i^2$
67. For which variable, determining number of terms is not possible?
 (a) Discrete variable (b) Continuous variable (c) Quantitative variable (d) Qualitative variable
- Answer the next three question based on the following information.**

A farmer collects growth (in cm) of 10 plants in a month and finds that
 $\sum x_i = 7$ and $\sum x_i^2 = 15$

68. Which is considered statistics?
 (a) Jaman obtained 75 in statistics (b) Shafiq lives at Road no. 5
 (c) Mean monthly income in a city is 60,000 taka (d) Width of a book is 10 cm
69. What is the value of $\sum(x_i + 4)$ if $x = \{2, 3\}$?
 (a) 23 (b) 47 (c) 22 (d) 13
70. If $x_1 = 2, x_2 = 3, x_3 = 5, x_4 = 7$ and $y_1 = 3, y_2 = 4, y_3 = 5, y_4 = 8$; $\sum_{i=2}^4 x_i y_i = ?$
 (a) 14 (b) 201 (c) 93 (d) 117
71. From the following table, $\sum_{i=1}^4 x_i y_i = ?$

X	1	5	3	2
Y	20	12	3	14

- (a) 14 (b) 201 (c) 99 (d) 109
72. What is the value of $\sum(x_i - 4)^2$?
 (a) 23 (b) 135 (c) 484 (d) 119
73. If the square of summation is subtracted the sum of square, the value is -
 (a) -8 (b) 34 (c) 8 (d) -34

74. Which one is not an example of ratio scale?

- (a) Room no. (b) Income (c) Number of accidents (d) Weight

75. Which one is discrete?

- (a) Weight (b) Amount of rainfall
(c) Temperature (d) No. of member in a family

76. Which type of scale of measurement are religion and blood group?

- (a) Interval (b) Ratio (c) Nominal (d) Ordinal

Answer the next two questions based on the following information

$$X = 20, 25, 30, 40$$

77. Find $\sum(X_i + 10)$

- (a) 150 (b) 155 (c) 125 (d) 250

78. $\sum(X_i - 30)^2$

- (a) 225 (b) 230 (c) 420 (d) 235

Answer the next two questions based on the following information

$$X = 3, 5, 7, 10$$

79. Find $\sum(X_i + 3)$

- (a) 28 (b) 32 (c) 37 (d) 40

80. $\sum(X_i - 5)^2$

- (a) 16 (b) 33 (c) 12 (d) 8

Answer the next two questions based on the following information

$$X = 6, 8, 10, 12$$

81. Find $\sum(X_i - 4)$

- (a) 20 (b) 30 (c) 32 (d) 22

82. $\sum(X_i + 2)^2$

- (a) 196 (b) 504 (c) 210 (d) 220

Answer the next two questions based on the following information

$$X = 4, 9, 13, 15$$

83. Find $\sum(2X_i)$

- (a) 68 (b) 70 (c) 82 (d) 74

84. $\sum(X_i - 10)^2$

- (a) 71 (b) 80 (c) 85 (d) 92

Answer the next three questions based on the following information.

The values of x_i and f_i are given below:

x_i	1	2	3	4
f_i	2	3	4	1

85. Find $\sum_{i=1}^4 f_i x_i$.

(a) 20

(b) 21

(c) 22

(d) 24

86. Compute $\sum_{i=1}^4 f_i x_i^2$.

(a) 30

(b) 35

(c) 66

(d) 64

87. Determine $\sum_{i=1}^4 f_i^2 x_i$.

(a) 74

(b) 49

(c) 78

(d) 65

Answer the next three questions based on the following information.

The values of x_i and f_i are given below:

x_i	2	4	6	8
f_i	2	2	5	4

88. Find $\sum_{i=1}^4 f_i x_i$.

(a) 50

(b) 74

(c) 56

(d) 60

89. Compute $\sum_{i=1}^4 f_i x_i^2$.

(a) 256

(b) 274

(c) 476

(d) 300

90. Determine $\sum_{i=1}^4 f_i (x_i - 5)^2$.

(a) 61

(b) 48

(c) 52

(d) 58

1.1 Situation Set

Answer the next TWO questions based on the following information

The table below shows monthly sales (in thousands of dollars) for two products in four regions.

Product	North	South	East	West
A	12	15	10	8
B	18	22	14	16

91. Calculate total sales of Product A across all regions.

(a) 40

(b) 42

(c) 45

(d) 48

92. Calculate total sales of all products across all regions.

(a) 95

(b) 100

(c) 105

(d) 115

2 Collection, Organization, and Presentation of Data

93. How many sources of data are there?
 (a) 5 (b) 4 (c) 3 (d) 2
94. What is the raw material of research?
 (a) Data (b) Theory (c) Graph (d) Mean
95. Data obtained through direct observation is called—
 (a) Primary data (b) Secondary data (c) Original Data (d) Informal data
96. Which formula is used to find angles for Pie Chart?
 (a) $\theta_i = \frac{f_i}{N} \times 100$ (b) $\theta_i = \frac{f_i}{100} \times 360$ (c) $\theta_i = \frac{f_i}{N} \times 360$ (d) $\theta_i = \frac{f_i}{N-1} \times 360$
97. Who invented Stem and Leaf plot?
 (a) Karl Pearson (b) R.A. Fisher (c) David Cox (d) John Tukey
98. If all the rats in Sylhet is a population, all the rats in Sylhet Airport is –
 (a) Data (b) Sample (c) Statistics (d) Frequency
99. Which rule is suggested by H.G. Sturges for determining number of class (k)?
 (a) $K = 1 + 3.322 \log N$ (b) $K = 1 + 3.222 \log N$ (c) $K = 1 - 3.222 \log N$ (d) $K = 1 + 2.332 \log N$
100. To show runs per over in a cricket match, which diagram can be used?
 (a) Histogram (b) Bar Diagram (c) Ogive (d) Frequency polygon

2.1 Situation Set

Answer the next THREE questions based on the following information

Radius of 80 trees are recorded and this frequency distribution is constructed.

Radius (cm)	0-10	10-20	20-30	30-40
No. of Trees	20	15	21	24

101. How many trees have radius between 10 and 30?
 (a) 30 (b) 15 (c) 36 (d) 21
102. How many trees have radius at least 20?
 (a) 44 (b) 45 (c) 24 (d) 21
103. What percent of trees have radius between 20 and 40?
 (a) 44% (b) 56% (c) 46% (d) 53%

Answer the next two questions based on the following plot

Data: 18, 21, 22, 23, 24, 26, 31, 33, 33, 35, 37, 42

Stem	Leaf
1	8
2	1 2 3 4 6
3	1 3 3 5 7
4	2

Key: 2 | 1 means 21

104. How many data values are greater than 30 in the stem-and-leaf plot?

- (a) 3 (b) 4 (c) 5 (d) 6

105. What is the median of the data shown in the stem-and-leaf plot?

- (a) 26 (b) 31 (c) 30 (d) 29

Answer the next two questions based on the following plot

Stem	Leaf
1	3 7 8
2	0 2 5 9
3	1 1 4 6
4	0 3

Key: 2 | 5 = 25

106. What is the mode of the data set?

- (a) 31 (b) 22 (c) 13 (d) None (no mode)

107. What is the median of the data shown in the stem-and-leaf plot?

- (a) 26 (b) 31 (c) 30 (d) 29

Answer the next THREE questions based on the following information.

The heights of 100 plants were measured, and this frequency distribution was constructed.

Height (cm)	0-20	20-40	40-60	60-80
No. of Plants	25	30	20	25

108. How many plants have height between 20 and 60?

- (a) 50 (b) 30 (c) 20 (d) 25

109. How many plants have height at least 40?

- (a) 50 (b) 45 (c) 40 (d) 25

110. What percent of plants have height between 20 and 80?

- (a) 80% (b) 75% (c) 60% (d) 50%

Answer the next THREE questions based on the following information.

The weights of 120 fruits were recorded and this frequency distribution was constructed.

Weight (grams)	0-50	50-100	100-150	150-200
No. of Fruits	30	35	25	30

111. How many fruits weigh at least 100 grams?

- (a) 55 (b) 50 (c) 60 (d) 65

112. How many fruits weigh less than 100 grams?

- (a) 68 (b) 70 (c) 65 (d) 50

113. What percent of fruits weigh between 50 and 150 grams?

- (a) 50% (b) 55% (c) 60% (d) 75%

Answer the next two questions based on the following information

Class Interval	<10	10-20	20-30	30-40
Frequency	6	3	7	4

114. What is relative frequency of the class with the highest frequency?

- (a) 0.25 (b) 0.45 (c) 0.40 (d) 0.35

115. Which curve is suitable for

- (a) Histogram (b) Bar Diagram (c) Pie Chart (d) Ogive

116. Example of primary data —

- i. A student collected data for research
- ii. A professor had a student collect data for them
- iii. A researcher collected data from a newspaper.

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

117. Which of the following is an example of secondary data?

- i. Data obtained from a published journal
- ii. Data collected by a government agency and used by a researcher
- iii. Data gathered directly through interviews

Which one is correct?

- (a) i and ii (b) ii and iii (c) i and iii (d) i, ii and iii

118. Which of the following represents primary data?

- i. A scientist collects soil samples for analysis
- ii. Data compiled in a textbook
- iii. A business owner surveys customers directly

Which one is correct?

- (a) i and iii (b) i and ii (c) ii and iii (d) i, ii, and iii

119. Which of these are examples of secondary data?

- i. A report sourced from census data
- ii. A student conducting a direct experiment
- iii. Statistics extracted from a government database

Which one is correct?

- (a) i and iii (b) i and ii (c) ii and iii (d) i, ii, and iii

120. Which one is true of primary data?

- i. Original
- ii. Suitable
- iii. Reliable

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

121. Which statement is true about secondary data?

- i. Already published
- ii. Economical
- iii. Always up-to-date

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

122. Which one is true about secondary data?

- i. Easy to collect
- ii. Collected by someone else
- iii. Free from bias

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

123. Which is an advantage of primary data?

- i. Specific to the study
- ii. More reliable
- iii. Less time-consuming

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

3 Measures of Central Tendency

3.1 General Questions

124. Which statement is correct

- (a) Quartiles are well defined (b) Outliers affect Median
(c) Median is always present in data (d) Quadratic mean is widely used

125. Which measure is suitable for open-ended distribution?

- (a) Median (b) Mode (c) Geometric Mean (d) Arithmetic mean

126. Which is not a measure of central tendency?

- (a) Arithmetic mean (b) Mode (c) Range (d) Quadratic mean

127. When is the statement $AM = GM = HM$ true?

- (a) When the values are natural numbers (b) When all the values are equal
(c) When all the values have equal frequency (d) When mode is greater than median

128. If a value is zero, which measure is not usable?

- (a) Arithmetic Mean (b) Harmonic Mean (c) Geometric Mean (d) Mode

129. How many measure of central tendency are there?

- (a) 2 (b) 3 (c) 4 (d) 5

130. Which measure of central tendency is suitable for qualitative variable?

- (a) Arithmetic Mean (b) Harmonic Mean (c) Quadratic Mean (d) Mode

145. Which measure might have more than one value?
 (a) Arithmetic mean (b) Geometric mean (c) Quadratic mean (d) Mode
146. Which relationship is correct?
 (a) $AM \times GM = HM^2$ (b) $AM \times HM = GM^2$ (c) $AM \times HM = GM^3$ (d) $AM \div GM = HM^2$
147. The arithmetic mean and geometric mean of two non-zero positive numbers are 15 and 10, respectively. What is harmonic mean?
 (a) 6.61 (b) 6.67 (c) 7.66 (d) 6.76
148. For two non-zero positive numbers, the harmonic mean is 8 and the geometric mean is 12. What is the arithmetic mean?
 (a) 16 (b) 18 (c) 20 (d) 22
149. For two non-zero positive numbers, the harmonic mean is 10 and the arithmetic mean is 25. What is the geometric mean?
 (a) 7.07 (b) 20 (c) 25 (d) 30

3.2 Arithmetic Mean

150. If $\bar{X} = 20$, what is the value of $\bar{Y}$, if $Y = 4X$?
 (a) 24 (b) 5
 (c) 80 (d) Cannot be determined
151. If $\bar{X} = 12$, what is $\bar{Y}$ when $Y = 3X$?
 (a) 4 (b) 12 (c) 24 (d) 36
152. If $\bar{X} = 30$, what is $\bar{Y}$ when $Y = 2X + 5$?
 (a) 35 (b) 60 (c) 65 (d) 70
153. If $Y = 4X - 8$ and $\bar{Y} = 20$, find $\bar{X}$.
 (a) 3 (b) 5 (c) 7 (d) 9
154. If each observation is multiplied by 3 and then increased by 4, the arithmetic mean of the original data will be:
 (a) Multiplied by 3 (b) Increased by 4
 (c) Multiplied by 3 and then increased by 4 (d) Unchanged
155. Let $Y = aX + b$, where a and b are constants. If $\bar{X} = 8$ and $\bar{Y} = 30$, and $a = 3$, find b .
 (a) 3 (b) 6 (c) 9 (d) 12
156. If $\bar{X} = 25$, what is $\bar{Y}$ when $Y = X - 10$?
 (a) 10 (b) 15 (c) 25 (d) 35
157. If each value of a variable is multiplied with 4, what would be the new arithmetic mean?
 (a) 4 times (b) twice (c) Thrice (d) Unchanged
158. If $\sum(x_i - k) = 0$, what is the value of k ?
 (a) n (b) $\bar{x}$ (c) x (d) $n\bar{x}$
159. If $\sum(x_i - a)^2$ is minimized, then the value of a is:
 (a) $\bar{x}$ (b) 0 (c) Median (d) Mode

160. Find the arithmetic mean: 6, 9, 12, $\dots$, 84
 (a) 40 (b) 45 (c) 50 (d) 55
161. The arithmetic mean of first 10 natural numbers is:
 (a) 6 (b) 8.5 (c) 5.5 (d) 5.6
162. Arithmetic Mean of first 25 natural numbers is –
 (a) 12 (b) 13 (c) 14 (d) 26
163. An equation is: $y = 5x + 9$. If $\bar{x} = 20, \bar{y} = ?$
 (a) 100 (b) 209 (c) 109 (d) 29
164. An equation is: $y = 5x + 9$. If $\bar{x} = 20$, what is $\bar{y}$?
 (a) 100 (b) 209 (c) 109 (d) 29
165. Given the relationship $y = 2x - 4$, and $\bar{x} = 15$, find the value of $\bar{y}$.
 (a) 26 (b) 34 (c) -26 (d) 35
166. Arithmetic Mean of two numbers is 25. If a number is 40, what is the other number?
 (a) 40 (b) 50 (c) 25 (d) 10
167. The Arithmetic Mean of two numbers is 30. If one number is 40, what is the other number?
 (a) 20 (b) 30 (c) 40 (d) 60
168. The Arithmetic Mean of two numbers is 35. If one number is 50, what is the other number?
 (a) 25 (b) 20 (c) 40 (d) 70
169. Number of students in two classes are 50 and 55 and their combined arithmetic mean (AM) of marks is 82. If AM of the first class is 75, what is the AM of the other class?
 (a) 88.36 (b) 88.40 (c) 84.55 (d) 78.33
170. The summation of deviation of each value from their arithmetic mean is –
 (a) 0 (b) 1 (c) 2 (d) 4
171. For grouped data, which formula is correct for Arithmetic Mean?
 (a) $\bar{X} = \frac{\sum f_i x_i}{\sum f_i}$ (b) $\bar{X} = \frac{\sum x_i}{N}$ (c) $\bar{X} = \frac{\sum f_i x_i}{n}$ (d) $\bar{X} = \frac{\sum f_i}{N}$
172. Arithmetic mean of the series 2, 12, 22, $\dots$, 92 is–
 (a) 45 (b) 46 (c) 47 (d) 55
173. What is the arithmetic mean of first n odd natural numbers?
 (a) $\frac{n+1}{n}$ (b) n (c) n+1 (d) $\frac{n+1}{2}$
174. What is the arithmetic mean of first n even natural numbers?
 (a) $\frac{n+1}{2}$ (b) n + 1 (c) n (d) $\frac{n-1}{2}$
175. The arithmetic mean of first n natural numbers–
 (a) $\frac{n}{2}$ (b) $\frac{n+1}{2}$ (c) $\frac{n^2}{2}$ (d) $\frac{n^2-1}{2}$
176. Arithmetic means of three groups having equal no. of items are 30, 32, and 34. What is the combined mean?
 (a) 30.33 (b) 32.67 (c) 32.00 (d) 33.00

3.3 Harmonic Mean

177. Which formula is correct for harmonic mean?

- (a) $\frac{n}{\sum_{i=1}^n \frac{f_i}{x_i}}$ (b) $\frac{f_i}{\sum_{i=1}^n \frac{f_i}{x_i}}$ (c) $\frac{\sum f_i}{\sum_{i=1}^n \frac{f_i}{x_i}}$ (d) $\frac{\sum f_i}{\sum_{i=1}^n \frac{1}{x_i}}$

178. What is the harmonic mean of these values: 10, 12, 13, 15, 20, 25

- (a) 12.49 (b) 14.93 (c) 14.39 (d) 13.49

179. A rate is defined as $R = \frac{c}{d}$; c and d are arbitrary numbers. If c is constant, which mean is used?

- (a) Arithmetic Mean (b) Geometric Mean
(c) Harmonic Mean (d) Weighted Geometric Mean

180. A rate is defined as $R = \frac{c}{d}$; c and d are arbitrary numbers. If d is constant, which mean is used?

- (a) Arithmetic Mean (b) Geometric Mean
(c) Harmonic Mean (d) Weighted Geometric Mean
(a) Arithmetic Mean (b) Geometric Mean
(c) Harmonic Mean (d) Weighted Geometric Mean

181. Which is the representation of Harmonic Mean?

- (a) Mean of Reciprocal (b) Reciprocal of Mean
(c) Reciprocal of Mean of Reciprocal (d) None of the above

3.4 Geometric Mean

182. Which data set is suitable for Geometric Mean?

- (a) 1, -1, 2, 4, 6, 7 (b) 1, 2, 4, 8, 16, 32 (c) 0, 1, 2, 3, 4, 6 (d) 1, 1, 2, 3, 4, 4, 5

183. Find geometric mean: 2, 4, 8, 16

- (a) 6.65 (b) 6.56 (c) 5.66 (d) 5.56

Answer the next three questions based on the following information

The data collected in a research is this: 1, 2, 4, 8, 16, 32

184. Which measure is suitable?

- (a) Arithmetic Mean (b) Geometric Mean (c) Median (d) Mode

185. What is the arithmetic mean of the data?

- (a) 8.5 (b) 10 (c) 8 (d) 10.5

186. What is the geometric mean?

- (a) 8.5 (b) 5.66 (c) 6.55 (d) 16

3.5 Mode

187. Which of the following may be used to determine mode?
(a) Histogram (b) Frequency Curve (c) Ogive (d) Frequency Polygon
188. What is the mode the set: 7, 8, 8, 9, 9, 13, 17, 9, 8, 8
(a) 17 (b) 9
(c) 8 (d) Cannot be determined
189. What is the mode of the data set: 4, 7, 2, 4, 9, 4, 2, 9?
(a) 2 (b) 4 (c) 9 (d) 7
190. Which of the following best defines the mode of a data set?
(a) The middle value when data are arranged in order
(b) The average of all the values
(c) The value that occurs most frequently
(d) The difference between highest and lowest values
191. Find the mode of the following frequency distribution:

Value	2	3	4	5	6
Frequency	3	5	2	7	1

- (a) 3 (b) 5 (c) 6 (d) 5
192. In a symmetrical unimodal distribution, which of the following is usually true?
(a) Mean < Median < Mode (b) Mean > Median > Mode
(c) Mean = Median = Mode (d) Mode > Mean

3.6 Median

193. Which can be measured from the Ogive?
(a) Arithmetic Mean (b) Geometric Mean (c) Median (d) Mode
194. Median can be determined from the—
(a) Histogram (b) Frequency curve (c) Ogive (d) Pie Chart

3.7 Partition Values

3.8 Situation Set

Answer the next three questions based on the following information

The following table shows weekly production of milk (in liters) by different varieties of cows.

Interval	10-20	20-30	30-40	40-50	50-60	60-70
Frequency	5	12	18	25	20	10

195. What is the median?

- (a) 43 (b) 44 (c) 45 (d) 50

196. What is the lower limit of class interval for first quartile?

- (a) 10 (b) 20 (c) 30 (d) 40

197. What is the 3rd quartile?

- (a) 55.75 (b) 43.75 (c) 53.15 (d) 53.75

Answer the next two (2) questions based on the following information

Class	≤ 20	20-25	25-50	50-60	69-70	≥ 70
Frequency	5	10	10	7	5	3
Cumulative Frequency	5	15	25	32	37	40

198. How many values are between 20 and 70?

- (a) 20 (b) 32 (c) 35 (d) 37

199. Which one is the median class?

- (a) 20-25 (b) 25-50 (c) 50-60 (d) 60-70

200. What is the median of the following values: 4, 5, 2, 1, 8, 3

- (a) 1.5 (b) 2 (c) 3.5 (d) 4

Answer the next three questions as per the following information.

42 44 59 64 70 72 74 91 94 are 9 values.

201. What is the 50th percentile?

- (a) 64 (b) 70 (c) 72 (d) 71

202. Below which value lie 70 percent values?

- (a) 42 (b) 44 (c) 59 (d) 74

203. Above which value lie 30% observations?

- (a) 3rd Quartile (b) Median (c) 30th Percentile (d) 70th percentile

Answer the next three questions as per the following information.

42 44 59 64 70 72 74 91 94 are 9 values.

204. What is the median?

- (a) 64 (b) 70 (c) 72 (d) 71

205. What is the first quartile?

- (a) 42.4 (b) 44.7 (c) 51.5 (d) 64.2

206. Above which value lie 60% observations?

- (a) 70.4 (b) 72.0 (c) 74.6 (d) 66.4

3.9 Multiple Completion

207. **Inappropriate for algebraic analysis–**

- i. Median
- ii. Mode
- iii. Geometric Mean

Which one is true?

- (a) i (b) ii (c) i & ii (d) ii & iii

208. **With negative observations, which cannot be used**

- i. Arithmetic Mean
- ii. Geometric Mean
- iii. Harmonic Mean

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

209. **A good measure of central tendency -**

- i. is loosely defined
- ii. takes into consideration all values
- iii. easily understandable

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

210. **A good measure of central tendency -**

- i. is not affected by extreme values
- ii. represents the entire dataset accurately
- iii. is difficult to compute

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

211. **A good measure of central tendency -**

- i. is stable for different samples
- ii. provides a single representative value
- iii. ignores extreme values completely

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

212. **Median is –**

- i. Affected by extreme values
- ii. Rigidly defined
- iii. Suitable for open-ended distributions

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

213. **Mode is –**

- i. The most frequently occurring value
- ii. Unaffected by extreme values
- iii. Always unique in a dataset

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

214. A rate is defined as $R = \frac{c}{d}$; c and d are arbitrary numbers. If neither c or d is constant, which mean is used?

- i. Weighted Arithmetic Mean
- ii. Weighted Harmonic Mean
- iii. Harmonic Mean

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

215. What is true of harmonic mean?

- i. uses all values in the data
- ii. undefined if the any value is zero
- iii. affected by extreme values

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

216. Arithmetic Mean is –

- i. Rigidly defined
- ii. Unaffected by sample fluctuation
- iii. Suitable for algebraic analysis

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

4 Measures of Dispersion

217. What is the variance of a constant c?

- (a) 1 (b) 0 (c) -1 (d) ∞

218. What is the correct formula for the variance of a sample?

- (a) $\frac{\sum x_i^2}{n} - \frac{\bar{x}}{n}$ (b) $\frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2$ (c) $\sum x^2$ (d) $\sum 2^x$

219. Two sets of data are X = 2, 4, 6 and Y = 4, 6, 8; which has higher variance?

- (a) X
(b) Y
(c) $V(X) = V(Y)$
(d) Cannot be determined from the given information

220. Which set of data looks more scattered?

- (a) -1, 1, 2 (b) -5, 6, 7 (c) -10, 10, 30 (d) 5, 8, 20

221. Which of the following is the best measure of dispersion?

- (a) Range (b) Mean deviation
(c) Standard deviation (d) Coefficient of variation

222. What is the minimum possible value of standard deviation?

- (a) ∞ (b) -1 (c) 0 (d) 1

223. For two values, range is found to be 8. What are the values of mean deviation and standard deviation
 (a) (2,4) (b) (4,4) (c) (4,8) (d) (8,8)
224. For two values, the range is found to be 12. What are the values of mean deviation and standard deviation
 (a) (2,4) (b) (4,4) (c) (6, 6) (d) (8,8)
225. The mean and coefficient of variation of a distribution are 5 and 30%, respectively. What is the value of standard deviation?
 (a) 1.5 (b) 6.5 (c) 7.6 (d) 10.2
226. What is the correct formula for the standard deviation?
 (a) $\sqrt{\frac{1}{n} \sum (x_i - \bar{x})^2}$ (b) $\sqrt{\frac{1}{n-1} \sum (x_i - \bar{x})^2}$ (c) $\frac{1}{n} \sum (x_i - \bar{x})$ (d) $\sqrt{\sum (x_i - \bar{x})}$
227. What is the standard deviation of the first n natural numbers?
 (a) $\sqrt{\frac{(n^2-1)}{6}}$ (b) $\sqrt{\frac{(n^2-1)}{12}}$ (c) $\sqrt{\frac{n(n+1)(2n+1)}{6n}}$ (d) $\sqrt{\frac{n(n+1)}{2}}$
228. The Mean Deviation of two unequal numbers is 3. What is their range?
 (a) 1.5 (b) 3 (c) 6 (d) 12
229. What is the standard deviation of first 10 natural numbers?
 (a) 2.87 (b) 3.02 (c) 0 (d) 2.78
230. Which measure is unit-free?
 (a) Range (b) Mean deviation
 (c) Standard deviation (d) Coefficient of variation
231. Which measure is suitable for an open-ended distribution?
 (a) Range (b) Mean deviation (c) Standard deviation (d) Quartile deviation

4.1 Situation Set

Answer the next two questions based on the following information.

The temperatures (in $^{\circ}C$ of two cities in a country are 30 and 35.

232. What is their Mean deviation?
 (a) 1.2 (b) 2.5 (c) 3.0 (d) 5.5
233. What is the coefficient of variation?
 (a) 2.7% (b) 8.3% (c) 5.8% (d) 7.7%

4.2 Multiple Completion

234. Dispersion has application in measuring —

- i. regression
- ii. correlation
- iii. range

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

235. **Standard deviation** —

- i. depends on all values
- ii. is not affected by outliers
- iii. can be analyzed algebraically

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

5 Moments, Skewness, and Kurtosis

5.1 Moments

236. **What is the purpose of moments?**

- (a) To find the center of data
- (b) To find dispersion
- (c) To find relationship between skewness and kurtosis
- (d) To uniquely identify a distribution

237. **The moments around the origin are called –**

- (a) Central moments (b) Raw moments
(c) First raw moment (d) Measures of dispersion

238. **The Range of Karl Pearson's measure of skewness –**

- (a) (0, 1) (b) (-1, 1) (c) (-3, 3) (d) (0, ∞)

239. **Which is not a type of Moments**

- (a) Central Moments (b) Raw Moments (c) Corrected Moments (d) Rectified Moments

240. **The second moment around w is –**

- (a) $\frac{\sum(x_i - \bar{x})^n}{w}$ (b) $\frac{\sum(x_i - \bar{x})^2}{w}$ (c) $\frac{\sum(x_i - w)^2}{n}$ (d) $\frac{\sum(x_i - w)^n}{2}$

241. **Which relationship is correct?**

- (a) $\mu'_1 = \bar{x} + a$ (b) $\mu'_1 = \bar{x} - a$ (c) $\mu'_2 = \bar{x} + a$ (d) $\mu_1 = \bar{x} - a$

242. **What is formula of rth raw moment for grouped data about a?**

- (a) $\frac{\sum f_i(x_i - a)^r}{n}$ (b) $\frac{\sum f_i(x_i - \bar{x})^r}{n}$ (c) $\frac{\sum(x_i - a)^r}{n}$ (d) $\frac{\sum(x_i + a)^r}{n}$

243. **Which quantity uniquely characterizes a distribution?**

- (a) Median (b) Quantile (c) Moments (d) Trend

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

244. **Which can be used to measure dispersion?**

- (a) μ'_2 (b) μ_1 (c) μ_2 (d) μ'_1

245. **The formula of coefficient of variance (CV) is –**

- (a) $\frac{\sqrt{\mu_2}}{n} \times 100$ (b) $\frac{\mu_2}{\mu_1} \times 100$ (c) $\frac{\sqrt{\mu_2}}{\bar{x}} \times 100$ (d) $\frac{\mu_3}{\sigma} \times 100$

246. **First moment around zero is –**

- (a) 0 (b) 1 (c) -1 (d) Arithmetic Mean

247. Which moment is equal to zero?
- (a) First raw moment around 1 (b) Second central moment
(c) First central moment (d) Second raw moment around 0
248. Which might have a negative value?
- (a) μ_4 (b) μ_3 (c) μ'_2 (d) μ_2
249. 2nd Central Moment is –
- (a) $\mu_2 - \mu'^2_1$ (b) $\mu_2 + \mu'_1$ (c) $\mu_2 - \mu'^2_1$ (d) $\mu'_2 - \mu'^2_1$
250. First central moment is equal to –
- (a) 1 (b) 0 (c) -1 (d) $\bar{x} - a$
251. First moment around a is equal to –
- (a) 1 (b) 0 (c) -1 (d) $\bar{x} - a$
252. The first raw moment about 3 is -5. What is the value of arithmetic mean?
- (a) 2 (b) -2 (c) 0 (d) 8
253. The first raw moment about 4 is -4. What is the value of arithmetic mean?
- (a) 2 (b) -2 (c) 0 (d) 8
254. The first raw moment about 0 is 2. What is the value of arithmetic mean?
- (a) 2 (b) -2 (c) 0 (d) 8
255. The arithmetic mean of a variable is 4. What is the first raw moment around 2?
- (a) 2 (b) -2 (c) 0 (d) 8
256. The arithmetic mean of a variable is 10. What is the first raw moment around 0?
- (a) 10 (b) -2 (c) 0 (d) 8
257. The arithmetic mean of a variable is 2.6. What is the first raw moment around 6?
- (a) 2.2 (b) -3.4 (c) 0.1 (d) 1.8
258. If the values in a dataset have mean 4.8, what is the first moment about the mean?
- (a) 0 (b) 4.8 (c) 1.0 (d) -4.8
259. The mean of a variable is 3.2. Find the first raw moment around 0.
- (a) 3.2 (b) -3.2 (c) 0 (d) 1.2
260. The first raw moment around 0 of a data set is 5. What is the arithmetic mean of the data?
- (a) 3 (b) 4 (c) 5 (d) 6
261. The first raw moment around 5 of a data set is 15. What is the arithmetic mean of the data?
- (a) 8 (b) 20 (c) 12 (d) 15
262. The first raw moment around 3 of a data set is 18. What is the arithmetic mean of the data?
- (a) 6 (b) 17 (c) 28 (d) 21

263. The first raw moment around 10 of a data set is 50. What is the arithmetic mean of the data?

- (a) 52 (b) 24 (c) 60 (d) 40

264. Moments can be–

- i. positive
ii. not negative
iii. positive or negative

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

265. The second central moment of the natural numbers 1 through 12 is –

- (a) 12.93 (b) 11.92 (c) 10.94 (d) 12.60

266. The second raw moment around zero of the natural numbers 5 through 15 is –

- (a) 120 (b) 115 (c) 100 (d) 110

5.2 Skewness


267. What kind of distribution is clustered around the mean?

- (a) Left-skewed (b) Symmetrical (c) Right-skewed (d) None of the above

268.

- (a) (b) Choice (c) Choice (d) Choice

269. The following graph is an example of –




- (a) Positive Skew (b) Negative Skew (c) No Skew (d) Not detectable

270. For a symmetrical distribution, what is the value of β_1 ?

- (a) 0 (b) 1 (c) -1 (d) ∞

Answer the next ? questions based on the following information



271. The curve (a) is an example of

- (a) Positive Skew (b) Negative Skew (c) No Skew (d) Not detectable

272. The curve (b) is an example of

- (a) Positive Skew (b) Negative Skew (c) No Skew (d) Not detectable

273. In Image (b), what is denoted by 4th value?
 (a) Mean (b) Median (c) Mode (d) All of the above
274. In Image (c), what is in 6th value?
 (a) Mean (b) Median (c) Mode (d) None of the above
275. What is the value corresponding to the position 3?
 (a) Mean (b) Median (c) Mode (d) None of the above
276. What is the value corresponding to the position 7?
 (a) Mean (b) Median (c) Mode (d) None of the above
277. If $\gamma_1 > 0$, the data is -
 (a) Negatively skewed (b) Positively skewed (c) Symmetric (d) Uncertain
278. Which relationship is correct?
 (a) $M_o = 2Me - \bar{x}$ (b) $M_o = 3Me - \bar{x}$ (c) $M_o = 3Me - 2\bar{x}$ (d) $M_o = 2Me - 3\bar{x}$
279. Characteristics of a skewed distributon are -
 i. $Mean \neq Median \neq Mode$
 ii. Differences of upper and lower quartiles from median are unequal
 iii. Frequency curve is asymmetric
280. In a distribution, $\mu_2 = 25, \mu_3 = 20$, and $\mu_4 = 2200$; the distribution is -
 (a) Negativelky skewed (b) leptokurtic (c) Platykurtic (d) Symmetric
281. For a data, $Q_3 = 41.6, Q_1 = 17.2, Median = 29$, & $AM = 30$; What is Coefficient of skewness?
 (a) 24.4 (b) 1 (c) 0.03 (d) 29.45
282. In case of positive skewness, which one is correct?
 (a) $Mean > Median > Mode$ (b) $Mean < Median < Mode$
 (c) $Mean = Median = Mode$ (d) $Mean > Median < Mode$
283. For a symmetrical distribution, $\beta_1 =$ —
 (a) 1 (b) -1 (c) 0 (d) 3
284. $\sqrt{\beta_1} = -0.23$ implies—
 (a) Left Skew (b) Symmetry (c) Right Skew (d) Mesokurtic
285. $\gamma_1 = 0.43$ implies—
 (a) Left Skew (b) Symmetry (c) Right Skew (d) Mesokurtic
286. $\gamma_1 = 0.0001$ implies—
 (a) Left Skew (b) Symmetry (c) Right Skew (d) Mesokurtic
287. First 3 moments about 2 are 1, 2 and 8, respectively. What is the arithmetic mena?
 (a) 1 (b) 2 (c) 3 (d) 4
288. What is the second central moments of first 10 natural numbers?
 (a) 9.90 (b) 9.09 (c) 8.25 (d) 5.67
289. Frequencies of low and high values are smaller in - distribution
 (a) Positively skewed (b) Negatively skewed (c) Symmetric (d) Mesokurtic

290. **Frequencies of higher values are smaller and of low values are higher in – distribution**

- (a) Positively skewed (b) Negatively skewed (c) Symmetric (d) Mesokurtic

291. **Frequencies of higher values are higher and of low values are lower in – distribution**

- (a) Positively skewed (b) Negatively skewed (c) Symmetric (d) Mesokurtic

292. **In a positively-skewed distribution–**

- i. Frequencies of higher values are lower
- ii. Frequencies of low values are higher
- iii. Frequencies of higher values are higher

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

293. **In a negatively-skewed distribution–**

- i. Frequencies of higher values are lower
- ii. Frequencies of low values are lower
- iii. Frequencies of higher values are higher

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

294. **In a symmetric distribution–**

- i. Frequencies of higher values are lower
- ii. Frequencies of low values are higher
- iii. Frequencies of low values are lower

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

295. **Which formula is correct for determining skewness?**

- (a) $\gamma_1 = \sqrt{\frac{\mu_3^2}{\mu_2^3}}$ (b) $\gamma_1 = \sqrt{\beta_1^2}$ (c) $\gamma_1 = \sqrt{\frac{\mu_3}{\mu_2^3}}$ (d) $\frac{\mu_2}{\sqrt{\mu_3^2}}$

5.2.1 Situation Set

Answer the next THREE questions based on the following information

The arithmetic mean, Pearson's coefficient of skewness, and coefficient of variation of a distribution are 50, -0.4, and 40 %, respectively.

296. **What is the standard deviation?**

- (a) 50 (b) 20 (c) 30 (d) 40

297. **The Mode is –**

- (a) 45 (b) 50 (c) 60 (d) 58

298. **The Median is**

- (a) 52.67 (b) 5 (c) 20 (d) 10.39

Answer the next two questions based on the following information

The first and second moments of a distribution around 2 are 13 and 195, respectively.

299. What is the arithmetic mean?

- (a) 15 (b) 12 (c) 11 (d) 26

300. What is its variance?

- (a) 20 (b) 21 (c) 26 (d) 25

5.2.2 Multiple Completion

301. In a negatively skewed distribution –


- i. there is a long tail on the left
- ii. median is greater than mean
- iii. high values have frequencies

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

5.3 Kurtosis

302. Which curve is platykurtic?



- (a) A (b) B (c) C (d) None

303. How many types of kurtosis are there?

- (a) 2 (b) 3 (c) 4 (d) 5

304. The standard deviation of a mesokurtik distribution is 2. What is the value of the 4th central moment?

- (a) 4 (b) 8 (c) 16 (d) 48

305. $\beta_2 = \sqrt{9}$ implies data are–

- (a) Leptokurtic (b) Platykurtic (c) Mesokurtic (d) Symmetric

306. $\beta_2 = 4$ implies data are–

- (a) Leptokurtic (b) Platykurtic (c) Mesokurtic (d) Symmetric

307. $\beta_2 = 3$ implies data are–

- (a) Leptokurtic (b) Platykurtic (c) Mesokurtic (d) Symmetric

308. $\beta_2 = 1$ implies data are–

- (a) Leptokurtic (b) Platykurtic (c) Mesokurtic (d) Symmetric

309. The relationship between β_2 and γ_2 is –

- (a) $\beta_2 = \gamma_2 - 3$ (b) $\gamma_2 = \beta_2 - 3$ (c) $\gamma_2 = 3\beta_2$ (d) $\gamma_2 = \frac{\beta_2}{3}$

310. For a mesokurtik distribution, $\beta_2 = --$

- (a) 0 (b) -3 (c) 3 (d) 1

311. What is the relationship between γ_2 and β_2 ?

- (a) $\gamma_2 = \beta_2 + 3$ (b) $\gamma_2 = 2\beta_2 - 3$ (c) $\gamma_2 = \beta_2 - 1$ (d) $\gamma_2 = \beta_2 - 3$

5.4 Misc

312. What is formula of the left inner fence for a box and whisker plot?

- (a) $Q_1 - 1.5 \times IQR$ (b) $Q_3 + 1.5 \times IQR$ (c) $Q_1 - 3 \times IQR$ (d) $Q_3 + 1.5 \times IQR$

313. What is the formula of IQR?

- (a) $IQR = Q_3 + Q_1$ (b) $IQR = Q_3 - Q_1$ (c) $IQR = 2Q_3 - Q_1$ (d) $IQR = \frac{Q_3 - Q_1}{2}$

314. Which is not used in constructing Box & Whisker Plot?

- (a) Mode (b) X_L (c) $Q_1 \& Q_3$ (d) $Q_1, Q_2 \& Q_3$

315. In a symmatric distribution–

- i. Arithmetic Mean = Mode = Median
ii. $Q_2 - Q_1 = Q_3 - Q_2$
iii. $Q_1 - X_L = X_H - Q_3$

Which one is true?

- (a) i & ii (b) ii & iii (c) i & iii (d) i, ii & iii

5.5 Box and Whisker Plot

316. The following values represent the quartiles of a data set:

- $Q_1 = 25$
- $Q_2 = 50$
- $Q_3 = 75$

What is the interquartile range (IQR)?

- (a) 25 (b) 50 (c) 75 (d) 100

317. In a box and whisker plot, the following statements hold true:

- i. The length of the box represents the interquartile range (IQR).
ii. The whiskers extend from the minimum to the maximum data values.
iii. The median is represented by the top of the box.

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

5.6 Five Number Summary

318. In a given data set, the following values are recorded:

- i. The interquartile range is the difference between Q3 and Q1.
- ii. The median is always equal to the mean.
- iii. The maximum value in the five-number summary is the largest data point.

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

319. The five-number summary of a data set consists of the following:

- i. Minimum value
- ii. Maximum value
- iii. Median (Q2), First Quartile (Q1), Third Quartile (Q3)

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

320. Which is not included in five number summary?

- (a) Arithmetic Mean (b) X_H (c) Q_2 (d) Q_3

6 Correlation and Regression

6.1 Correlation

321. Who proposed the formula of correlation coefficient?

- (a) R. A. Fisher (b) Bowley (c) Spearman (d) Karl Pearson

322. Which of the following has the strongest correlation?

- (a) -0.92 (b) 0.67 (c) 0.50 (d) 0.91

323. The lowest possible value of the correlation coefficient —

- (a) 1 (b) 0 (c) $-\infty$ (d) -1

324. The linear association between two random variables is called —

- (a) Correlation (b) Regression (c) Randomness (d) Regularity

325. Which measures the strength of linear association between two random variables?

- (a) Correlation (b) Regression (c) Correlation coefficient (d) Regression coefficient

326. Karl Pearson's method of determining the strength of correlation is not applicable for —

- (a) Qualitative variable (b) Quantitative variable (c) Discrete variable (d) Continuous variable

327. For two independent variables, the value of the correlation coefficient is —

- (a) -1 (b) 1 (c) ∞ (d) 0

328. Two variables having changes in same direction at same rates display —

- (a) Perfect negative correlation (b) Partial positive correlation
(c) Perfect positive correlation (d) Partial negative correlation

329. **Two variables having changes in opposite direction at same rates display —**
 (a) Perfect negative correlation (b) Partial positive correlation
 (c) Perfect positive correlation (d) Partial negative correlation
330. **Two variables having changes in same direction at different rates display —**
 (a) Perfect negative correlation (b) Partial positive correlation
 (c) Perfect positive correlation (d) Partial negative correlation
331. **Two variables having changes in opposite direction at different rates display —**
 (a) Perfect negative correlation (b) Partial positive correlation
 (c) Perfect positive correlation (d) Partial negative correlation
332. **When a variable is changing, but another is not affected, it is called**
 (a) Perfect negative correlation (b) Partial positive correlation
 (c) Perfect positive correlation (d) Zero correlation
333. **TEXT**
 (a) Choice (b) Choice (c) Choice (d) Choice
334. **Which of the following coefficients indicates strong negative correlation?**
 (a) -0.5 (b) -2 (c) 0.67 (d) -0.94
335. **Which of the following has the strongest correlation?**
 (a) -0.5 (b) -2 (c) 0.67 (d) -0.94
336. **If $b_1 = 0.25$ and $b_2 = -0.43$, which implies higher impact of the independent variable?**
 (a) b_1 (b) b_2 (c) Cannot be determined (d) Equal impact

6.2 Situation Set

Answer the next two questions based on the following information

A study was conducted to find the impact of study hour on students' GPA and the following was found:

$$\sum (x_i - \bar{x})(y_i - \bar{y}) = 30, \sum (x_i - \bar{x})^2 = 45, \text{ and } \sum (y_i - \bar{y})^2 = 55$$

337. **What is the value of correlation coefficient?**
 (a) 0.50 (b) 0.60 (c) -0.60 (d) -0.50
338. **What is the value of b_{yx} ?**
 (a) 0.58 (b) -0.67 (c) 0.67 (d) -1.75

7 Time Series

339. **Which is not a time series data?**
 (a) Number of calls received per week (b) No. of road accidents on different days
 (c) No. of earthquakes in different regions (d) No. of particles decayed in each second

340. Which is not a time series data?

- (a) Daily closing prices of a stock (b) Annual temperature records of a city
(c) Number of students in a each class (d) Number of visitors to a website each day

341. Which is an example of time series data?

- (a) Number of calls received by a call center each month
(b) Height of children at different ages
(c) Total salary of all employees at a company
(d) Population of different countries in 2020

342. Which is a type of trend?

- i. Linear trend
ii. Non-linear trend
iii. Cyclic trend

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

343. Which can measure trend most precisely?

- (a) Graphical method (b) Semi-average method
(c) Moving average method (d) Quarter-average method

344. Which is the multiplicative time series model?

- (a) $Y_t = T_t \times S_t \times C_t \times R_t$ (b) $Y_t = T_t \times D_t \times C_t \times R_t$
(c) $Y_t = T_t \times P_t \times C_t \times R_t$ (d) $Y_t = T_t \times G_t \times C_t \times R_t$

345. In additive model, in the long run, $\sum R_t =$ --

- (a) 0 (b) 1 (c) Undefined (d) Infinity

346. In multiplicative time series model, in the long run, $\sum R_t =$ --

- (a) 0 (b) 1 (c) Undefined (d) Infinity

Answer the next two questions based on the following information

Commodity wise export shipments (In million US\$) of Frozen and live fish in Bangladesh are given below.

Months	2022-23 (July-Dec)	2023-24 (Jan-Jun)	2022-23 (July-Dec)
Amount	246.38	175.19	215.13

Table 1: Source:BB

347. Which component of time series is most evident?

- (a) Irregular variation (b) Cyclic variation (c) Trend (d) Seasonal variation

348. Which value is most probable in the next period?

- (a) 200 (b) 190 (c) 130 (d) 220

349. A linear trend goes along a --

- (a) a curved line (b) a wave (c) straight line (d) circle

350. Which of the following is an example of seasonal variation in a time series?
- (a) Increase in ice cream sales during summer (b) Rising fuel prices over decades
(c) Stock market crash (d) Unemployment rate changes due to war
351. Which business is most likely to experience strong seasonal variation in its sales?
- (a) A supermarket (b) A toy store (c) A furniture store (d) A gas station
352. Which of the following is an example of cyclic variation in a time series?
- (a) Boom and recession phases in an economy
(b) Increase in electricity consumption during summer
(c) High demand for umbrellas during the rainy season
(d) Sudden decline in stock prices due to a pandemic
353. Which of the following is an example of a trend in a time series?
- (a) Gradual increase in global average temperature
(b) Increase in ice cream sales during summer
(c) Fluctuations in stock prices due to news events
(d) Sudden drop in airline bookings due to a storm
354. Which type of trend is usually observed in a country's population growth over several decades?
- (a) Upward trend (b) Downward trend (c) Seasonal trend (d) Cyclic trend
355. Which of the following best represents a downward trend in a time series?
- (a) Declining birth rates in a country over several decades
(b) Increase in online shopping during holiday seasons
(c) Fluctuations in stock market prices
(d) Sudden rise in fuel prices due to a crisis
356. Which factor is most likely to contribute to an upward trend in a company's annual revenue?
- (a) Improved marketing strategies over time (b) Seasonal discounts and promotions
(c) Short-term fluctuations in customer demand (d) Unpredictable supply chain disruptions
357. Which factor is most likely to cause cyclic variation in a time series?
- (a) Festive shopping trends (b) Long-term business cycles
(c) Daily fluctuations in temperature (d) Random fluctuations in demand
358. A non-linear trend goes along a –
- (a) a curved line (b) a wave (c) a cubic pattern (d) Any of the above
359. Which measure of trend is subjective?
- (a) Semi-average method (b) Graphical method
(c) Moving average method (d) None of the above
- Answer the next THREE questions based on the following information**
360. What is the second value of semi-average method?
- (a) 85.40 (b) 90.37 (c) 91.73 (d) 89.78

Year	2016	2017	2018	2019	2020	2021	2022	2023
USD Exchange Rate	78.35	79.49	82.87	83.26	84.60	84.37	85.80	106.70

Table 2: Source–Investing.com

361. **What kind of a trend do the data have?**

- (a) Upward (b) Downward
(c) Both upward & downward (d) No trend

362. **Which component of time series is visible in the later part of the data?**

- (a) Seasonal Variation (b) General Trend (c) Irregular Variation (d) Cyclic Variation

Answer the next THREE questions based on the following information

Year	2015	2016	2017	2018	2019	2020	2021	2022
Average Temperature (°C)	22.5	23.0	24.2	24.5	25.0	25.5	26.0	27.0

Table 3: Source–National Weather Service

363. **What is the second value of the semi-average method?**

- (a) 25.75 (b) 26.00 (c) 25.88 (d) 24.29

364. **What kind of trend do the data show?**

- (a) Upward (b) Downward
(c) Both upward & downward (d) No trend

365. **Which component of the time series is most prominent in the data?**

- (a) Seasonal Variation (b) General Trend (c) Irregular Variation (d) Cyclic Variation

Answer the next THREE questions based on the following information

Year	2016	2017	2018	2019	2020	2021	2022	2023
USD Exchange Rate	78.35	79.49	82.87	83.26	84.60	84.37	85.80	106.70

Table 4: Source–Investing.com

366. **What is the second value of semi-average method?**

- (a) 85.40 (b) 90.37 (c) 91.73 (d) 89.78

367. **What kind of a trend do the data have?**

- (a) Upward (b) Downward
(c) Both upward & downward (d) No trend

368. **Which component of time series is visible in the later part of the data?**

- (a) Seasonal Variation (b) General Trend (c) Irregular Variation (d) Cyclic Variation

Answer the next THREE questions based on the following information

369. **What is the semi-average for the second period of the data?**

- (a) 160 (b) 165 (c) 180 (d) 190

370. **Which type of trend do these rainfall data indicate?**

- (a) Increasing (b) Decreasing (c) No trend (d) Fluctuating

Month	January	February	March	April	May	June	July	August
Rainfall (mm)	150	120	180	200	160	140	170	190

Table 5: Source: Meteorological Department

371. What is the primary variation component observed in the data?

- (a) Seasonal Variation (b) Trend Variation (c) Cyclic Variation (d) Irregular Variation

372. Time Series has how many components?

- (a) 2 (b) 3 (c) 4 (d) 5

373. Which component involves period more than one (01) year?

- (a) Seasonal Variation (b) Cyclic Variation (c) Irregular Variation (d) Random Variation

374. Which one is not a component of Time Series

- (a) Seasonal Variation (b) Cyclic Variation (c) General Trend (d) Regular Variation

375. A company is constantly getting greater revenue than previous year; this is–

- (a) Seasonal Variation (b) General Trend (c) Irregular Variation (d) Cyclic Variation

376. Which is not a method of finding general trend?

- (a) Graphical Method (b) Moving Average (c) Semi-Average (d) Moving Median

Answer the next two questions based on the following table:

Year	2007	2008	2009	2010	2011	2012
Sales	5	35	34	40	42	204

377. In Semi-Average method, what is the 2nd average?

- (a) 74 (b) 24.67 (c) 95.33 (d) 28

378. What is the last value of 3-yearly moving average?

- (a) 93.55 (b) 95.53 (c) 95.33 (d) 59.33

379. Which component of time series is affected by economic changes due to war?

- (a) Trend (b) Seasonal Variation (c) Irregular Variation (d) Cyclic Variation

380. Which component of a time series captures long-term upward or downward movement?

- (a) Trend (b) Seasonal Variation (c) Irregular Variation (d) Cyclic Variation

381. Which time series component represents fluctuations occurring at regular intervals within a year?

- (a) Trend (b) Seasonal Variation (c) Irregular Variation (d) Cyclic Variation

382. Which component of time series is affected by economic changes during a recession?

- (a) Trend (b) Seasonal Variation (c) Irregular Variation (d) Cyclic Variation

383. Which component of time series is most likely to be impacted by weather conditions like a monsoon season?

- (a) Trend (b) Seasonal Variation (c) Irregular Variation (d) Cyclic Variation

Year	2016	2017	2018	2019	2020
Car Sales	1200	1500	1700	1600	1800

384. Which component of time series would be influenced by government policy changes such as tax reforms?

- (a) Trend (b) Seasonal Variation (c) Irregular Variation (d) Cyclic Variation

Answer the next three questions based on the following table:

385. What is the first value of the 2-yearly moving average?

- (a) 1350 (b) 1300 (c) 1400 (d) 1250

386. What is the last value of the 3-yearly moving average?

- (a) 1600 (b) 1670 (c) 1630 (d) 1750

387. What is the semi-average for the first period of the data?

- (a) 1350 (b) 1400 (c) 1450 (d) 1300

388. Demand for warm clothes is higher in winter season and less in summer. Which component of time series deals with this change?

- (a) Trend (b) Seasonal Variation (c) Irregular Variation (d) Cyclic Variation

389. Death rates of a country for 7 years are given below:

Year	2009	2010	2011	2012	2013	2014	2015
Rate	5	7	6	8	7	12	13

In semi-average method, which year will be excluded?

- (a) 2012 (b) 2013 (c) 2015 (d) 2009

390. Which component of time series represents a natural disaster?

- (a) Seasonal Variation (b) General Trend (c) Irregular Variation (d) Cyclic Variation

391. How many models of time series are there to combine the components?

- (a) 2 (b) 3 (c) 4 (d) 5

392. Which one reflects an irregular variation?

- (a) Fluctuation in production due to war (b) Price hike due to famine
(c) Rise of Temperature to drought (d) Any of the above

7.1 Situation Set

Answer the next three questions based on the following table:

393. Death rates of a country for 7 years are given below:

Year	2009	2010	2011	2012	2013	2014	2015
Rate	5	7	6	8	7	12	13

In semi-average method, what is the first average?

- (a) 5 (b) 7 (c) 6 (d) 8

394. What is the first value of the 2-yearly moving average?

- (a) 5 (b) 6 (c) 7 (d) 8

395. What is the last value of the 3-yearly moving average?

- (a) 11.10 (b) 9.68 (c) 10.65 (d) 10.67

Answer the next three questions based on the following table:

The following table shows the population growth rate (in percentage) of a city over seven years.

Year	2015	2016	2017	2018	2019	2020	2021
Rate (%)	2.5	2.7	3.1	3.6	3.9	4.2	4.5

396. What is the average population growth rate over the 7 years?

- (a) 3.2% (b) 3.5% (c) 3.6% (d) 3.8%

397. What is the second value in the 3-yearly moving average?

- (a) 2.8% (b) 3.1% (c) 3.3% (d) 3.5%

398. Using the semi-average method, what is the second average?

- (a) 3.6% (b) 3.7% (c) 3.8% (d) 4.0%

Answer the next three questions based on the following table:

The following table shows the annual rainfall (in cm) recorded in a region over seven years.

Year	2010	2011	2012	2013	2014	2015	2016
Rainfall (cm)	85	90	88	92	95	100	105

399. What is the median annual rainfall for the given years?

- (a) 90 cm (b) 92 cm (c) 93 cm (d) 95 cm

400. What is the first value of the 2-yearly moving average?

- (a) 86.5 cm (b) 87 cm (c) 88.5 cm (d) 89 cm

401. Using the semi-average method, what is the first average?

- (a) 88 cm (b) 89 cm (c) 90 cm (d) 91 cm

Answer the next three questions based on the following table:

The following table shows the average monthly temperature (in °C) recorded in a city over seven months.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul
Temperature (°C)	12	14	18	22	26	30	32

402. What is the mean temperature over the given months?

- (a) 19.5°C (b) 20.5°C (c) 21.5°C (d) 22.5°C

403. What is the third value in the 3-monthly moving average?

- (a) 16°C (b) 18°C (c) 20°C (d) 22°C

404. Using the semi-average method, what is the second average temperature?

- (a) 24°C (b) 25°C (c) 26°C (d) 27°C

Answer the next three questions based on the following table:

The following table shows the monthly sales revenue (in thousand dollars) of a store over seven months.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul
Revenue (000\$)	50	55	60	70	75	80	85

405. Which month had the highest sales revenue?

- (a) May (b) Jun (c) Jul (d) Apr

406. What is the first value of the 2-monthly moving average?

- (a) 52.5 (b) 55 (c) 57.5 (d) 60

407. Using the semi-average method, what is the first average revenue?

- (a) 57.5 (b) 55 (c) 62.5 (d) 65

7.2 Multiple Completion

408. Which of the following are components of a time series?

- i. Trend
- ii. Seasonal Variation
- iii. Correlation

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

409. Which statements about time series models are correct?

- i. The additive model adds all the components
- ii. The multiplicative model also contains some additions.
- iii. Additive and multiplicative models produce identical forecasts.

Which one is correct?

- (a) ii (b) iii (c) i (d) i, ii and iii

410. Which of the following are methods of estimating trend in time series?

- i. Moving Average Method
- ii. Sem-average method
- iii. Simple Random Sampling

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

8 Published Statistics in Bangladesh

411. Limitations of published statistics in Bangladesh are –

- i. Wrong data collection method
- ii. Insufficient data
- iii. Lack of proper training

Which one is correct?

- (a) i and ii (b) i and iii (c) ii and iii (d) i, ii and iii

412. How many sources of published statistics are there in Bangladesh?

- (a) 2 (b) 3 (c) 4 (d) 6

413. Bangladesh Bureau of Statistics collect –

- (a) Official statistics (b) Non-official statistics(c) Semi-official statistics(d) None of the above

414. Which statistics are published by an NGO?

- (a) Official statistics (b) Non-official statistics(c) Semi-official statistics(d) None of the above

415. The primary source of official statistics in Bangladesh is –

- (a) WHO (b) BBS (c) CPD (d) UNDP

416. Which statistics are typically published by NGOs like World Wildlife Fund (WWF)?

- (a) Official statistics (b) Non-official statistics(c) Semi-official statistics(d) None of the above

417. Which organization typically publishes non-official statistics in the field of health?

- (a) UNICEF (b) World Health Organization (WHO)
(c) World Bank (d) United Nations (UN)

418. In Bangladesh, a census is usually done every – years

- (a) 20 (b) 15 (c) 10 (d) 12

419. Population census is –

- (a) Official statistics (b) Non-official statistics(c) Semi-official statistics(d) None of the above

420. In Bangladesh, which ministry present the budget?

- (a) Planning (b) Education (c) Finance (d) Agriculture

Answer Key:

1. (d) R.A. Fisher
2. (d) Database creation
3. (d) Red blood cells in a person's body
4. (c) Stars in the Milky Way
5. (b) Fish in the Pacific Ocean
6. (a) i and ii
7. (b) $\sum_{i=1}^{20} cx_i = nc \sum_{i=1}^{20} x_i$
8. (d) Regression
9. (c) Correlation
10. (c) Regression analysis
11. (b) Water molecules in the ocean
12. (a) Books in a school library
13. (b) Grains of sand on a beach
14. (d) Ordinal
15. (b) Ordinal
16. (c) Interval
17. (a) Nominal
18. (a) $y_i = \frac{x_i}{a}$
19. (c) 150
20. (a) 100
21. (c) 80
22. (a) 50
23. (c) Sample
24. (b) $b \sum_{i=1}^n x_i$
25. (c) 4
26. (d) Success rate
27. (c) Ratio scale
28. (d) Ratio
29. (d) Grade in a subject
30. (b) Number of cars in a parking lot
31. (b) Number of students in a class
32. (b) Number of books on a shelf
33. (a) Shoes sizes available in a store
34. (a) Grades on a multiple-choice test (A, B, C, D)
35. (a) Outcomes of rolling a die
36. (a) Counts of people in a room
37. (a) Number of languages spoken by a person
38. (d) No. of particles in atoms
39. (c) 206
40. (d) 122
41. (b) 65
42. (c) 42
43. (c) 54
44. (d) 45
45. (d) 84
46. (c) 8
47. (b) 62
48. (b) 6
49. (c) 90
50. (d) 435
51. (c) 2612
52. (d) 7264
53. (c) 344
54. (b) 330
55. (c) 24
56. (d) 50
57. (a) 108
58. (b) 174
59. (a) i and ii
60. (a) Temperature
61. (c) Gender
62. (c) Educational Level
63. (a) Temperature
64. (c) Ratio scale
65. (d) Grade in a subject
66. (a) $\prod x_i^2$
67. (b) Continuous variable
68. (c) Mean monthly income in a city is 60,000 taka
69. (d) 13
70. (c) 93
71. (c) 99
72. (d) 119
73. (d) -34
74. (a) Room no.
75. (d) No. of member in a family
76. (c) Nominal
77. (b) 155
78. (a) 225
79. (c) 37
80. (b) 33
81. (a) 20
82. (b) 504
83. (c) 82
84. (a) 71
85. (d) 24
86. (c) 66
87. (a) 74
88. (b) 74
89. (c) 476
90. (a) 61
91. (c) 45
92. (d) 115
93. (d) 2
94. (a) Data
95. (a) Primary data
96. (c) $\theta_i = \frac{f_i}{N} \times 360$
97. (d) John Tukey
98. (b) Sample

99. (a) $K = 1 + 3.322 \log N$ 124. (a) Quartiles are well defined 149. (d) 7.07 174. (b) $n + 1$
100. (b) Bar Diagram 125. (b) Mode 150. (c) 80 175. (b) $\frac{n+1}{2}$
101. (c) 36 126. (c) Range 151. (d) 36 176. (c) 32.00
102. (b) 45 127. (b) When all the values are equal 152. (c) 165 177. (a) $\frac{n}{\sum_{i=1}^n \frac{f_i}{x_i}}$
103. (a) 44% 128. (c) Geometric Mean 153. (c) 7 178. (c) 14.39
104. (d) 6 129. (d) 5 154. (c) Multiplied by 3 and then increased by 4 179. (c) Harmonic Mean
105. (b) 31 130. (d) Mode 155. (b) 6 180. (a) Arithmetic Mean
106. (a) 31 131. (b) Geometric Mean 156. (b) 15 180. (c) Harmonic Mean
107. (d) 29 132. (c) 7.5 157. (a) 4 times 181. (c) Reciprocal of Mean of Reciprocal
108. (a) 50 133. (b) 8 158. (b) $\bar{x}$ 182. (b) 1, 2, 4, 8, 16, 32
109. (b) 45 134. (d) Mode 159. (a) $\bar{x}$ 183. (c) 5.66
110. (b) 75% 135. (d) 110th Percentile 160. (a) 40 184. (b) Geometric Mean
111. (a) 55 136. (a) $\sum_{i=1}^n (X_i - \text{Median})^2$ 161. (c) 5.5 185. (d) 10.5
112. (c) 65 137. (b) Geometric Mean 162. (b) 13 186. (b) 5.66
113. (c) 60% 138. (a) All values are equal 163. (c) 109 187. (a) Histogram
114. (d) 0.35 139. (b) Median 164. (c) 109 188. (c) 8
115. (d) Ogive 140. (b) Geometric Mean 165. (a) 26 189. (b) 4
116. (a) i and ii 141. (b) Geometric Mean 166. (d) 10 190. (c) The value that occurs most frequently
117. (a) i and ii 142. (c) Median 167. (a) 20 191. (d) 5
118. (a) i and iii 143. (c) Median 168. (b) 20 192. (c) Mean = Median = Mode
119. (a) i and iii 144. (b) Harmonic mean 169. (a) 88.36 193. (c) Median
120. (d) i, ii and iii 145. (d) Mode 170. (a) 0 194. (c) Ogive
121. (a) i and ii 146. (b) $AM \times HM = GM^2$ 171. (a) $\bar{X} = \frac{\sum f_i x_i}{\sum f_i}$ 195. (b) 44
122. (a) i and ii 147. (b) 6.67 172. (c) 47 196. (c) 30
123. (a) i and ii 148. (b) 18 173. (b) n 197. (d) 53.75
198. (b) 32

199. (b) 25-50
 200. (c) 3.5
 201. (b) 70
 202. (d) 74
 203. (d) 70th percentile
 204. (b) 70
 205. (c) 51.5
 206. (c) 74.6
 207. (c) i & ii
 208. (c) ii and iii
 209. (c) ii and iii
 210. (a) i and ii
 211. (a) i and ii
 212. (b) i and iii
 213. (a) i and ii
 214. (a) i and ii
 215. (a) i and ii
 216. (b) i and iii
 217. (b) 0
 218. (b) $\frac{\sum x_i^2}{n} - \left(\frac{\sum x_i}{n}\right)^2$
 219. (c) $V(X) = V(Y)$
 220. (c) -10, 10, 30
 221. (c) Standard deviation
 222. (c) 0
 223. (b) (4,4)
224. (c) (6, 6)
 225. (a) 1.5
 226. (a) $\sqrt{\frac{1}{n} \sum (x_i - \bar{x})^2}$
 227. (b) $\sqrt{\frac{(n^2-1)}{12}}$
 228. (c) 6
 229. (a) 2.87
 230. (d) Coefficient of variation
 231. (d) Quartile deviation
 232. (b) 2.5
 233. (d) 7.7%
 234. (a) i and ii
 235. (b) i and iii
 236. (d) To uniquely identify a distribution
 237. (b) Raw moments
 238. (c) (-3, 3)
 239. (d) Rectified Moments
 240. (a) $\frac{\sum (x_i - \bar{x})^n}{w}$
 241. (b) $\mu'_1 = \bar{x} - a$
 242. (a) $\frac{\sum f_i (x_i - a)^r}{n}$
 243. (c) Moments
 243. (d) i, ii and iii
 244. (c) μ_2
 245. (c) $\frac{\sqrt{\mu_2}}{\bar{x}} \times 100$
 246. (d) Arithmetic Mean
247. (c) First central moment
 248. (b) μ_3
 249. (d) $\mu'_2 - \mu_1'^2$
 250. (b) 0
 251. (d) $\bar{x} - a$
 252. (b) -2
 253. (c) 0
 254. (a) 2
 255. (a) 2
 256. (a) 10
 257. (b) -3.4
 258. (a) 0
 259. (a) 3.2
 260. (c) 5
 261. (b) 20
 262. (d) 21
 263. (c) 60
 264. (b) i and iii
 265. (b) 11.92
 266. (d) 110
 267. (b) Symmetrical
 268. (a)
 269. (a) Positive Skew
 270. (a) 0
 271. (b) Negative Skew
272. (a) Positive Skew
 273. (d) All of the above
 274. (b) Median
 275. (c) Mode
 276. (a) Mean
 277. (b) Positively skewed
 278. (c) $M_o = 3Me - 2\bar{x}$
 280. (b) leptokurtic
 281. (d) 29.45
 282. (a) $Mean > Median > Mode$
 283. (c) 0
 284. (a) Left Skew
 285. (c) Right Skew
 286. (b) Symmetry
 287. (c) 3
 288. (c) 8.25
 289. (c) Symmetric
 290. (a) Positively skewed
 291. (b) Negatively skewed
 292. (a) i and ii
 293. (c) ii and iii
 294. (b) i and iii
 295. (a) $\gamma_1 = \sqrt{\frac{\mu_3^2}{\mu_2^3}}$
 296. (b) 20
 297. (d) 58
 298. (a) 52.67
 299. (a) 15

300. (c) 26
301. (d) i, ii and iii
302. (a) A
303. (b) 3
304. (d) 48
305. (c) Mesokurtic
306. (a) Leptokurtic
307. (c) Mesokurtic
308. (b) Platykurtic
309. (b) $\gamma_2 = \beta_2 - 3$
310. (c) 3
311. (d) $\gamma_2 = \beta_2 - 3$
312. (a) $Q_1 - 1.5 \times IQR$
313. (b) $IQR = Q_3 - Q_1$
314. (a) Mode
315. (d) i, ii & iii
316. (b) 50
317. (a) i and ii
318. (b) i and iii
319. (d) i, ii and iii
320. (a) Arithmetic Mean
321. (d) Karl Pearson
322. (a) -0.92
323. (d) -1
324. (a) Correlation
325. (c) Correlation coefficient
326. (a) Qualitative variable
327. (d) 0
328. (c) Perfect positive correlation
329. (a) Perfect negative correlation
330. (b) Partial positive correlation
331. (d) Partial negative correlation
332. (d) Zero correlation
333. (a) Choice
334. (d) -0.94
335. (d) -0.94
336. (c) Cannot be determined
337. (b) 0.60
338. (c) 0.67
339. (c) No. of earthquakes
340. (c) Number of students
341. (a) Number of calls received
342. (a) i and ii
343. (c) Moving average method
344. (a) $Y_t = T_t \times S_t \times C_t \times I_t$
345. (a) 0
346. (b) 1
347. (d) Seasonal variation
348. (b) 190
349. (a) a curved line
350. (a) Increase in ice cream sales
351. (b) A toy store
352. (a) Boom and recession phases
353. (a) Gradual increase in global average temperature
354. (a) Upward trend
355. (a) Declining birth rate
356. (a) Improved marketing strategies
357. (b) Long-term business cycles
358. (d) Any of the above
359. (b) Graphical method
360. (b) 90.37
361. (a) Upward
362. (c) Irregular Variation
363. (c) 25.88
364. (a) Upward
365. (b) General Trend
366. (b) 90.37
367. (a) Upward
368. (c) Irregular Variation
369. (b) 165
370. (b) Fluctuations
371. (a) Seasonal Variation
372. (b) 41.5°C
373. (b) Cyclic Variation
374. (d) Regular Variation
375. (b) General Trend
376. (d) Moving Median
377. (c) 95.33
378. (c) 95.33
379. (c) Irregular Variation
380. (a) Trend
381. (b) Seasonal Variation
382. (c) Irregular Variation
383. (b) Seasonal Variation
384. (b) 3
385. (a) 1350
386. (c) 1350
387. (a) 1350
388. (b) Seasonal Variation
389. (b) 2013
390. (c) Irregular Variation
391. (a) 2
392. (d) Any of the above
393. (c) 6
394. (b) 6
395. (c) 10.65
396. (b) 3.5%
397. (b) 3.1%
398. (c) 3.8%
399. (b) 92 cm
400. (a) 86.5 cm
401. (b) 89 cm
402. (c) 21.5°C
403. (b) 18°C
404. (c) 26°C
405. (c) Jul
406. (a) 52.5
407. (b) 55
408. (a) i and ii
409. (c) i
410. (a) i and ii
411. (d) i, ii and iii
412. (b) 3
413. (a) Official statistics
414. (c) Semi-official statistics
415. (b) BBS
416. (b) Non-official statistics
417. (b) World Health Organization (WHO)
418. (c) 10
419. (a) Official statistics
420. (c) Finance