


2.0 nm film: registered value on the original measured curve
6.3 nm film: registered value on the original measured curve
13.2 nm film: registered value on the original measured curve
(a) shaded: replicate 95 % interval (2.5-97.5 %)

(b) SS fixed window 5e-11...5e-10 A/µm (floor rule); (c) V_{th} constant current 1e-9 A/µm; (d) floor; bars = 95 % interval
(b) SS A1 window [r, 10 r] x I_{on} (fixed); (c) V_{th} constant current 5e-10 A/µm; bars = 95 % interval
(c) V_{th} extrapolation in the linear region (ELR); bars = 95 % interval
(c) power-law V_T (fit 1-3 V, first 400 replicates); bars = 95 % interval

Data: C:/Users/moham/Downloads/IWO_Local_Codex_Handoff (1)/IWO_PHYSICS_CONSTRAINED_MODEL_V1/analysis_2026-10-10_campaign/mc_results_2026-10-15/mc_measured_extraction_out.json, produced 2026-10-15T22:11:55Z by mc_measured_extraction.py (sha256 8c9c4aff; output sha256 2c354272), register A57 / D-80. Measured input: C:/Users/moham/Downloads/IWO_Local_Codex_Handoff (1)/IWO_PHYSICS_CONSTRAINED_MODEL_V1/data/experimental_clean.csv (sha256 e27a53e8; MEASURED, already seen: extraction uncertainty, not a blind test). Noise model ASSUMED (SMU specification not supplied); N = 2000 replicates per film (power law on 400), seed 20261016 + film index.