

Class19

Shayan Malekpour

Investigating pertussis cases by year

Q1. With the help of the R “addin” package `datapasta` assign the CDC pertussis case number data to a data frame called `cdc` and use `ggplot` to make a plot of cases numbers over time.


```
library(datapasta)
library(ggplot2)

cdc <- data.frame(
  Year = c(1922L, 1923L, 1924L, 1925L,
           1926L, 1927L, 1928L, 1929L, 1930L, 1931L,
           1932L, 1933L, 1934L, 1935L, 1936L,
           1937L, 1938L, 1939L, 1940L, 1941L, 1942L,
           1943L, 1944L, 1945L, 1946L, 1947L,
           1948L, 1949L, 1950L, 1951L, 1952L,
           1953L, 1954L, 1955L, 1956L, 1957L, 1958L,
           1959L, 1960L, 1961L, 1962L, 1963L,
           1964L, 1965L, 1966L, 1967L, 1968L, 1969L,
           1970L, 1971L, 1972L, 1973L, 1974L,
           1975L, 1976L, 1977L, 1978L, 1979L, 1980L,
           1981L, 1982L, 1983L, 1984L, 1985L,
           1986L, 1987L, 1988L, 1989L, 1990L,
           1991L, 1992L, 1993L, 1994L, 1995L, 1996L,
           1997L, 1998L, 1999L, 2000L, 2001L,
           2002L, 2003L, 2004L, 2005L, 2006L, 2007L,
           2008L, 2009L, 2010L, 2011L, 2012L,
           2013L, 2014L, 2015L, 2016L, 2017L, 2018L,
           2019L, 2020L, 2021L),
  No..Reported.Pertussis.Cases = c(107473, 164191, 165418, 152003,
                                   202210, 181411, 161799, 197371,
                                   166914, 172559, 215343, 179135, 265269,
```

```
180518,147237,214652,227319,103188,  
183866,222202,191383,191890,109873,  
133792,109860,156517,74715,69479,  
120718,68687,45030,37129,60886,  
62786,31732,28295,32148,40005,  
14809,11468,17749,17135,13005,6799,  
7717,9718,4810,3285,4249,3036,  
3287,1759,2402,1738,1010,2177,2063,  
1623,1730,1248,1895,2463,2276,  
3589,4195,2823,3450,4157,4570,  
2719,4083,6586,4617,5137,7796,6564,  
7405,7298,7867,7580,9771,11647,  
25827,25616,15632,10454,13278,  
16858,27550,18719,48277,28639,32971,  
20762,17972,18975,15609,18617,  
6124,2116)
```

```
)
```

```
ggplot(cdc) +  
  aes(x = Year, y = No..Reported.Pertussis.Cases) +  
  geom_point() +  
  geom_line() +  
  labs(x = "Year", y = "Cases", title = "Pertussis Cases Over Time")
```




A tale of two vaccines (wP & aP)

Q2. Using the ggplot `geom_vline()` function add lines to your previous plot for the 1946 introduction of the wP vaccine and the 1996 switch to aP vaccine (see example in the hint below). What do you notice?

```
ggplot(cdc) +
  aes(x = Year, y = No..Reported.Pertussis.Cases) +
  geom_point() +
  geom_line() +
  geom_vline(xintercept = 1946, linetype = "dashed", color = "red") +
  geom_vline(xintercept = 1996, linetype = "dashed", color = "blue") +

  labs(x = "Year", y = "Cases", title = "Pertussis Cases Over Time")
```



Q3. Describe what happened after the introduction of the aP vaccine? Do you have a possible explanation for the observed trend? cases decreased

Exploring CMI-PB data

The CMI-PB API returns JSON data

```
library(jsonlite)
```

```
subject <- read_json("https://www.cmi-pb.org/api/subject", simplifyVector = TRUE)
```

```
head(subject, 3)
```

	subject_id	infancy_vac	biological_sex	ethnicity	race
1	1	wP	Female Not	Hispanic or Latino	White
2	2	wP	Female Not	Hispanic or Latino	White
3	3	wP	Female	Unknown	White

	year_of_birth	date_of_boost	dataset
1	1986-01-01	2016-09-12	2020_dataset
2	1968-01-01	2019-01-28	2020_dataset

3 1983-01-01 2016-10-10 2020_dataset

```
# Q4. How many aP and wP infancy vaccinated subjects are in the dataset?  
table(subject$infancy_vac)
```

```
aP wP  
47 49
```

```
# Q5. How many Male and Female subjects/patients are in the dataset?  
table(subject$biological_sex)
```

```
Female    Male  
     66     30
```

```
# Q6. What is the breakdown of race and biological sex (e.g.number of Asian females, White
```

```
table(subject$biological_sex, subject$race)
```

	American Indian/Alaska Native	Asian	Black or African American
Female	0	18	2
Male	1	9	0

	More Than One Race	Native Hawaiian or Other Pacific Islander
Female	8	1
Male	2	1

	Unknown or Not Reported	White
Female	10	27
Male	4	13

```
## Working with Dates
library(lubridate)
```

Attaching package: 'lubridate'

The following objects are masked from 'package:base':

```
date, intersect, setdiff, union
```

```
time_length( today() - ymd("2000-01-01"), "years")
```

```
[1] 23.45243
```

Q7. Using this approach determine (i) the average age of wP individuals, (ii) the average age of aP individuals;

```
subject$age <- today () - ymd(subject$year_of_birth)
```

```
library(dplyr)
```

Attaching package: 'dplyr'

The following objects are masked from 'package:stats':

```
filter, lag
```

The following objects are masked from 'package:base':

```
intersect, setdiff, setequal, union
```

```
ap <- subject %>% filter(infancy_vac == "aP")
round( summary( time_length( ap$age, "years" ) ) )
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
23	25	26	26	26	27

```
wp <- subject %>% filter(infancy_vac == "wP")
round( summary( time_length( wp$age, "years" ) ) )
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
28	32	35	37	40	55

```
# Q8. Determine the age of all individuals at time of boost?
int <- ymd(subject$date_of_boost) - ymd(subject$year_of_birth)
age_at_boost <- time_length(int, "year")
head(age_at_boost)
```

```
[1] 30.69678 51.07461 33.77413 28.65982 25.65914 28.77481
```


Q9. With the help of a faceted boxplot or histogram (see below), do you think these two groups are significantly different? not significant difference.

```
# Or use wilcox.test()
x <- t.test(time_length( wp$age, "years" ),
            time_length( ap$age, "years" ))
x$p.value
```

```
[1] 1.316045e-16
```

```
ggplot(subject) +
  aes(time_length(age, "year"),
      fill=as.factor(infancy_vac)) +
  geom_histogram(show.legend=FALSE) +
  facet_wrap(vars(infancy_vac), nrow=2) +
  xlab("Age in years")
```

`stat_bin()` using `bins = 30`. Pick better value with `binwidth`.



Joining multiple tables

```
# Complete the API URLs...
specimen <- read_json("https://www.cmi-pb.org/api/specimen", simplifyVector = TRUE)
titer <- read_json("https://www.cmi-pb.org/api/ab_titer", simplifyVector = TRUE)
```

Q10. Complete the code to join `specimen` and `subject` tables to make a new merged data frame containing all specimen records along with their associated subject details:

```
meta <- inner_join(specimen, subject)
```

Joining with ``by = join_by(subject_id)``

```
dim(meta)
```

```
[1] 729 14
```

```
head(meta)
```


	specimen_id	subject_id	actual_day_relative_to_boost			
1	1	1	-3			
2	2	1	736			
3	3	1	1			
4	4	1	3			
5	5	1	7			
6	6	1	11			

	planned_day_relative_to_boost	specimen_type	visit	infancy_vac	biological_sex
1	0	Blood	1	wP	Female
2	736	Blood	10	wP	Female
3	1	Blood	2	wP	Female
4	3	Blood	3	wP	Female
5	7	Blood	4	wP	Female
6	14	Blood	5	wP	Female

	ethnicity	race	year_of_birth	date_of_boost	dataset
1	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
2	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
3	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
4	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
5	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
6	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset

	age
1	13679 days
2	13679 days
3	13679 days
4	13679 days
5	13679 days
6	13679 days

Q11. Now using the same procedure join `meta` with `titer` data so we can further analyze this data in terms of time of visit aP/wP, male/female etc.

```
abdata <- inner_join(titer, meta)
```

Joining with `by = join_by(specimen_id)`

```
dim(abdata)
```

```
[1] 32675    21
```

Q12. How many specimens (i.e. entries in `abdata`) do we have for each `isotype`?

```
table(abdata$isotype)
```

```

IgE  IgG  IgG1 IgG2 IgG3 IgG4
6698 1413 6141 6141 6141 6141

```

Q13. What do you notice about the number of visit 8 specimens compared to other visits?
lower than the other visits

```
table(abdata$visit)
```

```

 1    2    3    4    5    6    7    8
5795 4640 4640 4640 4640 4320 3920  80

```

Examine IgG1 Ab titer levels

```

ig1 <- abdata %>% filter(isotype == "IgG1", visit!=8)
head(ig1)

```

	specimen_id	isotype	is_antigen_specific	antigen	MFI	MFI_normalised
1	1	IgG1	TRUE	ACT	274.355068	0.6928058
2	1	IgG1	TRUE	LOS	10.974026	2.1645083
3	1	IgG1	TRUE	FELD1	1.448796	0.8080941
4	1	IgG1	TRUE	BETV1	0.100000	1.0000000
5	1	IgG1	TRUE	LOLP1	0.100000	1.0000000
6	1	IgG1	TRUE	Measles	36.277417	1.6638332

	unit	lower_limit_of_detection	subject_id	actual_day_relative_to_boost
1	IU/ML	3.848750	1	-3
2	IU/ML	4.357917	1	-3
3	IU/ML	2.699944	1	-3
4	IU/ML	1.734784	1	-3
5	IU/ML	2.550606	1	-3
6	IU/ML	4.438966	1	-3

	planned_day_relative_to_boost	specimen_type	visit	infancy_vac	biological_sex
1	0	Blood	1	wP	Female
2	0	Blood	1	wP	Female
3	0	Blood	1	wP	Female


4		0	Blood	1	wP	Female
5		0	Blood	1	wP	Female
6		0	Blood	1	wP	Female

	ethnicity	race	year_of_birth	date_of_boost	dataset
1	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
2	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
3	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
4	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
5	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset
6	Not Hispanic or Latino	White	1986-01-01	2016-09-12	2020_dataset

	age
1	13679 days
2	13679 days
3	13679 days
4	13679 days
5	13679 days
6	13679 days


Q14. Complete the following code to make a summary boxplot of Ab titer levels (MFI) for all antigens:

```
ggplot(ig1) +
  aes(x = MFI, y = antigen) +
  geom_boxplot() +
  facet_wrap(vars(visit), nrow = 2)
```




Q15. What antigens show differences in the level of IgG1 antibody titers recognizing them over time? Why these and not others? antigen 5 has significant difference and FIM2/3 are different.

```
## Better visualization
ggplot(ig1) +
  aes(MFI, antigen, col=infancy_vac ) +
  geom_boxplot(show.legend = FALSE) +
  facet_wrap(vars(visit), nrow=2) +
  theme_bw()
```




Another version :

```
ggplot(ig1) +
  aes(MFI, antigen, col=infancy_vac ) +
  geom_boxplot(show.legend = FALSE) +
  facet_wrap(vars(infancy_vac, visit), nrow=2)
```




Q16. Filter to pull out only two specific antigens for analysis and create a boxplot for each. You can chose any you like. Below I picked a “control” antigen (“**Measles**”, that is not in our vaccines) and a clear antigen of interest (“**FIM2/3**”, extra-cellular fimbriae proteins from *B. pertussis* that participate in substrate attachment).

```
filter(ig1, antigen=="Measles") %>%
  ggplot() +
  aes(MFI, col=infancy_vac) +
  geom_boxplot(show.legend = F ) +
  facet_wrap(vars(visit)) +
  theme_bw()
```



```
## and the same for antigen=="FIM2/3"
filter(ig1, antigen=="FIM2/3") %>%
  ggplot() +
  aes(MFI, col=infancy_vac) +
  geom_boxplot(show.legend = F) +
  facet_wrap(vars(visit)) +
  theme_bw()
```



Q17. What do you notice about these two antigens time courses and the FIM2/3 data in particular?

FIM2/3 has longer time course compared to Measles

Q18. Do you see any clear difference in aP vs. wP responses?

No clear difference

Obtaining CMI-PB RNASeq data

```
url <- "https://www.cmi-pb.org/api/v2/rnaseq?versioned_ensembl_gene_id=eq.ENSOG00000211896."


rna <- read_json(url, simplifyVector = TRUE)

#meta <- inner_join(specimen, subject)
ssrna <- inner_join(rna, meta)
```

Joining with `by = join_by(specimen_id)`

Q19. Make a plot of the time course of gene expression for IGHG1 gene (i.e. a plot of visit vs. tpm).

```
ggplot(ssrna) +  
  aes(visit, tpm, group=subject_id) +  
  geom_point() +  
  geom_line(alpha=0.2)
```




Q20.: What do you notice about the expression of this gene (i.e. when is it at it's maximum level)?

the expression is at maximum in 4th visit.

Q21. Does this pattern in time match the trend of antibody titer data?

yes it matches the plot.

```
ggplot(ssrna) +  
  aes(tpm, col=infancy_vac) +  
  geom_boxplot() +  
  facet_wrap(vars(visit))
```



```
## no obvious wP vs. aP differences here even if we focus in on a particular visit:
ssrna %>%
  filter(visit==4) %>%
  ggplot() +
    aes(tpm, col=infancy_vac) + geom_density() +
    geom_rug()
```

