

DISCOVERY THROUGH GOSSIP

MOHAMMAD MAHZOUN

February 19, 2020

CONTENTS

1	Introduction	2
1.1	Notation	2
1.2	Useful lemmas	2
2	Triangulation Results	3
2.1	Upper Bound	3
2.2	Lower Bound	3
2.3	Paragraphs	3
2.4	Math	4
3	Results and Discussion	4
3.1	Subsection	5
3.2	Figure Composed of Subfigures	5

LIST OF FIGURES

Figure 1	Discovery Methods	2
Figure 2	An example of a floating figure	5
Figure 3	A number of pictures.	6

LIST OF TABLES

Table 1	Table of Notations	2
---------	------------------------------	---




Figure 1: Discovery Methods

1 INTRODUCTION

We want to study Gossip based discovery processes in both directed and undirected graphs using push discovery (triangulation) and pull discovery (two-hop walk process).

We are interested in studying the time taken by process to converge to the transitive closure of the graph.

1.1 Notation

Table 1: Table of Notations

Notation	Description
δ_t	Minimum degree of G_t
$N_t^i(u)$	Set of nodes at distance i from u in G_t
$d_t(u)$	Degree of u in G_t
$d_t(u, S)$	Degree induced on S

1.2 Useful lemmas

Lemma 1. $|\cup_{i=1}^4 N_t^i(u)| \geq \min\{2\delta_t, n-1\}$ for all $u \in G_t$.

Proof. If $N_t^3 \neq \emptyset$, then $|\cup_{i=2}^4 N_t^i(u)| \geq \delta_t$ and $|N_t^1(u)| \geq \delta_t$. So $|\cup_{i=1}^4 N_t^i(u)| \geq 2\delta_t$ since the two sets are disjoint.

If $N_t^3 = \emptyset$, $N_t^1(u) \cup N_t^2(u) = n-1$ since G_t is connected. □

Lemma 2. Consider k Bernoulli experiments in which the success probability of the i th experiment is at least i/m where $m \geq k$. If X_i denotes the number of trials needed for experiment i to output a success and $X = \sum_{i=1}^k X_i$, then $\Pr[X > (c+1)m \ln m] < \frac{1}{m^c}$.

Proof. w.l.o.g assume that $k = m$. The problem can be seen as *coupon collector problem* where X_{m+i-1} is the number of steps to collect i th coupon. Consider the probability of not obtaining the i th coupon after $(c+1)m \ln m$ steps, we have: $(1 - \frac{1}{m})^{(c+1)m \ln m} < e^{-(c+1) \ln m} = \frac{1}{m^{c+1}}$. By union bound, the probability that some coupon has not been collected after $(c+1)m \ln m$ steps is less than $\frac{1}{m^c}$. □

2 TRIANGULATION RESULTS

2.1 Upper Bound

2.2 Lower Bound

Theorem 3 (Upper bound for triangulation process). *For any connected undirected graph, the triangulation process converges to a complete graph in $O(n \log^2 n)$ rounds with high probability.*

In order to prove Theorem 3, we prove that the minimum degree of the graph increases by a constant factor (or equals to $n - 1$) in $O(n \log n)$ steps. We say that a node v is **weakly tied** to a set of nodes S if $d_t(v, S) < \delta_0/2$, and **strongly tied** to a set of nodes S if $d_t(v, S) \geq \delta_0/2$.

Lemma 4. If $d_t(u) < \min\{n - 1, (1 + \frac{1}{4}\delta_0)\}$ and $w \in N_t^1(u)$ has at least $\frac{\delta_0}{4}$ edges to $N_t^2(u)$, then the probability that u connects to a node in $N_t^2(u)$ through w in round t is at least $\frac{1}{6n}$.

Proof. The probability that u connects to a node in $N_t^2(u)$ through w in round t is:

$$\begin{aligned} \frac{d_t(w, N_t^2(u))}{D_t(w)} \times \frac{1}{d_t(w)} &\geq \frac{d_t(w, N_t^2(u))}{D_t(w)} \times \frac{1}{n} \geq \frac{d_t(w, N_t^2(u))}{|N_t^1(u)| + d_t(w, N_t^2(u))} \times \frac{1}{n} \geq \\ &\frac{d_t(w, N_t^2(u))}{(1 + \frac{1}{4})\delta_0 + d_t(w, N_t^2(u))} \times \frac{1}{n} \geq \frac{\frac{\delta_0}{4}}{(1 + \frac{1}{4})\delta_0 + \frac{\delta_0}{4}} \times \frac{1}{n} = \frac{1}{6n} \end{aligned}$$

□

Lemma 5. If $d_t(u) < \min\{n - 1, (1 + \frac{1}{4}\delta_0)\}$ and $w \in N_t^1(u)$ is weakly tied to $N_t^2(u)$, and $v \in N_0^2(u) \cap N_0^1(w)$, then u connects to v through w in round t with probability at least $\frac{1}{4\delta_0^2}$.

Proof. Since w is weakly tied to $N_t^2(u)$ and $d_t(w)$ is at most $|N_t^1(u)| + d_t(w, N_t^2(u))$, we obtain that $d_t(w)$ is at most $(1 + \frac{1}{4})\delta_0 + \frac{\delta_0}{2}$. Therefore, the probability that u connects to v through w in round t equals:

$$\frac{1}{d_t(w)^2} \geq \frac{1}{((1 + \frac{1}{4})\delta_0 + \frac{\delta_0}{2})^2} \geq \frac{1}{\frac{7\delta_0}{4}} \geq \frac{1}{4\delta_0^2}.$$

□

To analyze the growth in the degree of a node u , we consider two overlapping cases. The first case is when more than $\delta_0/4$ nodes of $N_t^1(u)$ are strongly tied to $N_t^2(u)$, and the second is when less than $\delta_0/3$ nodes of $N_t^1(u)$ are strongly tied to $N_t^2(u)$.

Lemma 6 (Several nodes are strongly tied to two-hop neighbors). There exists a $T = O(n \log n)$ such that if more than

2.3 Paragraphs

Suspendisse vel felis. Ut lorem lorem, interdum eu, tincidunt sit amet, laoreet vitae, arcu. Aenean faucibus pede eu ante. Praesent enim elit, rutrum at, molestie non, nonummy vel, nisl. Ut lectus eros, malesuada sit amet, fermentum eu, sodales cursus, magna. Donec eu purus. Quisque vehicula, urna sed ultricies auctor, pede lorem egestas dui, et convallis elit erat sed nulla. Donec luctus. Curabitur et nunc. Aliquam dolor odio, commodo pretium, ultricies non, pharetra in, velit. Integer arcu est, nonummy in, fermentum faucibus, egestas vel, odio.

PARAGRAPH DESCRIPTION Sed commodo posuere pede. Mauris ut est. Ut quis purus. Sed ac odio. Sed vehicula hendrerit sem. Duis non odio. Morbi ut dui. Sed accumsan risus eget odio. In hac habitasse platea dictumst. Pellentesque non elit. Fusce sed justo eu urna porta tincidunt. Mauris felis odio, sollicitudin sed, volutpat a, ornare ac, erat. Morbi quis dolor. Donec pellentesque, erat ac sagittis semper, nunc dui lobortis purus, quis congue purus metus ultricies tellus. Proin et quam. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Praesent sapien turpis, fermentum vel, eleifend faucibus, vehicula eu, lacus.

DIFFERENT PARAGRAPH DESCRIPTION Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Donec odio elit, dictum in, hendrerit sit amet, egestas sed, leo. Praesent feugiat sapien aliquet odio. Integer vitae justo. Aliquam vestibulum fringilla lorem. Sed neque lectus, consectetur at, consectetur sed, eleifend ac, lectus. Nulla facilisi. Pellentesque eget lectus. Proin eu metus. Sed porttitor. In hac habitasse platea dictumst. Suspendisse eu lectus. Ut mi mi, lacinia sit amet, placerat et, mollis vitae, dui. Sed ante tellus, tristique ut, iaculis eu, malesuada ac, dui. Mauris nibh leo, facilisis non, adipiscing quis, ultrices a, dui.

2.4 Math

Quisque ullamcorper placerat ipsum. Cras nibh. Morbi vel justo vitae lacus tincidunt ultrices. Lorem ipsum dolor sit amet, consectetur adipiscing elit. In hac habitasse platea dictumst. Integer tempus convallis augue. Etiam facilisis. Nunc elementum fermentum wisi. Aenean placerat. Ut imperdiet, enim sed gravida sollicitudin, felis odio placerat quam, ac pulvinar elit purus eget enim. Nunc vitae tortor. Proin tempus nibh sit amet nisl. Vivamus quis tortor vitae risus porta vehicula.

$$\cos^3 \theta = \frac{1}{4} \cos \theta + \frac{3}{4} \cos 3\theta \quad (1)$$

Fusce mauris. Vestibulum luctus nibh at lectus. Sed bibendum, nulla a faucibus semper, leo velit ultricies tellus, ac venenatis arcu wisi vel nisl. Vestibulum diam. Aliquam pellentesque, augue quis sagittis posuere, turpis lacus congue quam, in hendrerit risus eros eget felis. Maecenas eget erat in sapien mattis porttitor. Vestibulum porttitor. Nulla facilisi. Sed a turpis eu lacus commodo facilisis. Morbi fringilla, wisi in dignissim interdum, justo lectus sagittis dui, et vehicula libero dui cursus dui. Mauris tempor ligula sed lacus. Duis cursus enim ut augue. Cras ac magna. Cras nulla. Nulla egestas. Curabitur a leo. Quisque egestas wisi eget nunc. Nam feugiat lacus vel est. Curabitur consectetur.

Definition 1 (Gauss). To a mathematician it is obvious that $\int_{-\infty}^{+\infty} e^{-x^2} dx = \sqrt{\pi}$.

Theorem 7 (Pythagoras). *The square of the hypotenuse (the side opposite the right angle) is equal to the sum of the squares of the other two sides.*

Proof. We have that $\log(1)^2 = 2 \log(1)$. But we also have that $\log(-1)^2 = \log(1) = 0$. Then $2 \log(-1) = 0$, from which the proof. \square

3 RESULTS AND DISCUSSION

Reference to Figure 2 on the following page.

Suspendisse vitae elit. Aliquam arcu neque, ornare in, ullamcorper quis, commodo eu, libero. Fusce sagittis erat at erat tristique mollis. Maecenas sapien libero, molestie et, lobortis in, sodales eget, dui. Morbi ultrices rutrum lorem. Nam elementum ullamcorper leo. Morbi dui. Aliquam sagittis. Nunc placerat. Pellentesque




Figure 2: An example of a floating figure (a reproduction from the *Gallery of prints*, M. Escher, from <http://www.mcescher.com/>).

tristique sodales est. Maecenas imperdiet lacinia velit. Cras non urna. Morbi eros pede, suscipit ac, varius vel, egestas non, eros. Praesent malesuada, diam id pretium elementum, eros sem dictum tortor, vel consectetur odio sem sed wisi.

3.1 Subsection

3.1.1 Subsubsection

WORD Definition

CONCEPT Explanation

IDEA Text

- First item in a list
- Second item in a list
- Third item in a list

Reference to Table 1 on page 2.

3.2 Figure Composed of Subfigures

Reference the figure composed of multiple subfigures as Figure 3 on the following page. Reference one of the subfigures as Figure 3b on the next page.

Nulla in ipsum. Praesent eros nulla, congue vitae, euismod ut, commodo a, wisi. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Aenean nonummy magna non leo. Sed felis erat, ullamcorper in, dictum non, ultricies ut, lectus. Proin vel arcu a odio lobortis euismod. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Proin ut est. Aliquam odio. Pellentesque massa turpis, cursus eu, euismod nec, tempor congue, nulla. Duis viverra gravida mauris. Cras tincidunt. Curabitur eros ligula, varius ut, pulvinar in, cursus faucibus, augue.

Nulla mattis luctus nulla. Duis commodo velit at leo. Aliquam vulputate magna et leo. Nam vestibulum ullamcorper leo. Vestibulum condimentum rutrum mauris. Donec id mauris. Morbi molestie justo et pede. Vivamus eget turpis sed nisl cursus tempor. Curabitur mollis sapien condimentum nunc. In wisi nisl, malesuada at, dignissim sit amet, lobortis in, odio. Aenean consequat arcu a ante. Pellentesque porta elit sit amet orci. Etiam at turpis nec elit ultricies imperdiet. Nulla facilisi.




Figure 3: A number of pictures with no common theme.

In hac habitasse platea dictumst. Suspendisse viverra aliquam risus. Nullam pede justo, molestie nonummy, scelerisque eu, facilisis vel, arcu.

Curabitur tellus magna, porttitor a, commodo a, commodo in, tortor. Donec interdum. Praesent scelerisque. Maecenas posuere sodales odio. Vivamus metus lacus, varius quis, imperdiet quis, rhoncus a, turpis. Etiam ligula arcu, elementum a, venenatis quis, sollicitudin sed, metus. Donec nunc pede, tincidunt in, venenatis vitae, faucibus vel, nibh. Pellentesque wisi. Nullam malesuada. Morbi ut tellus ut pede tincidunt porta. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam congue neque id dolor.

Donec et nisl at wisi luctus bibendum. Nam interdum tellus ac libero. Sed sem justo, laoreet vitae, fringilla at, adipiscing ut, nibh. Maecenas non sem quis tortor eleifend fermentum. Etiam id tortor ac mauris porta vulputate. Integer porta neque vitae massa. Maecenas tempus libero a libero posuere dictum. Vestibulum ante ipsum primis in faucibus orci luctus et ultrices posuere cubilia Curae; Aenean quis mauris sed elit commodo placerat. Class aptent taciti sociosqu ad litora torquent per conubia nostra, per inceptos hymenaeos. Vivamus rhoncus tincidunt libero. Etiam elementum pretium justo. Vivamus est. Morbi a tellus eget pede tristique commodo. Nulla nisl. Vestibulum sed nisl eu sapien cursus rutrum.

REFERENCES

- [1] A. J. Figueredo and P. S. A. Wolf. Assortative pairing and life history strategy - a cross-cultural study. *Human Nature*, 20:317–330, 2009.